Rapid Point Survey by Bayonet Blade¹

PAULE S. POISSONET, PHILIPPE M. DAGET, JACQUES A. POISSONET, AND GILBERT A. LONG²

Plant Ecologists, Centre d'Etudes Phytosociologiques et Ecologiques, Centre National de la Recherche Scientifique, Montpellier, France.

A bayonet was developed for rapid points in tall dense herbaceous vegetation (Fig. 1 and 2). It is a steel blade about 65 cm high, 4.5 cm broad and 2 mm thick. This blade is sharp-edged along one height which constitutes the line of sight. The upper part is T shaped and used as a plumb-line. The

reduces the disturbance of vegetation when the bayonet blade is vertically stuck into the ground. At each species, all contacts of any part of the vegetation of every species, with the sharp-

lower part is bevelled on the opposite

side to the sharp-edged height; which

edged part of the blade, are recorded. A trained observer can do 350 sightings a day. This apparatus has been used successfully in tall grassland vegetation of France. Data obtained by




Fig. 1. The bayonet.



Fig. 2. The bayonet in field use.

bayonet recording give very small deviation (Poissonet, 1969) as compared to a needle point frame (Long et al., 1972).

Literature Cited

Long, G. A., P. S. Poissonet, J. A. Poissonet, Ph. M. Daget, and M. P. Godron. 1972. Improved needle point frame for exact line transects. J. Range Manage. 25:228–229.

Poissonet, P., and J. Poissonet. 1969. Etude comparée de diverses méthodes d'analyse de la végétation des formations herbacées denses et permanentes. Conséquences pour les applications agronomiques. C.E.P.E.—C.N.R.S., Montpellier, France. 50: 1–120.

SRM Annual Meetings

1973—Boise, Idaho Rodeway Inn February 4–9 1974-Tucson, Arizona Downtowner Motor Inn February 3-8

¹ Received November 30, 1971.

² Authors acknowledge with thanks the guidance provided by Professor Naphtali Tadmor, from the department of Botany, Hebrew University of Jerusalem, in editing the paper.