

Ungulate herbivory on Utah aspen: Assessment of long-term exclosures

CHARLES E. KAY AND DALE L. BARTOS

Authors are Natural Resource Policy Associate, Institute of Political Economy and Adjunct Assistant Professor, Department of Political Science, Utah State University, Logan, Utah 84322; and Ecologist, USDA Forest Service, Rocky Mountain Research Station, Logan, Utah 84321.

Abstract

The role of livestock grazing and big-game browsing in the decline of aspen (*Populus tremuloides* Michx.) in the Intermountain West has long been questioned. All known aspen exclosures (n=8) on the Dixie and Fishlake National Forests in south-central Utah were measured during late summer of 1995 and 1996 to determine aspen stem dynamics, successional status, and understory species composition. Five of the exclosures were of a 3-part design with a total-exclusion portion, a livestock-exclusion portion, and a combined-use portion which permitted the effects of deer (*Odocoileus hemionus*) and elk (*Cervus elaphus*) herbivory to be measured separately from those of livestock. Aspen within all total-exclusion plots successfully regenerated and developed multi-aged stems without the influence of fire or other disturbance. Aspen subjected to browsing by wildlife, primarily mule deer, either failed to regenerate successfully or regenerated at stem densities significantly lower (2,498 stems ha⁻¹) than that on total-exclusion plots (4,474 stems ha⁻¹). On combined wildlife-livestock-use plots, most aspen failed to regenerate successfully, or did so at low stem densities (1,012 stems/ha⁻¹). Aspen successfully regenerated on ungulate-use plots only when deer numbers were low. Similarly, ungulate herbivory had significant effects on understory species composition. In general, utilization by deer tended to reduce shrubs and tall palatable forbs while favoring the growth of native grasses. The addition of livestock grazing, however, tended to reduce native grasses while promoting introduced species and bare soil. Thus, communities dominated by old-age or single-age trees appear to be a product of ungulate browsing, not a biological attribute of aspen as has been commonly assumed. There was no evidence that climatic variation affected aspen regeneration. Observed differences are attributed to varied histories of ungulate herbivory.

Key Words: *Populus tremuloides* communities, reproduction, decline, cattle grazing, deer browsing, elk browsing, undergrowth.

Aspen (*Populus tremuloides* Michx.) in the western United States does not commonly grow from seed because of its demanding seed bed requirements (Kay and White 1995,

Research was funded by U.S. Forest Service's Intermountain Research Station (now part of the Rocky Mountain Research Station) and we thank the many agency employees who helped with various aspects of this project, especially Robert B. Campbell, Jr. and Dr. Walter F. Mueggler (Scientist Emeritus). Also, Brad Williams and the Utah State Trust Lands Administration for granting access to their files on the Parker Mountain exclosure.

Manuscript accepted 28 June 1999.

Resumen

El papel del apacentamiento del ganado y el ramoneo de la fauna silvestre mayor en la disminución del "Aspen" (*Populus tremuloides* Michx.) en la región intermontañosa del oeste ha sido ampliamente cuestionado. A fines de los veranos de 1995 y 1996 se midieron todas las exclusiones conocidas (n=8) de "Aspen" en los Bosques Nacionales Dixie y Fishlake situados en la región sur-central de Utah. Las mediciones realizadas en estas exclusiones fueron para determinar la dinámica de tallos, el estado sucesional y la composición de especies herbáceas. Cinco de las exclusiones estuvieron dentro de un diseño de tres factores: exclusión total, exclusión contra el ganado y uso combinado, el cual permitía medir separadamente los efectos de la herbívora de venados (*Odocoileus hemionus*) y alces (*Cervus elaphus*) de la del ganado. El "Aspen" se regeneró exitosamente dentro de las exclusiones, y sin la influencia del fuego u otro factor de disturbio, desarrollo una estructura de tallos de diferentes edades. El "aspen" sujeto al ramoneo de la fauna silvestre, principalmente venado, fallo en regenerarse exitosamente o se desarrolló con densidades de tallos significativamente menores (2,498 tallos ha⁻¹) que el "aspen" de las parcelas totalmente excluidas (4,474 tallos ha⁻¹). En las parcelas de uso combinado, ganado-fauna, la mayoría del aspen no se regeneró satisfactoriamente o lo hizo con bajas densidades de tallos (1,012 tallos ha⁻¹). El "Aspen" se regeneró exitosamente en las parcelas utilizadas por ungulados solo cuando el número de venados fue bajo. En forma similar, la herbívora de los ungulados tuvo efectos significativos en la composición del estrato herbáceo. En general, la utilización por el venado tendió a reducir los arbustos y las hierbas altas palatables mientras que favoreció el crecimiento de los zacates nativos. Por otra parte, la adición del apacentamiento de ganado domestico, tendió a reducir los zacates nativos y promovió el desarrollo de especies introducidas y de suelo desnudo. Así, las comunidades dominadas por arboles viejos o de una sola edad parecen ser el producto del ramoneo por los ungulados y no un atributo biológico del "Aspen" como comúnmente se ha asumido. No hubo evidencia de que la variación climática afectara la regeneración del "Aspen". Las diferencias observadas se atribuyeron a diferentes historiales de herbívora de los ungulados.

McDonough 1979). Because individual trees are relatively short-lived (< 150 years), long-lived aspen clones are often dependent on periodic disturbance such as fire to stimulate vegetative regeneration via root suckering, and to reduce conifer competition (Bartos and Mueggler 1981, Shepperd and Smith 1993).

Aspen has been declining throughout the Intermountain West since shortly after European settlement (Kay 1997a, Bartos and




Fig. 1. The decline of aspen on National Forests in Utah. Unpublished forest inventory data, Rocky Mountain Research Station, USDA Forest Service, used with permission.

Campbell 1998). On the Fishlake and Dixie National Forests in Utah, for example, there were historically over 304,000 ha of aspen; today there are only approximately 120,000 ha (Fig. 1). Furthermore, many aspen stands contain old-age or single-age trees and have not successfully regenerated for 80 years or longer (Mueggler 1989).

Several hypotheses have been advanced to explain this decline. These include fire suppression (Houston 1973, Despain et al. 1986), climate change (Despain et al. 1986, Romme et al. 1995), livestock grazing (Sampson 1919, Baker 1925), and browsing by mule deer (*Odocoileus hemionus*) and/or elk (*Cervus elaphus*) (Olmstead 1979, Shepperd and Fairweather 1994).

To test the ungulate herbivory hypothe-

ses, we measured all of the long-term, aspen-containing exclosures on the Fishlake and Dixie National Forests in south-central Utah. These exclosures, established during the 1930s to 1970s, were built to study the effect of livestock and/or wildlife use on aspen communities (Young 1956, Laycock 1969).

Methods

Laycock's (1975) list of range reference sites in Utah was first checked for aspen-containing exclosures on the Fishlake and Dixie National Forests. Forest Service employees on both national forests were then contacted to see if these exclosures still existed and if additional aspen-con-

taining exclosures were present on the 2 forests. These potential study sites were subsequently field-checked to see if the exclosures were still functional. We tried to locate in agency files all previous vegetation data, written description of permanent vegetation sampling schemes, and any old photographs.

Some of the exclosures were of a 2-part design where an inside area was fenced to exclude livestock, primarily cattle, but deer and elk had access. Adjacent unfenced outside plots were grazed by both wild and domestic ungulates. Other exclosures were of a 3-part design where 1 area was fenced to exclude all ungulates (treatment 1), another was fenced to exclude livestock (treatment 2), and outside areas were open to wild and domestic ungulates (treatment 3).

At each exclosure, any permanent aspen plots previously established by the agencies were resampled and permanent photo-points rephotographed. Sampling was done during late summer of 1995 and 1996. Next, 2 x 30 m belt transects were placed in representative aspen communities within each of the various parts of each exclosure (Kay 1990). The varied size of exclosures did not always allow sampling with an equal number of transects (Table 1). Each treatment at the following exclosures were sampled by 3 belt transects: Pot Holes, Blind Lake, Riddle Swale, and Hancock. Each treatment on Park Pasture was sampled by 4 transects, and Parker Mountain by 6. At Woodchuck, treatments 1 and 2 were sampled by 3 transects and treatment 3 by 2; at Grindstone, treatments 1 and 2 were also sampled by 3 transects, but treatment 3 by 4. To facilitate recording data, each 30 m transect was subdivided into 3 segments and the number of live aspen stems was recorded using the following size classes within each segment: (1) stems less than 2 m tall, (2) stems greater than 2 m tall but

Table 1. Location and description of aspen containing exclosures in south-central Utah, compiled in 1995–1996.

Exclosure	Land Ownership ¹	Location			Year established	Size (m)		Elevation	Main aspect	Vegetation type ²
		Township	Range	Section		Livestock exclusion	Total exclusion			
Pot Holes	DNF	30S	3E	27	1958	62x63	None	(m) 2743	NW	A,S/G
Park Pasture	DNF	31S	5E	22	1957	88x94	None	2652	NE	A,G
Blind Lake	DNF	30S	4E	24	1958	44x117	None	2984	SE	A,G
Riddle Swale	DNF	33S	1W	19	1957	65x76	65x113	2554	N	A,S/G,C
Woodchuck	DNF	31S	2 1/2W	33	1947	63x63	63x63	2804	S	A,S/G
Grindstone	FNF	29S	4W	29	1934	46x63	46x63	2816	S	A,C
Hancock	FNF	26S	1E	11	1962	61x61	61x61	3054	SE	A,C
Parker Mountain	UT	28S	1W	26	1974	100x100	100x100	2804	E	A,S/G

¹DNF = Dixie National Forest, FNF = Fishlake National Forest, UT = Utah State School Trust Lands.

²Vegetation types within the exclosures. A = aspen, S/G = sagebrush/grasslands, G = grasslands, and C = conifers

less than 5 cm diameter at breast height (DBH), (3) stems between 6 and 10 cm DBH, (4) stems between 11 and 20 cm DBH, and (5) stems greater than 21 cm DBH. Ages of aspen within each size class were determined by coring the larger and cutting the smaller stems and counting annual rings.

The number and species of conifers were recorded on the 2 x 30 m belt transects using the same size classes. While a 2 x 30 m belt transect is adequate to measure aspen stem dynamics and conifer seedlings (Bartos et al. 1994), a plot of that size tends to underestimate the more widely spaced conifers. To overcome this problem, the total percent conifer canopy cover was estimated in each aspen stand following Mueggler (1988).

Aspen and conifer stem counts on each of the 2 x 30 m belt transects were used to produce a mean number of stems per hectare by size classes for each treatment at each site. The treatment means of the various aspen size class data were then compared using repeated measures analysis of variance. Each site was considered an experimental unit for this analysis.

Frequency and canopy-cover of understory plants, bare soil, and rock were recorded from 0.1 m² plots at 1 m intervals to the right of each of the belt-transect's centerline (Daubenmire 1959). Readings from all the 0.1m² understory plots were averaged within each treatment and then compared using Student's t-test.

Other information recorded at each study site included: Universal Mercator Grid coordinates, elevation, aspect, old and new bark damage, highlining, and sucker browsing. The last 3 items provided an estimate of past ungulate use. Elk and moose (*Alces alces*) strip-off and eat the bark of aspen. Such bark damage usually occurs during winter when other foods are in short supply. Mule deer do not strip aspen bark, but mule deer, elk, and moose all browse aspen when the plants are available.

To determine the effects of deer herbivory on Indian paintbrush (*Castilleja miniata* Dougl.) we measured the height, crown diameter, total number of flower stalks, and the number of grazed flower stalks. These measurements were made just at the Hancock site.

Results

Many of the aspen-containing exclosures listed by Laycock (1969) for south-central Utah no longer exist or were not

maintained. Thus, only 8 exclosures containing aspen were evaluated (Table 1). Prior data were found for the Grindstone Flat exclosure. Photographs showing the originally enclosed aspen communities were found for 4 exclosures (Pot Holes, Park Pasture, Grindstone Flat, and Parker Mountain).

Pot Holes

With protection from cattle grazing, the aspen stand successfully established new stems without fire or other disturbance, and increased in size inside the exclosure as evidenced by photographs. The new suckers established in the early 1970s when mule deer numbers were low (Fig. 2). Later when deer numbers expanded the animals consumed all the lower branches from the newly grown aspen, termed highlining, and prevented new suckers from exceeding 1 m in height. The aspen stand inside the exclosure is now composed of

multi-sized, multi-aged stems, while little regeneration has occurred outside the exclosure (Table 2).

The largest live aspen were 28 cm DBH and 85+ years old. The 6–7 cm DBH stems were 20–21 years old and the 4 cm stems were 10–12 years old. There was fresh deer sign within the exclosure in August 1995. At this elevation (2,799 m), the conifers are mostly ponderosa pine (*Pinus ponderosa* Lawson) with a few Douglas fir (*Pseudotsuga menziesii* (Mirbel) Franco).

Major differences in understory species composition inside and outside this exclosure were significant ($P < 0.05$). Shrubs, such as rabbitbrush (*Chrysothamnus* spp. Nutt.) had lower frequency and canopy cover inside the exclosure, while needle-and-thread grass (*Stipa comata* Trin. & Rupr.) was more frequent inside the fenced plot, as was total grass canopy cover. Indian paintbrush (*C. linariaefolia* Benth.) was more frequent where live-

Table 2. Mean aspen stem densities by size classes and estimated conifer canopy cover inside and outside exclosures in south-central Utah in 1995–96.

Exclosure	Mean aspen stem density by size classes					Conifer canopy cover
	<2m tall	2m tall to 5cm DBH	6 to 10cm DBH	11 to 20cm DBH	>20cm DBH	
	----- (stems ha ⁻¹) -----					(%)
Pot Holes						
Livestock exclusion	7,348	3,791	2,221	618	333	<1
Outside	9,018	668	0	55	55	<1
Park Pasture						
Livestock exclusion	1,221	4,676	3,758	1,378	250	<1
Outside	459	835	835	0	584	<1
Blind Lake						
Livestock exclusion	9,574	8,684	3,228	1,057	556	<1
Outside	8,963	6,568	167	55	668	<1
Riddle Swale						
Total exclusion	7,228	4,670	2,780	1,890	168	<1
Livestock exclusion	8,613	2,446	834	0	449	<1
Outside	1,890	0	111	0	222	5
Woodchuck						
Total exclusion	5,177	4,787	2,394	1,169	390	None
Livestock exclusion	946	0	56	779	390	None
Outside	2,088	0	0	1,086	0	None
Grindstone Flat						
Total exclusion	NA ¹	2,505	3,396	1,614	333	30
Livestock exclusion	NA	0	222	835	668	50
Outside	NA	0	0	167	752	60
Hancock						
Total exclusion	5,288	7,738	1,225	557	445	11
Livestock exclusion	2,728	0	0	0	557	31
Outside	4,676	0	0	0	500	30
Parker Mountain						
Total exclusion	974	2,672	1,308	84	500	None
Livestock exclusion	4,509	389	0	0	84	None
Outside	1,030	28	0	0	306	None
Means						
Total Exclusion	4,668a ²	4,474a	2,221a	1,063a	734a	–
Livestock Exclusion	4,991a	2,498b	1,289ab	583a	822a	–
Outside	4,018a	1,012b	139b	170a	772a	–

¹NA = Stems less than 2m tall were not counted because most had been consumed by a recent fire – see text.

²Means within columns followed by the same letter are not different at $P < 0.08$ by analysis of variance on means.

stock had been excluded, while *Antennaria microphylla* Rydb. was more prevalent outside the exclosure. There was also more bare soil and less litter outside the exclosure than inside.

Park Pasture

When the Park Pasture exclosure was erected on the east side of Boulder Mountain, the area was a heavily used sheep-cattle allotment. Sheep were eliminated in 1963 and cattle herbivory subsequently reduced (U.S. Forest Service 1995). Historically, the area was also grazed by high numbers of mule deer, but that herd has since declined (Fig. 2). When the exclosure was built elk were rare, but they are now common. In fact, several fresh elk beds were observed inside the exclosure when the site was visited during August 1995.

There were no young aspen stems in 1957, but with exclusion of livestock, aspen regenerated and spread inside the exclosure (Fig. 3). This exclosure is also on an aspen-grassland ecotone with few conifers. Ponderosa pine, Douglas-fir, spruce (*Picea* spp. A. Dietr.), and sub-alpine fir (*Abies lasiocarpa* (Hook.) Nutt.) are the most common species in the immediate area. The largest aspen were 38–40 cm DBH and were 105–115 years old. Inside the exclosure there were numerous 10–15 cm DBH stems that were 25–30 years old, but there were none outside (Table 2). These began growth during the mid-1960's when combined high deer and cattle grazing apparently prevented aspen regeneration outside the exclosure. When deer numbers declined during the early to mid-1970s (Fig. 2), aspen regenerated both inside and outside the exclosure, but trees were about 5-times more numerous where cattle also were excluded (Table 2). The 6–10 cm DBH stems were all approximately 20 years old. Rebounding deer populations during the 1980s highlined the regenerated aspen and prevented new suckers from growing taller. Little difference, however, existed in understory species composition or frequency.

Blind Lake

The Blind Lake exclosure on the north side of Boulder Mountain is in an area comprised primarily of aspen and small meadows. Aspen stems inside the exclosure are multi-sized and multi-aged, and began regenerating shortly after the exclosure was erected (Table 2). The elimination of cattle grazing reduced browsing on the aspen suckers and allowed them to grow into the larger size classes. Aspen




Fig. 2. The number of deer and elk harvested by hunters in Utah from 1925 to 1995. (Utah Div. of Wildlife Resources' Annual big Game Reports—data smoothed to show historical trends.)

outside the exclosure regenerated approximately 20 years ago when deer populations plummeted (Fig. 2), despite continued access by cattle. The largest aspen were 30–38 cm DBH and 95–105 years old, while the 4–5 cm DBH stems were 18–25 years of age. Photographic evidence indicates that the aspen clone inside the exclosure has increased in area displacing a largely grassland community.

There was significantly less snowberry (*Symphoricarpos oreophilus* Gray) ($P<0.01$) and lovage (*Ligusticum porteri* Coult. & Rose) ($P<0.01$), and significantly more bare soil ($P<0.01$) and dandelion (*Taraxacum officinale* Weber ex Wiggers) outside the exclosure. In general, there were more palatable forbs (U.S. Forest Service 1937, Nelson and Leege 1982, Wallmo and Regelin 1981) inside than outside the exclosure, while less palatable and non-native forbs were more common outside than inside. There was also more vegetative cover inside than outside the exclosure ($P<0.01$).

Riddle Swale

Riddle Swale is a 3-part exclosure with a total-exclusion portion, a livestock-exclusion portion, and an outside plot open to both wildlife and cattle. The site is located on the west side of Escalante Mountain at the ecotone between aspen, sagebrush, and pinyon-juniper. A few Utah juniper (*Juniperus osteosperma* (Torr.) Little), pinyon pine (*Pinus monophylla* Torr. & Frem. In Frem.), ponderosa pine, and Douglas-fir are found both inside and outside the exclosure.

Inside the total-exclusion part aspen began regenerating shortly after the exclosure was constructed in 1957; the stems are now multi-sized and multi-aged (Table 2). In the livestock-exclusion portion of the exclosure, aspen only regenerated when deer numbers declined around 20 years ago (Fig. 2). As the deer recovered, however, they highlined all the previous aspen regeneration inside the livestock exclosure and prevented any new aspen suckers from growing more than 1 m tall. Where both cattle and deer grazed, few aspen stems attained more than 2 m in height (Table 2). The largest aspen at this site were 23–26 cm DBH and 72–78 years old. The 8–10 cm DBH stems were 20–25 years old, and the 3–5 cm stems were 14–20 years old.

There were also major differences in understory species composition. Deer use only and combined use eliminated Indian paintbrush, while combined big game and cattle use significantly ($P<0.01$) reduced the canopy cover of native grasses compared to the exclosures. Conversely, sagebrush increased under deer use, but more so under combined use. Bare soil and rock were not apparent where ungulate herbivory was absent, but increased to 6.4% cover where only wildlife grazed and to 27.2% cover with combined deer-cattle use ($P<0.01$).

Woodchuck

Woodchuck is a 3-way exclosure built on the northeast slope of Mount Dutton. Although this exclosure is at over 2,800 m



Fig. 3. Photographs of the Park Pasture enclosure: (a) taken in September 1958 shortly after the enclosure was constructed and showing no aspen regeneration; and (b) in August 1995 showing spread of aspen inside the enclosure.

in elevation, it is surrounded by extensive sagebrush-grasslands and lacks conifers. Cattle have used this allotment historically, and deer numbers have also been high. Elk have recently become common on Mount Dutton but do not appear to use this isolated patch of aspen.

Aspen within the total-exclusion part is multi-sized and multi-aged (Table 2), and began regenerating shortly after the enclosure was constructed in 1947. No new aspen stems have attained heights of 2 m or more in the livestock-exclusion or combined-use areas. The deer at this site have not allowed aspen to regenerate even where cattle have been excluded. The largest aspen were 25–30 cm DBH and 100+ years old, while aspen in the 15–16 cm size class were 65–70 years of age. Apparently, some event in the late 1920s and early 1930s allowed some aspen regeneration in this stand before it was enclosed.

Deer use significantly reduced wild rose (*Rosa woodsii* Lindl.), snowberry, lupine (*Lupinus caudatus* Kellogg), and Indian paintbrush, but favored the growth of native grasses, especially needle and thread (all $P < 0.01$). The addition of cattle grazing significantly reduced native grass cover (< 0.01) and significantly increased the amount of bare soil ($P < 0.01$). The canopy cover of introduced Kentucky bluegrass (*Poa pratensis* L.) increased from 1.9% in the total exclusion area, to 23.7% under deer only use ($P < 0.01$), and to 50.9% with the addition of cattle grazing ($P < 0.01$). Total vegetative understory cover declined significantly ($P < 0.01$) from 95.9% in the non-grazed area to 77.7% in the wildlife only area to 63.7% with combined use.

Grindstone Flat

The 3-part enclosure at Grindstone Flat (Table 1) was evaluated during the 1950s (Young 1956), and then remeasured by the U.S. Forest Service in 1975 (Mueggler and Bartos 1977). Part of the site was originally clear-cut prior to enclosure construction, but 1/3 of each enclosure, as well as all of the outside aspen, were left uncut. Only the uncut areas were analyzed to maintain comparability with the other sites.

Aspen regenerated inside the total-exclusion area, while deer use prevented aspen regrowth in the livestock-exclusion area, as well as the outside area (Table 2). Spruce and subalpine fir have heavily invaded the area (Mueggler and Bartos 1977).

During late June 1996 the enclosure and much of the surrounding lands were burned by the Pole Creek wildfire. Much of the vegetation inside the enclosure complex was burned, and the old enclosure pole-fence destroyed. The total-exclusion and livestock-exclusion areas were completely consumed by flames, as was the surrounding outside aspen. In contrast, the cut total-exclusion area was largely unburned while the cut livestock-exclusion area did not burn at all.

Since this is 1 of only 2 long-term enclosures on the Fishlake National Forest containing aspen, the Forest Service rebuilt the enclosure fences in 1997. We were able to measure aspen stem densities in September 1996 because the larger-sized live stems were too green to be consumed by the fire. Aspen stems less than 2 m tall, however, were not quantified because many of those were removed by

the fire. Understory species composition was not measured for the same reason.

Of most relevance in this enclosure comparison is that aspen in the total-exclusion area successfully regenerated and produced a multi-aged stand prior to the 1996 fire, while aspen in the livestock enclosure and outside combined use areas produced no new stems greater than 2 m tall. Conifer cover in the total-exclusion area was less than in the areas accessible to either deer or cattle (Table 2).

The largest aspen trees were 38–44 cm DBH, but age determination was difficult because of extensive heart rot. In 1975, Mueggler and Bartos (1977) recorded a 41 cm DBH aspen that was 175 years old, so the few remaining, unburned trees were likely approaching or near the maximum age (200 years) recorded for aspen in the Intermountain West (Jones and Schier 1985).

Hancock

This 3-part enclosure is located above Doctor Canyon, approximately 3 km southeast of Hancock Flat and 1 km northeast of Rust Spring. Unlike the other enclosures in this study, Hancock has been a sheep allotment and mule deer were the most abundant wild ungulate.

Aspen in the total-exclusion portion began regenerating shortly after the enclosure was constructed and today is multi-sized and multi-aged despite invasion by conifers (Table 2). Aspen has not regenerated successfully in either the wildlife-only or the combined-use areas; all of the aspen suckers that occur have been repeatedly browsed by deer and/or sheep. When this site was measured in July, mule deer

had already browsed many of the new aspen suckers in the livestock-exclusion portion of the enclosure. The largest aspen stems were 30 cm DBH and 105+ years old. Spruce canopy cover appeared greater on the grazed portion than on the ungrazed portion of the enclosure.

Deer grazing and combined use created major differences in understory species composition. Both deer grazing and combined use significantly ($P < 0.01$) reduced Indian paintbrush, dandelion, and total vegetative cover, and significantly ($P < 0.01$) increased cover of grass, yarrow (*Achillea millefolium* L.), rock, bare soil, and litter. Deer grazing also significantly ($P < 0.01$) reduced height, crown diameter, and number of flower stalks of individual Indian paintbrush plants. Deer had already grazed 63% of the Indian paintbrush flower stalks inside the livestock-exclusion portion of this enclosure when it was sampled in July.

Parker Mountain

When this site was fenced (1974) there was concern that browsing by jackrabbits (*Lepus* spp.) might effect aspen regeneration. To monitor this, each of the total-exclusion and livestock-exclusion areas were partitioned with additional fencing to prevent entry by lagomorphs on half of these areas.

The enclosure is on a joint cattle-sheep allotment where mule deer and antelope (*Antilocapra americana*) have been abundant. In recent years, however, the deer population has declined and most of the present wildlife use is by elk. The enclosures are also near the lower elevation of aspen and the site is surrounded by sagebrush-grasslands.

Exclusion of lagomorphs had no effect on aspen regeneration or understory composition so the 2 data sets were pooled. When protected from ungulates, aspen successfully regenerated on this site producing a multi-aged stand (Table 2). Aspen successfully regenerated in the livestock-exclusion part and on outside plots, especially after deer declined, but at significantly lower ($P < 0.05$) densities than on the total exclusion plot. The increasing elk population, however, highlined those stems and inflicted extensive bark damage (Krebill 1972) on the unprotected aspen. The largest aspen were 26–28 cm DBH and 90–95 years of age. Photographic evidence indicates that aspen in the total-exclusion part of this enclosure has spread into sagebrush-grasslands, an indication that climate had not limited expansion at this site.

Although this enclosure had been in place for only 22 years, there were significant ($P < 0.05$) differences in understory species composition. Indian paintbrush and littleflower penstemon (*Penstemon procerus* Dougl. ex Graham) had greater frequency and canopy cover in the total-exclusion area than on the wildlife exclusion area, and they were virtually eliminated by combined wildlife-livestock use. Conversely, native grasses were most abundant in the total-exclusion part (54%), were halved by wildlife-only use, and reduced to only 16% under combined use. The addition of livestock grazing also increased ($P < 0.05$) the amount of bare soil.

Discussion

Aspen Stem Dynamics

Aspen within all total-exclusion enclosures regenerated successfully without disturbance and developed multi-sized, multi-aged stems, even where the stand had been heavily invaded by conifers (i.e., the Hancock enclosure). Aspen subjected to browsing only by wildlife, primarily mule deer, either failed to produce new stems greater than 2 m tall, or regenerated at stem densities lower than on the total-exclusion plots. On combined wildlife-livestock use plots, most aspen failed to regenerate successfully, or did so at stem densities lower than on the livestock-exclusion plots. Aspen regenerated successfully on livestock-excluded and combined-use plots only when mule deer pop-

ulations were low.

Mule deer have been the most prevalent wild herbivore in south-central Utah since at least the 1930s. Moose were only recently transplanted into the area, and elk populations were very low until the 1980s. Deer numbers in Utah generally peaked during the early 1960s, but declined precipitously by 1975 (Fig. 2). This reduced the browsing pressure on aspen suckers, and many stands in southern Utah were able to regenerate successfully if livestock use was not excessive. An increase in deer during the 1980s prevented new aspen suckers from increasing in height, and the deer also consumed the lower branches from aspen that had regenerated earlier (Fig. 4). In the mid-1990s, mule deer populations again declined, but few aspen stands were able to regenerate successfully because increasing numbers of elk were foraging on the young suckers (Fig. 2). Thus, in some areas of Utah, deer browsing has been replaced by elk browsing. Elk also seem to have a greater preference for aspen than mule deer (Kay 1997b, Shepperd and Fairweather 1994). Episodes of aspen regeneration have been observed in other areas when mule deer (Julander and Low 1976, Olmstead 1979) or elk (Olmstead 1979, Kay and White 1995) numbers were low.

Understory Species Composition

Understory species composition of aspen stands was significantly affected by ungulate herbivory. In general, utilization by mule deer tended to eliminate palatable



Fig. 4. Highlining by mule deer in a 2-age class aspen stand on the Dixie National Forest.

Table 3. Mean frequency (Freq.) and canopy cover (C.C.) of undergrowth species on total-exclusion (wild & domestic ungulates), partial exclusion (livestock), and no-exclusion plots. (Means are based on only those sites where the species occurred; "n" = number of sites where species occurred; summation of cover values therefore are meaningless.)

	TOTAL EXCLUSION			PARTIAL EXCLUSION			NO EXCLUSION		
	n	Freq.	C.C.	n	Freq.	C.C.	n	Freq.	C.C.
	----- (%) -----			----- (%) -----			----- (%) -----		
TREES									
<i>Picea engelmannii</i> Parry ex Engelm.	1	18	10.6	2	45	30.3	2	23	14.8
SHRUBS									
<i>Artemisia tridentata</i> Nutt.	3	20	7.7	4	24	12.2	4	25	11.3
<i>Artemisia tripartita</i> Rydb.	1	3	0.4	2	2	0.2	2	15	4.5
<i>Chrysothamnus nauseosus</i> (Pallas) Britt.	1	1	0.1	2	4	1.1	2	8	2.1
<i>Chrysothamnus viscidiflorus</i> (Hook.) Nutt.	1	0	0.0	2	15	4.8	2	20	8.1
<i>Juniperus communis</i> L.	0	0	0	1	0	0.0	1	1	0.4
<i>Juniperus osteosperma</i> (Torr.) Little	1	7	3.3	1	0	0.0	1	8	5.9
<i>Potentilla fruticosa</i> L.	0	0	0	1	4	0.3	1	5	0.7
<i>Purshia tridentata</i> (Pursh) DC.	1	0	0.0	1	0	0.0	1	2	1.9
<i>Ribes</i> spp. L.	0	0	0	1	8	2.9	1	4	1.4
<i>Rosa woodsii</i> Lindl.	3	2	0.9	4	3	0.8	4	4	0.8
<i>Symphoricarpos oreophilus</i> Gray	3	17	8.0	5	19	8.4	5	11	3.3
GRAMINOIDS									
<i>Agropyron caninum</i> (L.) Beauv.	3	28	4.0	3	16	2.4	3	8	0.7
<i>Bromus ciliatus</i> L.	4	28	5.6	6	15	1.7	6	10	1.0
<i>Carex</i> spp. L.	2	18	0.1	1	17	4.2	1	0	0.0
<i>Festuca idahoensis</i> Elmer	2	1	0.1	3	42	9.7	3	39	6.5
<i>Festuca thurberi</i> Vasey	2	40	13.4	4	23	5.2	4	31	6.2
<i>Koeleria cristata</i> (L.) Pers.	1	4	0.5	1	35	4.2	1	24	2.6
<i>Muhlenbergia wrightii</i> Vasey ex Coult.	0	0	0	2	1	0.1	2	16	3.4
<i>Poa interior</i> (NY)	1	7	1.1	1	0	0.0	1	8	0.8
<i>Poa pratensis</i> L.	2	8	2.8	4	22	15.4	4	42	18.8
<i>Sitanion hystrix</i> (Nutt.) J. G. Sm.	3	18	2.6	5	19	3.0	5	18	2.5
<i>Stipa columbiana</i> Macoun	1	4	1.8	1	0	0.0	1	0	0.0
<i>Stipa comata</i> Trin. & Rupr.	3	42	11.7	6	46	14.1	6	30	4.1
<i>Stipa lettermanii</i> Vasey	1	17	2.3	1	11	2.2	1	0	0.0
FORBS									
<i>Achillea millefolium</i> L.	2	10	0.8	3	34	2.7	3	25	14.6
<i>Antennaria microphylla</i> Rydb.	0	0	0	1	0	0	1	54	11.9
<i>Aquilegia coerulea</i> James	1	8	2.3	1	3	0.2	1	0	0
<i>Aster</i> spp. L.	1	13	2.4	1	9	0.9	1	9	0.7
<i>Castilleja linariaefolia</i> Benth.	3	28	3.9	4	6	1.1	4	1	0.1
<i>Castilleja miniata</i> Dougl. ex Hook.	1	56	29.5	1	6	0.4	1	2	0.2
<i>Fragaria virginiana</i> Duchesne	0	0	0	1	16	1.2	1	3	0.2
<i>Galium</i> spp. L.	1	38	3.9	1	28	3.6	1	0	0.0
<i>Ligusticum porteri</i> Coult. & Rose	0	0	0	2	42	12.3	1	0	0.0
<i>Lupinus argenteus</i> Pursh	1	37	14.2	2	6	1.5	2	29	6.7
<i>Lupinus caudatus</i> Kellogg	1	82	40.4	1	0	0.0	2	18	4.4
<i>Penstemon procerus</i> Dougl. ex Graham	1	18	3.7	1	0	0.0	1	0	0.0
<i>Phlox</i> spp. L.	1	0	0.0	1	22	4.3	1	6	1.2
<i>Potentilla glandulosa</i> Lindl.	1	40	8.0	1	45	5.5	1	10	1.0
<i>Potentilla gracilis</i> Dougl. ex Hook.	0	0	0	1	2	1.3	1	0	0.0
<i>Senecio</i> spp. L.	1	10	2.0	1	0	0	1	0	0
<i>Taraxacum officinale</i> Weber ex Wiggers	1	29	4.6	3	14	2.6	3	31	8.7
<i>Vicia americana</i> Muhl.	0	0	0	1	83	25.2	1	72	26.5
ROCK	4	1	0.3	7	8	3.1	7	9	3.1
BARE SOIL	4	3	0.6	7	13	2.0	7	30	9.4
LITTER	4	63	25.3	7	66	26.8	7	69	28.9

tall forbs and shrubs while favoring the growth of native grasses and unpalatable forbs (U.S. Forest Service 1937, Nelson and Leege 1982, Wallmo and Regelin 1981) (Table 3). Indian paintbrush in particular was harmed by mule deer herbivory. The addition of livestock, primarily cattle, significantly reduced the native grass cover at most sites, while introduced grasses and bare soil increased. It is clear

that the combined level of ungulate use at all sites has not only altered the age structure of aspen stands but also altered understory species composition. It is equally clear that wildlife can dramatically affect the structure of aspen communities. Earlier open-range grazing studies have reported similar findings (Baker 1925, Weatherill and Keith 1969, Julander and Low 1976, Mueggler 1988, Smith et al. 1972).

Numerous hummingbirds were observed feeding on Indian paintbrush inside the total-exclusion area at the Hancock enclosure, but not inside the livestock-exclusion area or on outside plots. Thus, grazing-induced reduction of this flowering forb alone may have major effects on species not normally considered in range-wildlife studies.

Other Aspen Exclosure Studies

Aspen exclosure studies throughout the western United States and Canada depict similar findings. Coles (1965) and Mueggler and Bartos (1977) report similar results in central Utah. Kay (1990) measured 14 aspen exclosures in the Yellowstone Ecosystem where elk are the major herbivore. He found that all protected stands regenerated and developed multi-aged stands and that protection completely altered the composition of the undergrowth: shrubs and palatable forbs dominated inside the exclosures, whereas non-native grasses resistant to grazing dominated outside. In 3 out of 4 exclosures in Colorado's Rocky Mountain National Park, where deer and elk are the primary herbivores, aspen developed multi-aged stands while areas outside did not (Hess 1993, Baker et al. 1997). In South Dakota, aspen in Custer State Park expanded into grasslands where wildlife were excluded (Hoffman and Alexander 1987), and aspen in Wind Cave National Park develop multi-size class stands after exclusion of wild ungulates (Kay 1990).

From study of 4 exclosures in Alberta's Elk Island National Park where elk and moose are the major herbivores, Milner (1977) concluded that not only was aspen regeneration restricted by browsing, but palatable shrubs were more abundant under protection. Trottier and Fehr (1982), evaluating exclosures in Canada's Banff National Park, concluded that aspen regeneration was limited by elk browsing and that protected plots had both greater shrub density and more diverse height than the unprotected plots. Comparing repeat photographs taken of an exclosure (after 50 years protection) near Banff, Kay et al. (1994) observed that a dense multi-aged aspen stand had grown up inside the exclosure while no aspen stems had regenerated successfully outside. These same authors also reported that aspen regenerated successfully where protected for approximately 10-years within the game-proof fenced Trans Canada Highway right-of-way through Banff's lower Bow Valley, but did not where elk were free to browse.

The consensus of all of these studies is that deer and elk can significantly hinder aspen regeneration and change understory species composition. Moreover, livestock use has an additive negative effect on aspen regeneration and understory composition. Excessive use by wildlife tends to reduce woody species and palatable forbs. Utilization by mule deer favors native grasses. Excessive elk use has a negative effect on grasslands because elk utilize a broader array of forages than deer (Nelson

and Legee 1982). Excessive use by cattle also tends to reduce the abundance of native grasses and increase the amount of non-native species and bare soil.

Climate Change

The exclosures also demonstrate that climatic variation has had little effect on reproduction in aspen communities compared to that imposed by ungulates. Since the exclosure fence usually bisects a single aspen clone, differences between protected and open areas cannot be a product of either genetics or climate. The more abundant vegetation inside the exclosures, especially on total-exclusion plots, alters the microclimate, but that is an incorporated variable caused by the plant's response to the elimination of ungulate browsing, not the cause of the vegetation's response. Such microclimatic conditions would prevail in any aspen stand not subject to heavy ungulate use, whether in an exclosure or not.

If climatic variation is having an overriding effect on aspen community dynamics as proposed by others (Romme et al. 1995), we would expect aspen inside exclosures in south-central Utah to show signs of stress, especially since many of the exclosures are situated at the lower elevation of aspen or at grassland-aspen ecotones. However, no signs of physiological stress were observed during this study. In fact, the area occupied by aspen has increased inside exclosures at the expense of grasslands (Fig. 3). Baker et al. (1997) reported no correlation between climatic variation and aspen regeneration, while Kay (1990) reported that enclosed aspen in the Yellowstone Ecosystem replaced grasslands even on south-facing hillsides. White et al. (1998) and Kay (1997b) similarly reported no correlation between climate and aspen regeneration in the Canadian Rockies.

Conclusions

1. Browsing by native and domestic ungulates has hindered aspen regeneration throughout south-central Utah.
2. Aspen need not always be burned or clear-cut to regenerate successfully.
3. Wild ungulates, primarily mule deer, can have a major effect on aspen stem dynamics and understory composition.
4. Livestock grazing, as historically practiced in southern Utah, has had widespread effects on aspen communities, including changes in understory species composition.

5. Combined wildlife-livestock use most severely alters aspen community dynamics.

6. Aspen stands in the Rocky Mountain west dominated by old or single-age trees are most likely a product of excessive ungulate browsing.

7. Managers should quantify the level of ungulate herbivory before treating aspen stands with fire or cutting, because developing suckers may be subject to repeated browsing. If ungulate browsing is excessive, treatment of aspen stands may only hasten their demise.

Literature Cited

- Baker, F.S. 1925. Aspen in the central Rocky Mountain region. USDA Agr. Bull. No. 1291.
- Baker, W.L., J.A. Monroe, and A.E. Hessler. 1997. The effects of elk on aspen on the winter range in Rocky Mountain National Park. *Ecography* 20:155-165.
- Bartos, D.L. and R.B. Campbell, Jr. 1998. Decline of aspen in the interior West—examples from Utah. *Rangelands*. 20:17-24.
- Bartos, D.L. and W.F. Mueggler. 1981. Early succession in aspen communities following fires in western Wyoming. *J. Range Manage.* 34:315-318.
- Bartos, D.L., J.K. Brown, and G.D. Booth. 1994. Twelve years biomass response in aspen communities following fire. *J. Range Manage.* 47:79-83.
- Coles, F.H. 1965. The effects of big game and cattle grazing on aspen regeneration. M.S. Thesis, Brigham Young Univ., Provo, Utah.
- Daubenmire, R. 1959. A canopy-coverage method of vegetational analysis. *Northwest Sci.* 33:43-64.
- Despain, D.G., D. Houston, M. Meagher, and P. Schullery. 1986. Wildlife in transition: Man and nature on Yellowstone's northern range. Roberts Rinehart Inc., Boulder, Colo.
- Hess, K. Jr. 1993. Rocky times in Rocky Mountain National Park: An unnatural history. University Press of Colorado, Niwot, Colo.
- Hoffman, G.R. and R.A. Alexander. 1987. Forest vegetation of the Black Hills National Forest of South Dakota and Wyoming: A habitat type classification. USDA Forest Serv. Res. Paper RM-276. Fort Collins, Colo.
- Houston, D.B. 1973. Wild fires in northern Yellowstone National Park. *Ecol.* 54:1111-1117.
- Jones, J.R. and G.A. Schier. 1985. Growth pp. 19-24. In: Debyle, N.V. and R.P. Winokur eds. *Aspen: Ecology and Management in the Western United States*. U.S. Forest Service Gen. Tech. Rep. RM-119. 382 pp.
- Julander, O. and J.B. Low. 1976. A historical account and present status of the mule deer in the West. p. 3-19. In: Workman, G.W. and J.B. Low, (eds.), *Mule deer decline in the West: A symposium*. Utah State Univ. Agr. Exp. Sta., Logan, Utah.

- Kay, C.E. 1990.** Yellowstone's northern elk herd: A critical evaluation of the "natural regulation" paradigm. Ph.D. Diss., Utah State Univ., Logan, Utah.
- Kay, C.E. 1997a.** Is aspen doomed? *J. Forest.* 95:4-11.
- Kay, C.E. 1997b.** The condition and trend of aspen, *Populus tremuloides*, in Kootenay and Yoho National Parks: Implications for ecological integrity. *Canad. Field-Nat.* 111:607-616.
- Kay, C.E., and C.W. White. 1995.** Long-term ecosystem states and processes in the central Canadian Rockies: A new perspective on ecological integrity and ecosystem management. p. 119-132. *In:* Linn, R.M., (ed.) Sustainable society and protected areas. The George Wright Society, Hancock, Mich.
- Kay, C.E., B. Patton, and C. White. 1994.** Assessment of long-term terrestrial ecosystem states and processes in Banff National Park and the Central Canadian Rockies. Resource Conservation, Parks Canada, Banff National Park, Banff, Alberta.
- Kreibill, R.G. 1972.** Mortality of aspen on the Gros Ventre elk winter range. USDA Forest Serv. Res. Paper INT-129. Ogden, Utah.
- Laycock, W.A. 1969.** Exclosures and natural areas on rangelands in Utah. USDA Forest Serv. Res. Paper INT-62. Ogden, Utah.
- Laycock, W.A. 1975.** Rangeland reference areas. Soc. for Range Manage., Range Sci. Series 3.
- McDonough, W.T. 1979.** Quaking aspen seed germination and early seedling growth. USDA Forest Serv. Res. Paper INT-234. Ogden, Utah.
- Milner, B.J. 1977.** Vegetation analysis of ungulate range exclosures, Elk Island National Park. Canadian Wildlife Serv., Edmonton, Alberta.
- Mueggler, W.F. 1988.** Aspen community types of the Intermountain region. USDA Forest Serv. Gen. Tech. Rep. INT-250. Ogden, Utah.
- Mueggler, W.F. 1989.** Age distribution and reproduction of aspen stands in the Intermountain Region. *West. J. Appl. Forest.* 4:41-45.
- Mueggler, W.F. and D.L. Bartos. 1977.** Grindstone Flat and Big Flat exclosures—a 41-year record of changes in clearcut aspen communities. USDA Forest Serv. Res. Paper INT-195. Ogden, Utah.
- Nelson, J.R. and T.A. Leege. 1982.** Nutritional requirements and food habits. p. 323-367. *In:* Thomas, J.S. and D.E. Toweill, (eds.), *Elk of North America: Ecology and management.* Stackpole Books, Harrisburg, Penn.
- Olmsted, C.E. 1979.** The ecology of aspen with reference to utilization by large herbivores in Rocky Mountain National Park. p. 89-97. *In:* Boyce, M.S., and L.D. Hayden-Wing, (eds.), *North American elk: Ecology, behavior, and management.* Univ. of Wyoming, Laramie, Wyo.
- Romme, W.H., M.G. Turner, L.L. Wallace, and J.S. Walker. 1995.** Aspen, elk, and fire in northern Yellowstone National Park. *Ecol.* 76:2097-2106.
- Sampson, A.W. 1919.** Effect of grazing upon aspen reproduction. USDA Bull. No.741.
- Shepperd W.D. and F.W. Smith. 1993.** The role of near-surface lateral roots in the life cycle of aspen in the central Rocky Mountains. *For. Ecol. and Manage.* 61:157-170.
- Shepperd, W.D. and M.L. Fairweather. 1994.** Impact of large ungulates in restoration of aspen communities in a southwestern ponderosa pine ecosystem. p. 344-347. *In:* Covington, W.S., and L.F. DeBano, (eds.), *Sustainable ecological systems: Implementing an ecological approach to land management.* USDA Forest Serv. Gen. Tech. Rep. RM-247. Fort Collins, Colo.
- Smith, A.D., P.A. Lucas, C.O. Baker, and G.W. Scotter. 1972.** The effects of deer and domestic livestock on aspen regeneration in Utah. Utah Division of Wildlife Resources Pub. 72-1. Salt Lake City, Utah.
- Sokal, R.R. and F.J. Rohlf. 1981.** Biometry: The principles and practice of statistics in biological research. 2nd ed. W.H. Freeman and Co., San Francisco, Calif.
- Trottier, G.C. and A. Fehr. 1982.** Re-evaluation of four range exclosures in Banff National Park, 1981. Canadian Wildlife Service, Edmonton, Alberta.
- U.S. Forest Service. 1937.** Range plant handbook. USDA, Forest Service, Washington, D.C. 841 pp.
- U.S. Forest Service. 1995.** Environmental assessment for revised east slope cattle allotment management. Dixie National Forest, Teasdale, Utah.
- Wallmo, O.C., and W. L. Regelin. 1981.** Rocky mountain and intermountain habitats. Part 1. Food habits and nutrition. pp. 387-398 in Wallmo, O.C., ed. *Mule and blacktailed deer of North America.* Univ. of Nebraska Press, Lincoln, Nebr. 605 pp.
- Weatherill, R.G. and L.B. Keith. 1969.** The effect of livestock grazing on an aspen forest community. Alberta Dept. of Lands and Forests, Fish and Wildlife Div. Tech. Bull. No. 1. Calgary, Alberta.
- White, C.A., C.E. Olmstead, and C.E. Kay. 1998.** Aspen, elk, and fire in the Rocky Mountain national parks of North America. *Wildl. Soc. Bull.* 26:449-462.
- Young, S. 1956.** Survey and evaluation of big game exclosures in Utah. Utah State Dept. of Fish and Game, Federal Aid Division, Dept. Inform. Bull. No. 24. Salt Lake City, Utah.