



## The Meteoritical Bulletin, No. 90, 2006 September

Harold C. CONNOLLY, JR.<sup>1,2\*</sup>, Jutta ZIPFEL<sup>3</sup>, Jeffrey N. GROSSMAN<sup>4</sup>, Luigi FOLCO<sup>5</sup>, Caroline SMITH<sup>6</sup>, Rhian H. JONES<sup>7</sup>, Kevin RIGHTER<sup>8</sup>, Michael ZOLENSKY<sup>8</sup>, Sara S. RUSSELL<sup>6</sup>, Gretchen K. BENEDIX<sup>6</sup>, Akira YAMAGUCHI<sup>9</sup>, and Barbara A. COHEN<sup>10</sup>

<sup>1</sup>Department of Physical Science, Kingsborough Community College, Oriental Boulevard, Brooklyn, New York 11235, USA  
and the Department of Earth and Environmental Sciences, The Graduate School of the City University of New York,  
New York, New York 10016, USA

<sup>2</sup>Department of Earth and Planetary Science, American Museum of Natural History, Central Park West, New York, New York 10024, USA

<sup>3</sup>Sektion Meteoritenforschung Forschungsinstitut und Naturmuseum Senckenberg, Senckenberganlage 25,  
D-60325, Frankfurt am Main, Germany

<sup>4</sup>U.S. Geological Survey, MS 954, Reston, Virginia 20192, USA

<sup>5</sup>Museo Nazionale dell'Antartide, Via Laterina 8, I-53100 Siena, Italy

<sup>6</sup>Department of Mineralogy, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

<sup>7</sup>Department of Earth and Planetary Sciences, University of New Mexico, Albuquerque, New Mexico 87131, USA

<sup>8</sup>Code ST, NASA Johnson Space Center, Houston, Texas 77058, USA

<sup>9</sup>Antarctic Meteorite Research Center, National Institute of Polar Research, 1-9-10 Kaga, Itabashi, Tokyo 173-8515, Japan

<sup>10</sup>Institute of Meteoritics, MSC03-2050 University of New Mexico, Albuquerque, New Mexico 87506, USA

\*Corresponding author. E-mail: [meteorite@kingsborough.edu](mailto:meteorite@kingsborough.edu)

*(Received 11 June 2006)*

### FROM THE CHAIR OF THE NOMENCLATURE COMMITTEE

#### Dr. Jutta Zipfel

Since 2005, the president of the Meteoritical Society has appointed seven new members, among whom are G. Benedix, H. Chennaoui, H. Connolly, W. Hsu, C. Smith, and A. Yamaguchi. I welcome these new members and wish them successful years on the committee. I also thank members that rotated off the committee during this time, R. Harvey, M. Kimura, D. Kring, Y. Lin, and H. Palme, for their time and effort spent working on the committee. On behalf of the entire Committee I extend special thanks to S. Russell and M. Grady, who also rotated off the committee and could not continue their work as editors of the Meteoritical Bulletin. When Sara and Monica took over in 2002, the number of non-Antarctic meteorites reported was 493. Soon after, in 2004, it reached a maximum of 857. We all owe them our highest respect and gratitude for keeping the system functioning through this difficult time.

During the last couple of years, the Nomenclature Committee of the Meteoritical Society experienced a number of significant changes. Harold Connolly was appointed new editor of the Meteoritical Bulletin in May 2005. At the same time, I took over the chair of the Nomenclature Committee from Jeff Grossman. During the first months, Jeff was of tremendous support in getting the committee, and especially

Harold and me, started. We had many discussions about what changes could make the system more efficient and how to implement these changes so that they were relatively transparent to submitters, associated editors, and the Nomenclature Committee members.

Harold worked hard to structure the editorship in a way that permitted the new goals to be achieved. Major changes include: 1) an increased number of associate editors (Caroline Smith and Gretchen Benedix were appointed as associate editors for meteorites from Northwest Africa and Akira Yamaguchi for meteorites from Asia and the Pacific); 2) a regular review and voting schedule, the dates of which are announced on our Web page; 3) templates for tables and short descriptions; 4) submissions through the editor only. Also, he has given this and future editions of the Meteoritical Bulletin a new look. Meteorites are listed for each continent separately. Single descriptions are divided into history, physical properties, petrography, geochemistry, classification, and type specimen. He has also added images and will invite specified classifiers to submit images of described meteorites for future bulletins. In addition, starting in 2006 the Meteoritical Bulletin will be published twice a year.

Jeff Grossman has created and installed a meteorite database and search engine on the society's website, [http://www.meteoriticalsociety.org/simple\\_template.cfm?code=pub\\_bulletin](http://www.meteoriticalsociety.org/simple_template.cfm?code=pub_bulletin). He will regularly update his database as it provides information on the publication status of a given meteorite.

This is especially important to authors of papers and abstracts because editors of various journals, including *Meteoritics & Planetary Science* and *Geochimica et Cosmochimica Acta*, will no longer publish papers that contain provisional or unofficial meteorite names. I encourage everyone working in the field of meteoritics to use our new online database. On behalf of the entire committee, we are most grateful to Jeff for his hard work in creating and maintaining this online service for all communities.

During the last two years it has been tremendously helpful to rely on the support of Herbert Palme, president of the Meteoritical Society. President Palme regards the Nomenclature Committee as currently the most important committee among all committees of the Meteoritical Society. With his support, we have finally brought about new structures to serve the needs of the various meteorite communities. In part, the increasing number of newly recovered and classified meteorites motivated the committee to make many of these changes. I would like to stress that meeting the needs of the various meteorite communities may require additional changes in the coming year. It is my aspiration that all meteorite communities—scientists, classifiers, collectors, dealers, and all interested in meteorites—will benefit equally from these.

#### FROM THE EDITOR

##### Harold C. Connolly, Jr.

In February 2006 I attended the Tucson Gem and Mineral Show, held in the great city of Tucson, Arizona, USA. It was an amazing learning experience. For the first time I met many dealers, collectors, and classifiers from around the world. I had the great pleasure of meeting and discussing many issues surrounding meteoritics with Ms. A. Black, Mr. M. Taylor, Mr. E. Olson, Mr. J. Scharder, Mr. Martin Horejsi, Mr. M. Farmer, Mr. R. Verish, Mr. G. Heslep, Mr. R. Wesel, M. B. Fectay, Mme. C. Bidaut, and many others. I attended Mr. M. Blood's auction, which was great fun and a wonderful learning experience. Some friends and colleagues were also in attendance such as Drs. A. Ruzicka and M. Hutson, Dr. W. Hsu and his wife, and Dr. A. Ehlmann and his wife. I would like to thank the director of the Southwest Meteorite Center, Dr. D. Lauretta and the curator of the center, Mr. M. Killgore, for their invitation to attend the show and to speak about nomenclature to the many meteorite communities attending the show. Also speaking were L. Welzenback from the Smithsonian Institution, who spoke on meteorite classification, and M. Killgore, who spoke about the Southwest Meteorite Center. Dr. Lauretta and Curator Killgore were wonderful hosts and really aided me with a common goal: building bridges between the many meteorite communities. The increasing, almost alarming, rate of meteorite recovery from around the world is and will continue to drive new thinking on issues related to meteorite science,

nomenclature, collecting, and dealing. I look forward to future visits to the Tucson show, seeing the many new friends and colleagues.

##### Noteworthy Recovery

The office of the editor, the entire Nomenclature Committee, and all of the Meteoritical Society congratulate Steve Arnold of Arkansas, USA, Phil Mani of Texas, USA, and their colleagues for the recovery of a whopping 650 kg of Brenham!

##### The New Look of the Meteoritical Bulletin

With this edition, the presentation of the Meteoritical Bulletin changes. We hope to continue to make changes, as we feel they are needed to meet our goals of better serving the communities. This edition presents the descriptions or basic classification information for over 400 meteorites, 99% of which are new recoveries, either finds or falls. Keeping with our goals of presenting consistent data tables and descriptions, some meteorites typically published in the Meteoritical Bulletin, such as those from the United States of America's Antarctic collection, will be published in the March 2007 edition. Due to a low number of meteorites recovered from Asia, Europe, and other planets, these are presented only in descriptive format.

In closing, I need to thank many people who help to produce this volume and who were supportive in the last year when many issues surrounding the Meteoritical Bulletin were in transition. It is critical to recognize the amazing team of associate editors. They are co-authors of the Meteoritical Bulletin, and they serve the society by performing what is often tedious work. This past year, they all worked extremely hard to improve communication between the Nomenclature Committee, the Meteoritical Bulletin, submitters, and classifiers, and I deeply thank them. I am very appreciative of Tim Jull, the editor of *Meteoritics & Planetary Science*, Agnieszka Baier and everyone else at the production office, for their support in transforming the Meteoritical Bulletin to its new format.

#### AFRICA

##### Specific Locations within Africa

##### Acfer 366

26°36'56"N, 03°56'14"E

Acfer region, Libya

Find: November 2002

Carbonaceous chondrite (CH3)

**History:** A single stone weighing 1456 g was found in the Acfer region of Libya by Filiberto Ercolani (deceased 2004).

**Petrography:** (V. Moggi-Cecchi, A. Salvadori, and G. Pratesi, *MSP*) The outer surface displays a small portion of fusion crust. The stone is composed of 60 vol% lithic and crystal fragments, 20 vol% chondrules, and 30 vol%

nonsilicate phases. Chondrules range from 30–300  $\mu\text{m}$  with a mean apparent diameter of 90  $\mu\text{m}$ . Chondrule textural types: cryptocrystalline (C) = 58%, granular olivine (GO) + granular olivine-pyroxene (GOP) = 22%, porphyritic olivine (PO) = 17%, and barred olivine (BO) = 3% (all vol%). Most of the larger chondrules are GO/GOP. Nonsilicate phases are mainly Fe,Ni metal (predominantly kamacite) and minor sulfides. The metal and sulfide grains range from 10–100  $\mu\text{m}$ , are rounded to irregular in texture, and are homogeneously distributed through the meteorite, within and outside of chondrules.

**Geochemistry:** EPMA of chondrule olivine indicate a fairly homogeneous composition ( $\text{Fo}_{97-100}$ ; mean  $\text{Fo}_{97}$ ) although olivine in fragments has a more variable composition ( $\text{Fo}_{80-99}$ ; mean  $\text{Fo}_{95}$ ). Low-Ca pyroxene in chondrules and fragments is relatively homogenous ( $\text{En}_{93-100}\text{Wo}_{1-5}$ ). High-Ca pyroxene is rare, occurring only in PO chondrules ( $\text{Fs}_{23.8}\text{Wo}_{31.9}$ ). Plagioclase is found in chondrules and as angular fragments, and ranges from  $\text{An}_{80}$  to  $\text{An}_{100}$ . Kamacite has a high Ni content (5–9 wt%). Sulfides are typically troilite (Ni = 1–2 wt%) although some pyrrhotite grains with Ni = 11 wt% are observed. Oxygen isotope: (I. Franchi and R. Greenwood, *OU*)  $\delta^{17}\text{O} = 1.781$ ,  $\delta^{18}\text{O} = 4.610$ ,  $\Delta^{17}\text{O} = -0.616$  (all ‰).

**Classification:** Carbonaceous chondrite (CH3); S1, moderate weathering.

**Specimens:** A total of 30 g type specimen, one thin section, and the main mass of 1410 g are on deposit at *MSP* (inventory number MSP 2273).

#### Acfer 374

26°36'52"N, 04°03'18"E

Acfer region, Libya

Find: November 2002

Carbonaceous chondrite (CO3)

**History:** Seven small fragments, weighing 118 g in total, were found in the Acfer area of Libya by an Italian dealer. The largest mass weighs 30 g.

**Physical Characteristics:** (V. Moggi-Cecchi, A. Salvadori, and G. Pratesi, *MSP*) The hand specimen has an outer surface that is brown-red while the interior is darker.

**Petrography:** Chondrules range in size from 30 to 450  $\mu\text{m}$  (mean 110  $\mu\text{m}$ ). Chondrules textural types: GO to GOP = 61%, PO = 17%, GOP = 12%, radial pyroxene (RP) = 4%, C = 4%, POP = 1%, and BO = 1% (all vol%). Matrix/chondrules ratio is ~0.5. Chondrules are set in a very fine-grained matrix composed of olivine, phyllosilicates, and pyroxene. Calcium-rich, aluminum-rich inclusions (CAIs), and amoeboid olivine inclusions (AOIs) account for about 10 vol% of the meteorite. Opaque phases (6–8 vol%) include Fe,Ni alloys, troilite, pentlandite, and pyrrhotite. Rare grains of awaruite ( $\text{Ni}_2\text{Fe}$ ), tetrataenite ( $\text{FeNi}$ ), schreibersite, and nickel phosphide are present.

**Geochemistry:** (SEM and EPMA) Olivine composition is variable ( $\text{Fo}_{55-99}$ , mean  $\text{Fo}_{94}$ ,  $n = 26$  [ $n$  refers to the number of

analyses performed]), with narrow compositional ranges in type I chondrules, mainly GO ( $\text{Fo}_{95-100}$ , mean  $\text{Fo}_{97.1}$ ) and wider in type II ( $\text{Fo}_{55-100}$ ). Mean  $\text{Cr}_2\text{O}_3$  content in 10 MgO-poor olivines ( $\text{FeO} > 2$  wt%) is 0.35 wt%. Olivine in AOIs is Fe-rich,  $\text{Fo}_{96-100}$ . Low-Ca pyroxene is predominantly enstatite ( $\text{En}_{85-100}$ ) with different mean values for crystals in and out of chondrules ( $\text{En}_{80.4}$  and  $\text{En}_{96.7}$ , respectively). POP chondrules have high-Ca pyroxenes ( $\text{En}_{62.43}\text{Wo}_{36.58}\text{Fs}_{0.90}$ ). Kamacite (Ni = 4–5.00, mean value 4.61 [both wt%]; Co = 0.1–1.00, mean value 0.16 [both wt%]). Oxygen isotope: (I. Franchi and R. Greenwood, *OU*)  $\delta^{17}\text{O} = -6.042$ ,  $\delta^{18}\text{O} = -2.306$ ,  $\Delta^{17}\text{O} = -4.843$  (all ‰).

**Classification:** Carbonaceous chondrite (CO3); S2, minimal weathering.

**Specimens:** All the fragments, two polished thin sections, and the main mass of 30 g are on deposit at the *MSP* (inventory number MSP 2283).

#### Hammadah al Hamra 337

29°00'00"N, 12°07'40"E

Hammadah al Hamra region, Libya

Find: 24 February 2001

Carbonaceous chondrite (CK4)

**History:** A single stone weighing 198 g was found in February 2001 in the Hammadah al Hamra region of Libya by Giovanni Pratesi (*MSP*) during a scientific expedition.

**Physical Characteristics:** (V. Moggi-Cecchi, A. Salvadori, and G. Pratesi, *MSP*) The main mass has a dark brown external surface with fusion crust in some areas. In hand sample the chondrules are set in a dark green matrix and CAIs (up to 1 mm) are present.

**Petrography:** The thin section contains few chondrules (not perfectly delineated and sometimes altered) set in fine-grained matrix with several coarser-grained olivine crystals. Chondrules range from 380 to 1800  $\mu\text{m}$  (mean value 700  $\mu\text{m}$  on  $n = 25$ ) and account for about 20% of the total volume. Chondrule textural types: POP = 72%, PO = 12%, and GOP = 16% (all vol%). Very rare and extremely altered AOIs (2 vol%) and abundant CAIs (~1 mm) are observed. Plagioclase crystals can be found in GO and PO chondrules. Metal alloys are extremely rare (<0.01 vol%). Magnetite (inside and outside chondrules) is abundant (~8 vol%). Sulfides (mainly pyrrhotite) are rare and are located in the matrix. Rare kamacite and moncheite ( $\text{Pt,Pd}(\text{Te,Bi})_2$ ) are also present.

**Chemistry:** (EPMA) Olivine compositions are quite homogeneous ( $\text{Fo}_{68-75}$ ; mean  $\text{Fo}_{74}$ ), with the exception of rare zoned olivine crystals ( $\text{Fo}_{68-93}$ ) in PO chondrules. Low-Ca pyroxenes in GOP chondrules are  $\text{En}_{90-100}$ . High-Ca pyroxenes in POP chondrules have an augitic composition ( $\text{Wo}_{50}\text{En}_{50-60}\text{Fs}_{0-10}$ ). Plagioclase and devitrified mesostasis is albitic ( $\text{An}_{25-50}$ ). Magnetite ( $\text{Cr}_2\text{O}_3 = 2-3$  wt%): Sulfides (Ni = 1–2 wt%); kamacite with low Ni. Oxygen isotope: (I. Franchi, R. Greenwood, *OU*)  $\delta^{17}\text{O} = -3.668$ ,  $\delta^{18}\text{O} = -0.369$ ,  $\Delta^{17}\text{O} = -3.496$  (all ‰).

**Classification:** Carbonaceous chondrite (CK4); S1, moderate to extensive weathering.

**Specimens:** The main mass (174 g), a 24 g type specimen, and two thin sections are on deposit at *MSP* (inventory number MSP 1592).

**Mafuta** **16°54'09"S, 30°24'26"E**

Mafuta Farm, Makonde District, Zimbabwe

Find: 1 December 1984

Iron (IID)

**History:** On 1 December 1984, while clearing rocks from his fields with an assistant, Mr. Murray Alexander found an iron meteorite on Mafuta Farm, Makonde District, northern Zimbabwe. In 1993, Mr. Alexander agreed to have a slice cut for scientific studies. The cutting was done in Harare and a slice (about  $9.5 \times 4.5 \times 1$  cm) was sent to Vienna for further studies.

**Physical Characteristics:** (C. Koeberl, *UVien*) The original mass and dimensions were 71.5 kg and  $40 \times 20$  cm, respectively. The exterior has little rust and a brownish color.

**Petrography:** A medium octahedrite (kamacite band width between 0.4 and 1.1 mm) and rich in schreibersite (up to 5 mm), but it does not contain any large sulfide exposed on the large slice. The kamacite is recrystallized to an equigranular intergrowth of ~0.5 mm grains.

**Geochemistry:** Bulk composition: (C. Koeberl, *UVien*; INAA J. Wasson *UCLA*; PIXE W. Przybylowicz, *NACSA*) First sample: Ni = 9.71, Co = 0.61 (both wt%), Cr = 32, Ga = 67, As = 4.8, Os = 13.3, Ir = 15.0 (all ppm), and Au = 615 ppb. Second sample: Ni = 10.03, Co = 0.67 (both wt%), Cr = 62, Cu = 246, Ga = 72.7, As = 5.0, W = 2.92, Ir = 15.4, Pt = 19.8 (all ppm), and Au = 645 ppb.

**Classification:** (J. Wasson, *UCLA*) Iron (IID).

**Specimens:** A total of 221 g of sample is on deposit at the *NHNV* (inventory number M6694). Mr. M. Alexander of Mafuta Farm, Zimbabwe, holds the main mass.

## Northeast Africa

### Northeast Africa 002

Possibly Libya

Find: 2004

Iron (IID, anomalous)

**History:** A 5480 g iron meteorite was purchased in 2004.

**Petrography:** (J. Wasson, *UCLA*) The meteorite consists of small domains  $0.81 \times 2$  cm, filled with a fine Widmanstätten pattern of kamacite bandlets, often in parallel clusters of several bands, each ~0.2 mm wide. No inclusions are present. Cracks are present between several of the domains. Bulk composition: Cr = 138, Co = 6.6, Ni = 102 (all mg/g); Cu = 259, Ga = 71, As = 4.2, W = 3.0, Ir = 23, and Au = 0.56 (all  $\mu\text{g/g}$ ).

**Classification:** Iron (IID) with anomalous structure. The bulk composition is similar to that of IID-an Arltunga, which has an anomalous (but still finer) structure.

**Specimens:** A 131 g full slice type specimen is on deposit at *UCLA*. John Birdsell holds the main mass.

### Northeast Africa 003

**30°28'N, 13°33'E**

Libya

Find: November 2000/December 2001

Achondrite (lunar, mare basalt and basaltic breccia)

**History:** A dark gray, 6 g stone was found by a prospector 12 km NW from Al Qaryah ash Sharqiyah in Wadi Zamzam area, Libya, in December 2001. A stone of 118 g with incomplete fusion crust was found nearby in December 2001.

**Petrography:** (J. Haloda and P. Tycova, *PCU*) The meteorite contains two adjacent parts, mare basalt and basaltic breccia, both of which were investigated in detail. The main portion (~75 vol%) of the meteorite is coarse-grained, low-Ti olivine-rich basalt, showing porphyritic texture of olivine (Fo<sub>19-73</sub>), zoned pyroxene (En<sub>5-71</sub>Wo<sub>6-38</sub>), and plagioclase (An<sub>84-92</sub>) with late-stage mesostasis containing silica, Fe-rich pyroxene and pyroxferroite, plagioclase, ilmenite, troilite, and apatite. Opaque phases include chromite, Ti-rich chromite, ulvöspinel, ilmenite, troilite, and trace Fe,Ni metal; shock veins and impact melt pockets are present. All plagioclase is totally converted to maskelynite. Mineral modes are (vol%) olivine = 17.5, pyroxene = 60.6, plagioclase = 18.2, ilmenite = 1.2, spinel = 0.8, mesostasis + impact melt = 1.8.

**Geochemistry:** Bulk composition: SiO<sub>2</sub> = 44.7, TiO<sub>2</sub> = 1.3, Cr<sub>2</sub>O<sub>3</sub> = 0.5, Al<sub>2</sub>O<sub>3</sub> = 8.0, FeO = 21.8, MnO = 0.3, MgO = 13.6, CaO = 9.2, Na<sub>2</sub>O = 0.3, K<sub>2</sub>O = 0.1 (all wt%), Fe/Mn = 81. Concentration of selected elements (INAA; R. Korotev, *WashU*) Sc = 50.8, Co = 50.5, Ni = 84, Hf = 1.1, Ta = 0.15, Th = 0.43, U = 0.2 (all ppm); REE pattern is flat at  $10 \times \text{CI}$  with slight negative Eu anomaly, not depleted in LREE. Adjacent part is basaltic breccia (~25 vol%) consisting of well-consolidated glassy impact-melt matrix containing scattered mineral fragments of chemical composition identical with the coarse-grained low-Ti olivine-rich basalt and two larger clasts of low-Ti mare basalt lithologies. The low-Ti basaltic clasts are finer-grained and petrologically more evolved. No regolith component or highland material is present.

**Classification:** Achondrite (lunar mare basalt and basaltic breccia).

**Specimens:** A 20 g type specimen and two polished thin sections are on deposit at *PCU*. An anonymous finder holds the main mass.

## Northwest Africa

### Northwest Africa 869

Northwest Africa

Find: 2000 or 2001

Ordinary chondrite (L4-6)

**History:** It is quite clear that meteorite collectors in Northwest Africa have discovered a large L chondrite strewn

field at an undisclosed location. At least 2 metric tons of material comprising thousands of individuals has been sold under the name NWA 869 in the market places of Morocco and around the world. Individual masses are known to range from <1 g to >20 kg. It is certain that NWA 869 is paired with other NWA meteorites, although no systematic survey has been done. It is also possible that some stones sold as NWA 869 are not part of the same fall, although dealers are confident that most of the known masses are sufficiently distinctive from other NWA meteorites in terms of surface and internal appearance that the error rate should be fairly low. Scientists are advised to confirm the classification of any specimens they obtain before publishing results under this name.

**Petrography and Geochemistry:** (A. Rubin, *UCLA*) A fragmental breccia of type 4–6 material; one thin section dominated by an L5 lithology gave olivine (Fa<sub>24.2</sub>). **Classification:** Ordinary chondrite (L4–6); W1, S3. **Specimens:** A 189.3 g type specimen is on deposit at *UCLA*.

#### Northwest Africa 999

Morocco

Find: 2000

Achondrite (eucrite)

**History:** A 330 g brecciated stone was purchased at the Tucson show by D. Gregory from an anonymous Moroccan dealer.

**Petrology:** (P. Warren, *UCLA*) An extremely fine-grained (0.01–0.1 mm) stone with thoroughly exsolved pyroxenes, some pyroxenes (<1 vol%) up to 0.6 mm in maximum dimension. Microphenocrysts are compositionally identical to the groundmass pyroxenes.

**Geochemistry:** Low-Ca pyroxene (En<sub>32.0</sub>Wo<sub>2.3</sub>); high-Ca pyroxene (En<sub>27.8</sub>Wo<sub>44.1</sub>). Bulk chemistry: (INAA, *UCLA* two chips) Al<sub>2</sub>O<sub>3</sub> = 12.5 wt%, Na = 2.83 and 2.84 (both wt%); Sc = 32 and 33; Sm = 2.1 and 2.2 (both µg/g); Fe/Mn = 36, 37.

**Classification:** Achondrite (eucrite); low to moderate shock.

**Specimens:** A 20 g type specimen is on deposit at *UCLA*. Gregory holds the main mass.

#### Northwest Africa 1006

Morocco

Find: 2001

Achondrite (ureilite)

**History:** On 27 August 2001, M. Farmer purchased a 24.5 g sample in Quarzazate, Morocco.

**Petrography:** (P. Warren, *UCLA*) The meteorite is mostly composed of olivine and pigeonite with minor orthopyroxene. The Px/(Px+Ol) ratio is ~0.4. Olivines have rims with reduced chemistry.

**Geochemistry:** Olivine (cores Fo<sub>89.6</sub>; CaO = 0.33; Cr<sub>2</sub>O<sub>3</sub> = 0.64 [both wt%; *n* = 47]), pigeonite (En<sub>82.8</sub>Wo<sub>7.9</sub>; Cr<sub>2</sub>O<sub>3</sub> = 0.99, Al<sub>2</sub>O<sub>3</sub> = 0.57, MnO = 0.47 [all wt%; *n* = 17]), orthopyroxene (En<sub>86.4</sub>Wo<sub>4.7</sub>; Cr<sub>2</sub>O<sub>3</sub> = 0.95, Al<sub>2</sub>O<sub>3</sub> = 0.62 [both wt%; *n* = 3; all analyses by EMP]). Based on INAA data

(*UCLA*), typical ureilite trace-element composition (except for weathering effects): Sc = 9.4, Ni = 720, Zn = 330 (all µg/g); Ir = 124, Au = 16.1 (both ng/g).

**Classification:** Achondrite (ureilite); low shock.

**Specimens:** A 5.86 g type specimen is on deposit at *UCLA*. M. Farmer holds the main mass.

#### Northwest Africa 1460

Morocco

Find: June 2002

Achondrite (Martian, basaltic shergottite)

**History:** In January 2002, A. Habibi provided several small fragments of a 70.2 g complete stone with a fresh, black fusion crust to A. and G. Hupé for analysis. N. Oakes later purchased the stone for more detailed investigation.

**Physical Characteristics:** A complete stone weighing 70.2 g with a fresh black fusion crust. Dimensions of the stone are 47 × 34 × 27 mm.

**Petrography:** (A. Irving and S. Kuehner, *UWS*) The stone is largely unweathered and coarse-grained, with large, pale yellow-green pyroxene and glassy maskelynite laths clearly visible. Texture is subophitic to intersertal.

**Geochemistry:** Pyroxenes are zoned with cores of orthopyroxene (Fs<sub>20</sub>Wo<sub>3</sub>; FeO/MnO = 30.7) mantled by augite (Fs<sub>27</sub>Wo<sub>31</sub>; FeO/MnO = 32.2) with rims of Fe-rich pigeonite (Fs<sub>55</sub>Wo<sub>18</sub> to Fs<sub>85</sub>Wo<sub>13</sub>; FeO/MnO = 41.0–36.8). Plagioclase (Ab<sub>51</sub>Or<sub>1–2</sub>, maskelynite) has patchy compositional zoning. Accessory minerals are merrillite, Cl-F-bearing apatite, exsolved Fe-Ti oxides (ilmenite lamellae in titanomagnetite), ilmenite, chromite, pyrrhotite, K-rich glass, silica, and baddeleyite (occurs as blades up to 50 µm long). Fine-grained symplectitic intergrowths composed of fayalite+silica+hedenbergite occurs at the margins of pigeonite grains. Other symplectitic intergrowths of fayalite+silica also occur, commonly at the boundaries between merrillite and pyroxene. Textures, mineralogy, and mineral compositions are essentially identical to NWA 480, with which this sample is paired. Cosmogenic isotopes: (K. Nishizumi, *SSL*) Give a cosmic ray exposure age of 2.6 ± 0.2 Ma. Radiogenic isotopes and formation age: (L. Nyquist and C.-Y. Shih, *JSC*) A preliminary Sm-Nd isochron age of 352 ± 30 Ma and a Rb-Sr isochron age of 313 ± 3 Ma. **Classification:** Achondrite (Martian, basaltic shergottite); minimal weathering.

**Specimens:** A 2.5 g type specimen is on deposit at *JSC*. A 2.6 g type specimen and one polished thin section are on deposit at *MNB*. A 5.7 g type specimen and two polished thin sections are on deposit at *UWS*. A 3.5 g type specimen is on deposit at *NAU*. An anonymous collector holds the main mass.

#### Northwest Africa 1462

Morocco

Find: 2002

Achondrite (ureilite)

**History:** On 3 March 2002, D. Gregory purchased a 203 g sample in Erfoud, Morocco.

**Petrography:** (P. Warren, *UCLA*) The meteorite is mostly composed of olivine and pigeonite. The Px/(Px+Ol) ratio is ~0.3. Olivines have rims with reduced chemistry.

**Geochemistry:** Olivine ( $\text{Fe}_{78.1}$ ;  $\text{CaO} = 0.30$ ,  $\text{Cr}_2\text{O}_3 = 0.30$  [both wt%;  $n = 10$ ]), pigeonite ( $\text{En}_{68.1}\text{Wo}_{14.4}$ ;  $\text{Cr}_2\text{O}_3 = 1.23$ ,  $\text{Al}_2\text{O}_3 = 1.93$ ,  $\text{MnO} = 0.44$  [all wt%;  $n = 12$ ]). Based on INAA data (*UCLA*), typical ureilite trace-element composition: Sc = 9.4, Ni = 900, Zn = 159 (all  $\mu\text{g/g}$ ); Ir = 360, Au = 31 (all ng/g).

**Classification:** Achondrite (ureilite); low shock.

**Specimens:** A 20.2 g type specimen is on deposit at *UCLA*. D. Gregory holds the main mass.

#### Northwest Africa 1918

Morocco

Find: January 2003

Achondrite (eucrite, basaltic)

**History:** D. Gregory purchased a complete, fusion-crust stone weighing 136 g from a Moroccan dealer in January 2003.

**Petrography:** (A. Irving and S. Kuehner, *UWS*) Unbrecciated, with subophitic texture of dark grey pigeonite grains and groups of small anorthite laths, ilmenite (with rare baddeleyite inclusions), troilite, and Ni-free metal.

**Geochemistry:** Some of the large pigeonite grains have ribbon-like clinopyroxene exsolution lamellae within host orthopyroxene ( $\text{Fs}_{61.6}\text{Wo}_{1.4}$ ;  $\text{FeO/MnO} = 33.7$ ); clinopyroxene ( $\text{Fs}_{29.6}\text{Wo}_{39.2}$ ;  $\text{FeO/MnO} = 29.2$ ) occurs also as discrete grains along with silica.

**Classification:** Achondrite (eucrite, basaltic).

**Specimens:** A 20.2 g type specimen and one polished thin section are on deposit at *UWS*. Gregory holds the main mass.

#### Northwest Africa 1923

Morocco

Find: March 2003

Achondrite (eucrite, gabbroic)

**History:** D. Gregory purchased an extremely fresh, unbrecciated stone weighing 112 g with black fusion crust from a Moroccan dealer in March 2003.

**Petrography:** (A. Irving and S. Kuehner, *UWS*) Coarse-grained, noncumulus, igneous texture with primary grain contacts. Subequal amounts anorthite and pale yellow pigeonite with minor chromite.

**Geochemistry:** Anorthite ( $\text{An}_{95}$ ); low-Ca pyroxene grains consist of orthopyroxene ( $\text{Fs}_{37.8}\text{Wo}_{2.9}$ ;  $\text{FeO/MnO} = 25.8$ ) with sparse blades of exsolved clinopyroxene ( $\text{Fs}_{16.3}\text{Wo}_{43.2}$ ;  $\text{FeO/MnO} = 23.8$ ).

**Classification:** Achondrite (eucrite, gabbroic).

**Specimens:** A 20.4 g type specimen and one polished thin section are on deposit at *UWS*. Gregory holds the main mass.

#### Northwest Africa 1929

Morocco

Find: May 2003

Achondrite (howardite)

**History:** A 922.2 g, partially crusted, complete stone was purchased in Erfoud, Morocco, in May 2003.

**Petrography:** (T. Bunch and J. Wittke, *NAU*) A breccia of 72 vol% cumulate eucrite clasts, 8% subophitic clasts, 14% diogenites, and 6% melt clasts. Pervasive solid-state recrystallization of plagioclase and pyroxenes with localized melt pockets and veins in clasts.

**Geochemistry:** Coarse-grained gabbroic eucrite: Pyroxene ( $\text{Fs}_{45-40}\text{Wo}_{7-20}$ ); plagioclase ( $\text{An}_{91.2-95.3}$ ); metal (Ni = 0.97; Cr = 0.87 [both wt%]). Diogenite: Pyroxene ( $\text{Fs}_{43-54}\text{Wo}_{2.5-3.6}$ ; Fe/Mn = 36, 37).

**Classification:** Achondrite (howardite); high shock.

**Specimens:** A 22.02 g type specimen is on deposit at *NAU*. Farmer holds the main mass.

#### Northwest Africa 2200

Morocco

Find: August 2004

Achondrite: (lunar, feldspathic breccia)

**History:** A completely crusted, 552 g, ellipsoidal stone was found in the Atlas Mountains, Morocco, and purchased in Erfoud by a Moroccan dealer for D. Gregory in August 2004.

**Petrography and Geochemistry:** (S. Kuehner and A. Irving, *UWS*) Breccia consisting of coarse, greyish-to-whitish lithic and mineral clasts in a darker glassy-to-finely crystalline matrix. Lithic clasts are mainly very fine-grained, quench-textured, feldspathic rocks that probably result from impact melting of anorthositic to gabbroic anorthositic precursors. A small percentage of the clasts are ophitic-textured mare basalts. Mineral clasts include anorthitic plagioclase, olivine ( $\text{Fa}_{30-60}$ ), exsolved pigeonite, irregular grains of metal (Ni = 10–45 wt%), Ti-rich chromite, Ti-poor chromite, pyroxene-like glass, schreibersite (Ni = 5 wt%), clinopyroxene, ilmenite, troilite, and rare zirconolite. Clinopyroxene and orthopyroxene grains in mineral and lithic clasts have Fe/(Fe+Mg) = 0.258–0.482 with Ti/(Ti+Cr) = 0.53–0.75. FeO/MnO ratios measured for olivine (99.7, 105.5), clinopyroxene (73.7), and orthopyroxene (65.4) are unmistakably within the ranges for these minerals in known lunar rocks. Feldspar grains in mineral and lithic clasts have a narrow compositional range of  $\text{An}_{95.8-97.4}$ .

**Classification:** Achondrite (lunar, feldspathic breccia).

**Specimens:** A 20.5 g type specimen, one polished thin section, and one polished mount are on deposit at *UWS*. D. Gregory holds the main mass.

#### Northwest Africa 2210

Morocco

Find: December 2004

## Carbonaceous chondrite (CH3)

**History:** D. Gregory purchased a black, magnetic stone weighing 74 g in December 2004 from D. Bessey, who had obtained this material from a Moroccan dealer in Erfoud.

**Petrography:** (A. Irving and S. Kuehner, *UWS*) Relatively fine grained (10–89  $\mu\text{m}$ ) with glass spheres ( $\text{En}_{99}\text{Fo}_{99}$ ) and chondrules (size range 10–80  $\mu\text{m}$ ), together with broken fragments of the same, in a sparse matrix.

**Geochemistry:** Abundant metal (15–20% by volume) which is mainly kamacite (5–10 wt% Ni), but in places taenite (Ni = ~20 wt%). Most of the chondrules contain very magnesian olivine ( $\text{Fa}_1$ ) and/or orthopyroxene ( $\text{Fs}_1$ ) in a mesostasis of glass, but some chondrules and fragments have more ferroan silicates (up to  $\text{Fa}_{50}$  and  $\text{Fs}_{50}$ ). Some cryptocrystalline chondrule fragments are composed of very fine intergrowths of low-Ca and high-Ca pyroxene and glass spheres have compositions intermediate between forsterite and enstatite. Other phases identified include Cr-Al-rich diopside, troilite, feldspathic glass, and rare, Al-rich chromite. No CAIs were observed. Oxygen isotopes: (D. Rumble, *CIW*) Analyses of two whole rock fragments by laser fluorination gave, respectively,  $\delta^{18}\text{O} = 3.95, 4.11$ ;  $\delta^{17}\text{O} = 1.07, 1.05$ ;  $\Delta^{17}\text{O} = -1.02, -1.12$  (all ‰).

**Classification:** Carbonaceous chondrite (CH3).

**Specimens:** A 14.8 g type specimen and one polished thin section are on deposit at *UWS*. Gregory holds the main mass.

## Northwest Africa 2225

Morocco

Find: April 2004

Achondrite (ureilite)

**History:** D. Bessy purchased one mass of 40 g in Erfoud, Morocco.

**Petrography:** (Paul Warren, *UCLA*) The stone is composed mostly of olivine and pigeonite ( $(\text{Px}/\text{Px}+\text{Ol}) = \sim 0.25$ ) with minor carbon-rich matrix.

**Geochemistry:** Olivine ( $\text{Fo}_{82.3}$ );  $\text{CaO} = 0.36$ ,  $\text{Cr}_2\text{O}_3 = 0.71$  (both wt%,  $n = 25$  cores only); pigeonite ( $\text{En}_{75.2}\text{Wo}_{9.8}$ ;  $\text{Al}_2\text{O}_3 = 0.85$ ,  $\text{Cr}_2\text{O}_3 = 1.22$ ,  $\text{MnO} = 0.41$  [all wt%,  $n = 13$  analyses]). Bulk trace elements:  $\text{Sc} = 7.0$ ,  $\text{Ni} = 1140$ ,  $\text{Zn} = 280$  (all  $\mu\text{g/g}$ );  $\text{Ir} = 231$ ,  $\text{Au} = 9.8$  (both  $\text{ng/g}$ ).

**Classification:** Achondrite (ureilite).

**Specimens:** A 8.3 g type specimen is on deposit at *UCLA*. D. Bessey hold the main mass of 20 g. N. Gessler holds 11 g.

## Northwest Africa 2646

Algeria or Morocco

Find: December 2004

Achondrite (Martian, plagioclase-olivine clinopyroxenite)

**History:** A 9.3 g, broken stone (allegedly part of a larger specimen found at an unknown site in Algeria or Morocco) was purchased from a Moroccan dealer in December 2004 for N. Oakes.

**Petrography and Geochemistry:** (T. Bunch and J. Wittke, *NAU*; A. Irving and S. Kuehner, *UWS*) The specimen is

relatively coarse-grained and has an overall gray color, but grain surfaces have a thin whitish coating. It is modally heterogeneous and is composed of ~40.7% pigeonite, ~24.3% augite, ~21.6% olivine, and ~11.4% maskelynite, with 2% chromite, ilmenite, merrillite, and pyrrhotite. Euhedral to subhedral chromite ( $\text{Cr}/(\text{Cr}+\text{Al}) = 0.869$ ,  $\text{Mg}/(\text{Mg}+\text{Fe}) = 0.12$ ) and augite ( $\text{Fs}_{19-23}\text{Wo}_{26-36}$ ;  $\text{FeO}/\text{MnO} = 22-27$ ;  $\text{Cr}_2\text{O}_3 = 0.6$ ,  $\text{Al}_2\text{O}_3 = 2.2$  [both wt%]) appear to have crystallized first, followed by subhedral to anhedral olivine ( $\text{Fa}_{38-44}$ ;  $\text{FeO}/\text{MnO} = 35-56$ ). These are poikilitically enclosed in large oikocrysts of pigeonite ( $\text{Fs}_{24.4}\text{Wo}_{5.7}$  zoned to  $\text{Fs}_{34.4}\text{Wo}_{12.1}$ ;  $\text{FeO}/\text{MnO} = 26-32$ ); augite and olivine chadacrysts tend to be clustered. Laths of plagioclase (now maskelynite,  $\text{An}_{58.4-60.7}\text{Or}_{0.9}$ ), some in clusters, occur interstitially to pigeonite grains. All maskelynite grains are rimmed by zones up to 40  $\mu\text{m}$  in width consisting of a mixture of fine grained calcite, hydrous Al-rich silicate (possibly kaolinite), and very minor calcium chloride, which appear to be replacing the primary feldspar. This specimen has affinities to the six recognized “lherzolitic shergottites,” but differs from them in having a higher abundance of plagioclase and more ferroan mafic mineral compositions.

**Classification:** Achondrite (Martian, plagioclase-olivine clinopyroxenite); minimal weathering.

**Specimens:** A 3.4 g type specimen and one polished thin section are on deposit at *NAU*. One polished mount is on deposit at *UWS*. N. Oakes holds the main mass.

## Northwest Africa 2656

Morocco or Algeria

Find: 2003

Achondrite (acapulcoite)

**History:** A 386 g broken stone with weathered fusion crust (part of a larger ~7.5 kg mass found in 2003) was purchased in Erfoud, Morocco, in 2004 for N. Oakes.

**Petrography:** (T. Bunch and J. Wittke, *NAU*; A. Irving, *UWS*) The specimen is recrystallized into homogeneous polygonal and subhedral grains with a grain size of <1 mm and a ~ equal distribution of phases.

**Composition:** Olivine ( $\text{Fa}_{8.0}$ ;  $\text{FeO}/\text{MnO} = 16-19$  [ $n = 25$ ]), orthopyroxene ( $\text{Fs}_{8.4}\text{Wo}_{2.4}$ ; [ $n = 17$ ]), plagioclase ( $\text{An}_{18.3-21.0}\text{Or}_{2.8-3.4}$ ), chromite ( $(\text{Cr}/\text{Cr}+\text{Al}) = 0.85$ ;  $\text{Mg}/(\text{Mg}+\text{Fe}) = 0.41$ ). Troilite, schreibersite, and kamacite are also present. Oxygen isotopes: (D. Rumble, *CIW*) Replicate analyses by laser fluorination gave  $\delta^{17}\text{O} = 1.71, 1.69$ ;  $\delta^{18}\text{O} = 5.05, 5.04$ ;  $\Delta^{17}\text{O} = -0.953, -0.973$  (all ‰).

**Classification:** Achondrite (acapulcoite); low shock, moderate weathering.

**Specimens:** A 21 g type specimen and one polished thin section are on deposit at *NAU*. Oakes holds the main mass.

## Northwest Africa 2700

Morocco

Find: 2004

Achondrite (lunar, olivine gabbro with regolith breccia)

**History:** A light green to dark complete stone of 31.7 g was purchased in Erfoud, Morocco, in November 2004.

**Petrography and Geochemistry:** (T. Bunch and J. Wittke, *NAU*) The specimen consists of olivine gabbro and regolith breccia lithologies. The cumulate olivine gabbro contains ~50 vol% olivine ( $\text{Fa}_{29.3-34.7}$ ;  $\text{FeO/MnO} = 94$ ), pigeonite ( $\text{Fs}_{22-28.3}\text{Wo}_{5.6-10}$ ;  $\text{FeO/MnO} = 52$ ), augite ( $\text{Fs}_{13.2}\text{Wo}_{38.5}$ ), plagioclase and minor maskelynite ( $\text{An}_{89}$ ), Ba-rich alkali feldspar ( $\text{Or}_{92}\text{An}_4$ ;  $\text{BaO} = 8.9$  wt%), Cr-spinel, ilmenite, phosphate, and troilite. The breccia lithology is dominated by small olivine gabbro fragments and also contains subvolcanic basalt clasts with zoned pyroxenes ( $\text{Fs}_{44}\text{Wo}_{29}$  to  $\text{Fs}_{58}\text{Wo}_{23}$ ;  $\text{FeO/MnO} = 57$ ); plagioclase ( $\text{An}_{90}$ ); ilmenite, and Fe-rich, low-Ca pyroxene ( $\text{Fs}_{80.8}\text{Wo}_{14}$ ); symplectites of fayalite ( $\text{Fa}_{91}$ ); hedenbergite ( $\text{Fs}_{60.3}\text{Wo}_{32.7}$ ); silica; clear to yellow glass spherules; agglutinates, high silica fayalitic rocks ( $\text{Fa}_{95.6}$ ); ulvöspinel;  $\text{K}_2\text{O}$ -rich glass ( $\text{K}_2\text{O} = 8.8$ ;  $\text{SiO}_2 = 77$  [both wt%]), and pure  $\text{SiO}_2$ .

**Classification:** Achondrite (lunar, olivine gabbro with regolith breccia) where the olivine gabbro is moderately shocked and minimally weathered. Note: This sample may be paired with NWA 773.

**Specimens:** A 6.8 g type specimen and two thin sections are on deposit at *NAU*. An anonymous finder holds the main mass.

#### Northwest Africa 2703

Morocco

Find: 2004

Achondrite (ureilite)

**History:** A complete partially crusted stone weighing 121 g was purchased in Erfoud, Morocco.

**Petrography:** (J. Wittke and T. Bunch, *NAU*): A fine-grained ( $\leq 1$  mm) ureilite that has a low level of reduction.

**Composition:** Olivine (cores,  $\text{Fa}_{12.3}$ ;  $\text{FeO/MnO} = 22$ ), pigeonite ( $\text{Fs}_{10.6}\text{Wo}_{5.1}$ ;  $\text{FeO/MnO} = 14$ ), clinopyroxene ( $\text{Fs}_{6.5}\text{Wo}_{37}$ ;  $\text{FeO/MnO} = 13$ ). Metal, sulfide, and graphite ribbons also present.

**Classification:** Achondrite (ureilite); low shock; minimal weathering.

**Specimens:** A 20.5 g type specimen is on deposit at *NAU*. G. Hupé holds the main mass.

#### Northwest Africa 2705

Morocco

Find: 2004

Achondrite (ureilite)

**History:** A complete 70 g stone was purchased in Erfoud, Morocco.

**Petrography:** (J. Wittke and T. Bunch, *NAU*) Specimen shows well-formed, basal deformation lamellae in olivine with small enclaves of slightly rotated domains. The olivines and pyroxenes contain a large amount of fine-grained graphite inclusions. Olivine margins show limited reduction. Modal analyses of silicates (vol%): Olivine = 95, pigeonite = 3, orthopyroxene = 2.

**Composition:** Olivine (cores,  $\text{Fa}_{22.3}$ ;  $\text{FeO/MnO} = 40-47$ ), pigeonite (small grains,  $\text{Fs}_{7.8}\text{Wo}_{7.7}$ ;  $\text{FeO/MnO} = 8$ ), orthopyroxene ( $\text{Fs}_{3.5}\text{Wo}_{2.7}$ ).

**Classification:** Achondrite (ureilite); moderate shock, minimal weathering.

**Specimens:** A 16.8 g type specimen is on deposit at *NAU*. G. Hupé holds the main mass.

#### Northwest Africa 2708

Morocco

Find: 2004

Carbonaceous chondrite (CK4)

**History:** A single partially crusted stone of 528 g was purchased in Erfoud, Morocco, in January 2005.

**Petrography:** (J. Wittke and T. Bunch, *NAU*) Specimen contains rare, partially altered CAI; melt clasts and chondrules with fairly sharp margin/matrix boundaries.

**Geochemistry:** Matrix olivine ( $\text{Fa}_{29.7\pm 2.7}$ ), magnetite ( $\text{Cr}_2\text{O}_3 = 4.6$ ,  $\text{Al}_2\text{O}_3 = 2.19$  [both wt%]), poorly crystallized plagioclase ( $\text{An}_{78.4}$ ), pentlandite ( $\text{Co} = 1.87$  wt%). Chondrules: Olivine ( $\text{Fa}_{30.6\pm 1.95}$ ), Ca-rich pyroxene ( $\text{Fs}_{43.5}\text{Wo}_{48.7}$ ), magnetite ( $\text{Cr}_2\text{O}_3 = 3.2$ ,  $\text{Al}_2\text{O}_3 = 1.0$  [both wt%]). Magnetite in melt clasts contains Ni-bearing, exsolved ilmenite.

**Classification:** Carbonaceous chondrite (CK4); host = low shock; clasts = moderate to highly shocked, moderate weathering.

**Specimens:** A 20.3 g type specimen is on deposit at *NAU*. G. Hupé holds the main mass.

#### Northwest Africa 2711

Morocco

Find: 2004

Mesosiderite

**History:** A complete stone of 433 g was purchased in Erfoud, Morocco.

**Petrography:** (J. Wittke and T. Bunch, *NAU*) The specimen contains a relatively even distribution of metal and silicates with coarse orthopyroxene grains with margins that are recrystallized and contain numerous merrillite inclusions. The matrix is recrystallized. Modal content (vol%): Orthopyroxene = 66, metal = 16, plagioclase = 10, Ca pyroxene = 4, merrillite = 2, chromite and sulfide = 2.

**Composition:** Orthopyroxene ( $\text{Fs}_{26.8}\text{Wo}_{3.1}$ ;  $\text{FeO/MnO} = 31$ ); plagioclase ( $\text{An}_{92.9}$ ).

**Classification:** Achondrite (mesosiderite); low shock, minimal weathering.

**Specimens:** A 24 g type specimen is on deposit at *NAU*. G. Hupé holds the main mass.

#### Northwest Africa 2724

Northwest Africa

Find: 2004

Achondrite (eucrite, polymict)

**History:** A 3804 g crusted complete stone was purchased in Erfoud, Morocco, in 2004.



**Petrography:** (J. Wittke and T. Bunch, *NAU*) The specimen mostly consists of cumulate basalt clasts that range from coarse-grained (~2 mm) to very fine-grained (~0.1 mm) with less than 5 vol% of subophitic basalts observed. Cumulate basaltic clasts are unusually low in ilmenite, chromite, and metal compared with other cumulate eucrites.

**Composition:** Despite the large range in grain size, pyroxene compositions show a rather narrow range: low-Ca pyroxenes ( $\text{Fs}_{59.1-62}\text{Wo}_{3.2-3.8}$ ), pigeonite ( $\text{Fs}_{46.5-51}\text{Wo}_{12.4-16.6}$ ), Ca-rich pyroxenes ( $\text{Fs}_{40-2.1}\text{Wo}_{30.5-40.5}$ ), plagioclase ( $\text{An}_{89.4-92.7}$ ).

**Classification:** Achondrite (eucrite, polymict); low shock, minimal weathering.

**Specimens:** A 34 g type specimen is on deposit at *NAU*. E. Thompson holds the main mass.

#### Northwest Africa 2727

Morocco or Algeria

Find: June/July 2005

Achondrite (lunar, mare basalt/gabbro breccia)

**History:** Four stones of 30.6 g, 11.6 g, 64 g, and 85 g were purchased from Moroccan dealers in Erfoud for a consortium of North American collectors in June and July of 2005.

**Petrography and Geochemistry:** (T. Bunch and J. Wittke, *NAU*; A. Irving and S. Kuehner, *UWS*; R. Korotev, *WUSL*) All stones are very similar and consist of clast-dominated polymict breccias composed of >80 vol% olivine-phyric basalt and gabbroic/diabasic clasts (0.2 cm to several cm across) within a finer breccia matrix. The basalt clasts show a wide range in mineral compositions, but all contain phenocrysts of olivine ( $\text{Fa}_{28-99}$ ;  $\text{FeO/MnO} = 98.9$ ) and some also have phenocrysts of pyroxferroite or chromite all in a fine-grained matrix consisting of intergrown pigeonite, pyroxferroite, K-Ba feldspar, ilmenite, merrillite, baddeleyite, troilite, silica, and glass. The gabbroic clasts range in texture from coarser-grained (>3 mm) hypidiomorphic gabbro to finer-grained (~1 mm) diabasic clasts. Both types of gabbroic lithologies consist mainly of pigeonite ( $\text{Fs}_{23.3-31.3}\text{Wo}_{8.7-11.5}$ ;  $\text{FeO/MnO} = 60-69$ ) and subhedral to anhedral olivine ( $\text{Fa}_{34.1-41}$ ;  $\text{FeO/MnO} = 85-99$ ) with less abundant augite ( $\text{Fs}_{24.1-47.5}\text{Wo}_{24.4-32.1}$ ) and partly maskelite, blocky to tabular plagioclase ( $\text{An}_{81-94}$ ). The breccia matrix consists mainly of gabbroic debris with fragments of basalt, silica polymorph, symplectites, subparallel intergrowths of anorthite + pyroxferroite + ilmenite and shock-melted material. Bulk compositions: (R. Korotev, *WUSL*) INAA of 11 subsamples show that they vary considerably in bulk composition, with the most Fe-rich subsample nearly indistinguishable from NWA 3160 basalt. All other subsamples are compositionally equivalent to mixtures of NWA 3160 basalt and the regolith breccia lithology of NWA 773, but with slightly lower concentrations of incompatible elements. Note: Based on petrography, mineral compositions, and bulk compositions, these stones are paired with NWA 3160 and may be paired with NWA 773.

**Classification:** Achondrite (lunar, mare basalt/gabbro breccia).

**Specimens:** A 20.2 g type specimen and two polished thin sections are on deposit at *NAU*. A 0.5 g type specimen is on deposit at *WUSL*. Oakes, Reed, Boswell, and Turecki hold the main masses.

#### Northwest Africa 2736

Northwest Africa

Find: September 2004

Achondrite (aubrite)

**History:** M. Killgore purchased a 171.51 g stone in September 2004.

**Petrography:** (D. Hill, K. Domanik, and J. Lowe, *UAz*; I. Franchi, *OU*) The meteorite is made up primarily of enstatite (~60 vol%) and plagioclase (30 vol%) that are 50–100  $\mu\text{m}$  in size with an equigranular igneous texture. The groundmass is cross-cut by an extensive network of (oxidized) kamacite and daubreelite veins that follow silicate grain boundaries. An unusual feature of this meteorite is large, euhedral graphite up to 500  $\mu\text{m}$  in length within kamacite veins and interstitial to the silicates. Several round inclusions up to 2 mm in diameter that contain bars of enstatite and plagioclase are observed.

**Composition:** Enstatite ( $\text{En}_{99}\text{Fs}_0\text{Wo}_1$ ; avg. mg# = 9), plagioclase ( $\text{Ab}_{78.8}\text{An}_{15.9}\text{Or}_{5.3}$ ). Oxygen isotopes: (I. Franchi, *OU*)  $\delta^{17}\text{O} = 2.017$ ,  $\delta^{18}\text{O} = 3.732$ ,  $\Delta^{17}\text{O} = 0.076$  (all ‰).

**Classification:** Achondrite (aubrite); low shock, moderate weathering.

**Specimens:** A total of 37.14 g of type specimens are on deposit at *UAz*. M. Killgore holds the main mass.

#### Northwest Africa 2737

Morocco

Find: 2000

Achondrite (Martian, chassignite)

**History:** In August 2000, meteorite collectors discovered a stone fragmented into nine pieces (308 g, 128 g, 74 g, 47 g, 38 g, 6.4 g, 3.3 g, 2.0 g, and 4.3 g for a total mass of 611 g) in the western part of the Sahara.

**Petrography and Geochemistry:** (P. Beck, Ph. Gillet, B. Reynard, B. van de Moortele, *ENSL*; J.A. Barrat, M. Bohn, J. Cotton, *UBO*) Olivine ( $\text{Fo}_{78.2-79.1}$ ;  $\text{Mn/Fe} = 0.018$ ; ~89.6 vol%), chromite (4.6 vol%), low-Ca pyroxene ( $\text{En}_{78.5}\text{Wo}_{2.7}\text{Fs}_{18.8}$  to  $\text{En}_{76.6}\text{Wo}_{3.2}\text{Fs}_{20.2}$ ), high-Ca pyroxenes ( $\text{En}_{73.5}\text{Wo}_{8.0}\text{Fs}_{18.5}$  to  $\text{En}_{64.0}\text{Wo}_{22.1}\text{Fs}_{13.9}$ ;  $\text{Mn/Fe} 0.030$  [total of low- and high-Ca pyroxene ~4.1 vol%]), and sanidine glass (~1.6 vol%) with traces of apatite. The texture is that of a cumulate dominated by mm-size anhedral to subhedral olivine crystals, sometimes poikilitically enclosed in augite ( $\text{En}_{54.6}\text{Wo}_{32.8}\text{Fs}_{12.6}$  to  $\text{En}_{46.7}\text{Wo}_{44.1}\text{Fs}_{9.2}$ ). Oxygen isotopes: (I. Franchi, R. Greenwood, *OU*)  $\delta^{17}\text{O} = 2.40$ ,  $\delta^{18}\text{O} = 4.02$ ,  $\Delta^{17}\text{O} = 0.315$ ;  $\delta^{17}\text{O} = 2.30$ ,  $\delta^{18}\text{O} = 3.85$ ,  $\Delta^{17}\text{O} = 0.295$  (all ‰,  $n = 2$ ). Furthermore, NWA 2737 displays trace element abundances similar to Chassigny. For example, its REE

pattern resembles that of Chassigny but with a more pronounced LREE enrichment.

**Classification:** Achondrite (Martian, chassignite); highly shocked.

**Specimens:** A 20 g type specimen is on deposit at *ENSL*. B. Fectay and C. Bidaut of La mémoire de la Terre hold the main mass.

#### Northwest Africa 2758

Morocco

Find: 2004

Achondrite (eucrite, polymict breccia)

**History:** Four partial stones weighing a total of 458 g were purchased in Erfoud, Morocco, in October 2004.

**Petrography and Geochemistry:** (J. Wittke and T. Bunch, *NAU*) The sample contains an assortment of cumulate (maximum grain size = 1.8 mm), subophitic (maximum grain size = 0.2 mm), and shock-melted to crystallized basalts. Cumulate basalts: Pyroxenes ( $\text{Fs}_{49.3}\text{Wo}_{2.8}$ ;  $\text{Fs}_{38.8}\text{Wo}_{13.9}$ ;  $\text{Fs}_{48.5}\text{Wo}_{21.2}$ ;  $\text{FeO/MnO}$  for all = 31–40), plagioclase ( $\text{An}_{87}$ ); chromite ( $\text{Cr}/(\text{Cr}+\text{Al}) = 0.77$ ). Subophitic basalts: Pyroxenes ( $\text{Fs}_{39.1-45.6}\text{Wo}_{8.7-15.8}$ ;  $\text{FeO/MnO} = 30-36$ ), plagioclase ( $\text{An}_{89}$ ), chromite ( $\text{Cr}/(\text{Cr}+\text{Al}) = 0.87$ ).

**Classification:** Achondrite (eucrite, polymict breccia); low to high shock, minimal weathering.

**Specimens:** A 46 g type specimen is on deposit at *NAU*. Olsen holds the main mass.

#### Northwest Africa 2784

Morocco

Find: 2004

Achondrite (eucrite, polymict breccia)

**History:** A 141 g stone was purchased in Erfoud, Morocco, in October 2004.

**Petrography:** (T. Bunch and J. Wittke, *NAU*) A brecciated cumulate host (grain size up to 2.1 mm) with clasts of subophitic (maximum grain size, 0.12 mm) and recrystallized granular (average grain size, 0.03 mm) basalts.

**Geochemistry:** All analyzed lithologies are very similar in pyroxene and plagioclase composition. Orthopyroxenes ( $\text{Fs}_{62-64.1}\text{Wo}_{2.1-2.8}$ ), Ca-rich pyroxene ( $\text{Fs}_{27-31.1}\text{Wo}_{43.7-44.8}$ ), and plagioclase ( $\text{An}_{87.8-89.7}$ ). Pigeonite exsolution lamellae are too thin to analyze; cumulate contains Cr-rich ilmenite ( $\text{Cr}_2\text{O}_3 = 15.1 \text{ wt\%}$ ) and Cr-poor ilmenite. Other lithologies contain chromite of similar composition,  $\text{Cr}/(\text{Cr}+\text{Al}) = 0.85$ , and minor ilmenite.

**Classification:** Achondrite (eucrite, polymict breccia); low shock (no maskelynite), minimal weathering.

**Specimens:** A 25 g type specimen is on deposit at *NAU*. Birdsell holds the main mass.

#### Northwest Africa 2794

Morocco

Find: 2005

Achondrite (howardite)

**History:** A 145 g single stone with partial fusion crust was purchased in Erfoud, Morocco, in 2005.

**Petrography:** (J. Wittke and T. Bunch, *NAU*) Contains fine to medium cumulate basalt clast size (<3 mm diameter) and even finer-grained subophitic basalts (<1 mm) with diogenite fragments up to 8 mm.

**Geochemistry:** Cumulate basalt pyroxenes: Orthopyroxene ( $\text{Fs}_{49.4-64.7}\text{Wo}_{2.1-2.5}$ ), exsolved pigeonite ( $\text{Fs}_{45.1-54.4}\text{Wo}_{13.1-6.7}$ ), Ca pyroxene ( $\text{Fs}_{36.0-47.1}\text{Wo}_{19.6-24.1}$ ). All pyroxene  $\text{FeO/MnO} = 29-34$  ( $n = 27$ ). Diogenite: Orthopyroxene ( $\text{Fs}_{26.7-30}\text{Wo}_{2.2-2.5}$ ;  $\text{FeO/MnO} = 24-30$ ).

**Classification:** Achondrite (howardite); moderate weathering and low to moderate shock.

**Specimens:** A 21.4 g type specimen is on deposit at *NAU*. G. Hupé holds the main mass.

#### Northwest Africa 2795

Morocco

Find: 2005

Achondrite (diogenite)

**History:** A 329 g partial stone with moderately weathered fusion crust was purchased in Erfoud, Morocco, in 2005.

**Petrography:** (J. Wittke and T. Bunch, *NAU*) Brecciated fine to coarse grain size (<1 cm).

**Geochemistry:** Orthopyroxene ( $\text{Fs}_{26.2}\text{Wo}_{2.5}$ ;  $\text{FeO/MnO} = 29.5$ ), minor Ca-rich pyroxene ( $\text{Fs}_{18.5}\text{Wo}_{28.8}$ ), plagioclase ( $\text{An}_{87.7}$ ), chromite ( $\text{Cr}/(\text{Cr}+\text{Al}) = 0.79$ ), ilmenite, and kamacite present.

**Classification:** Achondrite (diogenite), moderately shocked (some maskelynite and high strain in orthopyroxene).

**Specimens:** A 20.4 g type specimen is on deposit at *NAU*. G. Hupé holds the main mass.

#### Northwest Africa 2853

Morocco

Find: 2005

Achondrite (howardite)

**History:** A 1006 g fully crusted stone was purchased in Erfoud, Morocco, in 2005.

**Petrography:** (T. Bunch and J. Wittke, *NAU*) Contains heterogeneously distributed orthopyroxene fragments that tend to be large (2–9 mm in diameter). Diogenite modes for three thin sections give 0, 9, and 15 vol%. Eucrites have a fine- to coarse-grained size range and consist of subophitic and cumulate textured basalts. Most pyroxenes recrystallized into small, polygonal aggregates; plagioclase melted, although original grain boundaries are preserved and crystallized into fine-grained fibrous to radiating textures.

**Geochemistry:** Host: Pigeonite ( $\text{Fs}_{45.3-51.7}\text{Wo}_{5.8-9.0}$ ;  $\text{FeO/MnO} = 29-33$ ), exsolution lamellae ( $\text{Fs}_{33.6-39.4}\text{Wo}_{19.0-25.4}$ ). Diogenite: Orthopyroxenes ( $\text{Fs}_{22.6-25.6}\text{Wo}_{1.8-2.3}$ ;  $\text{FeO/MnO} = 24-27$ ), plagioclase ( $\text{An}_{89-94.3}$ ).

**Classification:** Achondrite (howardite); high shock.

**Specimens:** A 22 g type specimen is on deposit at *NAU*. Regelman holds the main mass.

**Northwest Africa 2890**

Northwest Africa

Find: 2004

Achondrite (howardite)

**History:** An anonymous finder recovered a single stone of 132 g within the North African Sahara in 2004.

**Petrography and Geochemistry:** (A. Greshake, *MNB*) Polymict breccia with eucritic and diogenitic fragments embedded in a clastic matrix and contains dark impact melt fragments. Eucrite clasts consist mainly of frequently exsolved Ca pyroxene and plagioclase ( $\text{An}_{96.4-97.3}$ ), pigeonite ( $\text{Fs}_{37.3-53.0}\text{Wo}_{9.3-13.4}$ ), augite ( $\text{Fs}_{23.0-31.1}\text{Wo}_{39.6-43.3}$ ), with minor olivine, Mg-Al-chromite, and ilmenite. Diogenitic lithologies are dominated by orthopyroxene ( $\text{Fs}_{18.8-32.0}$ ) that are not always zoned.

**Classification:** Achondrite (howardite); low-shock, minimal weathering.

**Specimens:** A 22.1 g type specimen and one thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

**Northwest Africa 2895**

Northwest Africa

Find: 2004

Achondrite (ureilite)

**History:** A single stone of 43 g was found 2004 by an anonymous finder in the North African Sahara.

**Petrography and Geochemistry:** (A. Greshake, *MNB*) The sample is dominantly composed of large pigeonite ( $\text{Fs}_{9.9-10.8}\text{Wo}_{5.2-10.1}$ ;  $\text{Cr}_2\text{O}_3 = 1.1 \text{ wt}\%$ ) often enclosing small pyroxenes and less abundant olivines.

**Geochemistry:** Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = +3.27$ ,  $\delta^{18}\text{O} = +8.243$ ,  $\Delta^{17}\text{O} = -1.016$  (all ‰).

**Classification:** Achondrite (ureilite); low shock, moderate weathering.

**Specimens:** A 8.9 g type specimen and one thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

**Northwest Africa 2897**

Northwest Africa

Find: 2004

Rumuruti-like (R3–6)

**History:** An anonymous finder recovered one small stone of 23.3 g in 2004 in the North African Sahara.

**Petrography:** (A. Greshake, *MNB*) Olivine ( $\text{Fa}_{1.2-59.8}$ ), low-Ca pyroxene ( $\text{Fs}_{1.5-30}\text{Wo}_{0.1-4.9}$ ), augite ( $\text{Fs}_{7-17}\text{Wo}_{24.4-48.5}$ ), and plagioclase ( $\text{An}_{4.9-14}$ ).

**Geochemistry:** Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = +5.542$ ,  $\delta^{18}\text{O} = +5.06$ ,  $\Delta^{17}\text{O} = +2.91$  (all ‰), which are close to Rumuruti itself.

**Classification:** Rumuruti-like (R3–6); low shock, minimal weathering.

**Specimens:** A 5.75 g type specimen and one polished thin

section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

**Northwest Africa 2898**

Northwest Africa

Find: 2003

Ordinary chondrite (H7)

**History:** An anonymous finder recovered a single stone of 136 g within the North African Sahara in 2003.

**Petrography:** (A. Greshake, *MNB*) The meteorite has a totally recrystallized texture, with abundant  $120^\circ$  triple junctions of olivine ( $\text{Fa}_{17.7}$ ), low-Ca pyroxene ( $\text{Fs}_{15.6}\text{Wo}_{3.6}$ ), and feldspar ( $\text{An}_{12.3-18.9}$ ). No chondrules were observed.

**Geochemistry:** Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = +2.905$ ,  $\delta^{18}\text{O} = +4.4$ ,  $\Delta^{17}\text{O} = +0.617$  (all ‰).

**Classification:** Ordinary chondrite (H7); S2, W1/2.

**Specimens:** A 23 g type specimen and one polished thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

**Northwest Africa 2899**

Northwest Africa

Find: 2004

Ordinary chondrite (H-related impact-melt rock)

**History:** A single stone of 11 g was found 2004 by an anonymous finder in the North African Saharan desert.

**Petrography:** (A. Greshake, *MNB*) The stone is composed of small, mostly euhedral olivines ( $\text{Fa}_{6.1-23.9}$ ) embedded into a glassy groundmass. Most olivines show a pronounced compositional zoning with Mg-rich cores and Fe-rich rims or vice versa. Small spherical Fe,Ni sulfides are often found in the groundmass and Fe,Ni metal is present. Chondrules were not observed.

**Geochemistry:** Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = +3.088$ ,  $\delta^{18}\text{O} = +4.898$ ,  $\Delta^{17}\text{O} = +0.541$  (all ‰).

**Classification:** Ordinary chondrite (H-related impact-melt rock) with shock-related crystallized impact melt, W0/1.

**Specimens:** A 2.5 g type specimen and one polished thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

**Northwest Africa 2900**

Northwest Africa

Find: 2004

Carbonaceous chondrite (CV3)

**History:** An anonymous finder recovered a single stone of 1375 g within the North African Sahara in 2004.

**Petrography:** (A. Greshake, *MNB*) The stone has a grayish appearance and contains up to centimeter-size dark- and light-colored inclusions and is composed of large chondrules, CAIs, and mineral fragments—all set into a fine-grained, grayish matrix.

**Geochemistry:** Olivine ( $\text{Fa}_{14.2}$ , range  $\text{Fa}_{0.2-31.1}$ ), low-Ca

pyroxene ( $\text{Fs}_{0.8-22.9}$ ). Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = -1.361$ ,  $\delta^{18}\text{O} = +3.027$ ,  $\Delta^{17}\text{O} = -2.935$  (all ‰).

**Classification:** Carbonaceous chondrite (CV3); low shock, moderate weathering.

**Specimens:** A 20.5 g type specimen and one thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

#### Northwest Africa 2901

Northwest Africa

Find: 2004

Carbonaceous chondrite (CV3)

**History:** An anonymous finder recovered a single stone of 308 g within the North African Sahara in 2004.

**Petrography:** (A. Greshake, *MNB*) The sample contains large chondrules, CAIs, and mineral fragments set into a fine-grained, dark-appearing matrix.

**Geochemistry:** Olivine ( $\text{Fa}_{25.9}$ , range  $\text{Fa}_{0.4-32.1}$ ), low-Ca pyroxene ( $\text{Fs}_{20.8-22.7}$ ). Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = -1.226$ ,  $\delta^{18}\text{O} = +3.587$ ,  $\Delta^{17}\text{O} = -3.091$  (all ‰).

**Classification:** Carbonaceous chondrite (CV3); low shock, extensive weathering.

**Specimens:** A 21.1 g type specimen and one thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

#### Northwest Africa 2902

Northwest Africa

Find: 2003

Ordinary chondrite (L-related, impact-melt rock)

**History:** An anonymous finder recovered several stones weighing 1000 g in total within the North African Sahara in 2003.

**Petrography:** (A. Greshake, *MNB*) Shows both Fe,Ni metal-poor and Fe,Ni metal-rich lithologies that both consist of small olivines, pyroxenes, and opaque phases embedded into a glassy Si,Al-rich groundmass. Ca-rich pyroxenes are rare. Olivine ( $\text{Fa}_{12.9-23.9}$ ) and pyroxene ( $\text{Fs}_{11.2-19.7}\text{Wo}_{1.1-2.7}$ ) are often compositionally zoned and sometimes pyroxene is overgrown by olivine. No chondrules are observed.

**Geochemistry:** Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*) Metal-poor lithology:  $\delta^{17}\text{O} = +3.24$ ,  $\delta^{18}\text{O} = +4.095$ ,  $\Delta^{17}\text{O} = +1.106$  (all ‰). Metal-rich lithology:  $\delta^{17}\text{O} = +3.452$ ,  $\delta^{18}\text{O} = +4.505$ ,  $\Delta^{17}\text{O} = +1.109$  (all ‰).

**Classification:** Ordinary chondrite (L-related, impact-melt rock); shock-related crystallized impact melt, W1/2.

**Specimens:** A 22.7 g type specimen and one polished thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

#### Northwest Africa 2904

Northwest Africa

Find: Spring 2003

Achondrite (eucrite, polymict)

**History:** Four stones totaling 29.04 g were recovered in the spring of 2003 from the western part of the Sahara in Morocco.

**Classification and Mineralogy:** (A. Greshake and M. Kurz, *MNB*) A polymict breccia consisting of lithic and mineral clasts set into a more fine-grained, brecciated groundmass of dominantly plagioclase and often exsolved Ca pyroxene; plagioclase frequently contains pyroxene and ilmenite inclusions. Plagioclase,  $\text{An}_{84.6-90.7}$ ; low-Ca pyroxene,  $\text{Fs}_{44.5-55.8}\text{Wo}_{1.8-2.1}$ ; Ca pyroxene,  $\text{Fs}_{22.3-43.4}\text{Wo}_{15.6-42.6}$ ; minor phases include Ti-rich chromite, silica, and ilmenite. Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = +2.16$ ,  $\delta^{18}\text{O} = +4.58$ ,  $\Delta^{17}\text{O} = -0.222$  (all ‰) and (R. Clayton and T. Mayeda, *UChi*):  $\delta^{17}\text{O} = +2.13$ ,  $\delta^{18}\text{O} = +4.74$ ,  $\Delta^{17}\text{O} = -0.33$  (all ‰).

**Classification:** Achondrite (eucrite, polymict); moderate to high shock, extensive weathering.

**Specimens:** A 9.95 g type specimen and one polished thin section are on deposit at *MNB*. Main mass with an anonymous finder.

#### Northwest Africa 2913

Morocco

Find: 2005

Achondrite (eucrite, monomict melt breccia)

**History:** A 100.2 g complete stone was purchased in Erfoud, Morocco, in May 2005.

**Petrography:** (J. Wittke and T. Bunch, *NAU*) Specimen contains medium-grained (<2 mm), centimeter-size, cumulate basalt clasts set in a microbreccia melt matrix. Clasts are moderately shocked with partial maskelynitization of plagioclase and mechanical twinning, optical mosaicism and reduced birefringence in pyroxenes. The dark matrix is a flow mixture of FeO-rich glasses, very small cumulate basalt fragments, and patches of quench-textured microbasalt.

**Geochemistry:** Monomict cumulate basalt: host Ca-poor pyroxene ( $\text{Fs}_{55.4-57.8}\text{Wo}_{5.1-2.9}$ ),  $\text{FeO/MnO} = 28$ ; exsolved augite ( $\text{Fs}_{24.2}\text{Wo}_{44.7}$ ),  $\text{FeO/MnO} = 36$ ; plagioclase/maskelynite ( $\text{An}_{90.8}$ ); chromite  $\text{Cr}/(\text{Cr}+\text{Al}) = 0.86$ .

**Classification:** Achondrite (eucrite, monomict melt breccia); moderate shock, minimal weathering.

**Specimens:** A 21 g type specimen is deposited at *NAU*. Birdsell holds the main mass.

#### Northwest Africa 2914

Morocco

Find: April 2005

Achondrite (eucrite, monomict breccia)

**History:** Three paired partial stones weighing a total of 122 g were purchased in Erfoud, Morocco, in April 2005.

**Petrography:** (T. Bunch and J. Wittke, *NAU*) Cataclastic cumulate basalt that shows a remarkable range of shock features that include alternating bands of cataclastic and melt flows, mylonite-like swirls and lenticular masses, and

rounded, finely crushed breccia clasts that are separated by dark, glassy to cryptocrystalline bands.

**Geochemistry:** Host: orthopyroxene ( $\text{Fs}_{59.3-62.7}\text{Wo}_{3.1-1.9}$ ;  $\text{FeO/MnO} = 31$ ), exsolved augite ( $\text{Fs}_{26.8-27.8}\text{Wo}_{43-42.1}$ ;  $\text{FeO/MnO} = 28$ ), plagioclase ( $\text{An}_{92}$ ); chromite ( $\text{Cr}/(\text{Cr}+\text{Al}) = 0.88$ ).

**Classification:** Achondrite (eucrite monomict breccia) with moderate to high degree of shock and low degree of weathering.

**Specimens:** A 21.1 g type specimen is on deposit at *NAU*. Birdsell holds the main mass.

#### Northwest Africa 2923

Morocco

Find: July 2005

Achondrite (diogenite polymict breccia)

**History:** A 606 g complete stone fully crusted with moderately fresh fusion crust was purchased in Erfoud, Morocco, in July 2005.

**Physical Characteristics:** (T. Bunch and J. Wittke, *NAU*) A light buff-color interior matrix with clear, yellow, green, dark diogenite clasts.

**Petrography:** Modal analyses of a 27 cm<sup>2</sup> surface area: diogenites (grain size <2.7 cm) = 83, olivine microbasalts = 6, shock melt clasts = 5, recrystallized cumulate basalts = 4, chromite, sulfide, ilmenite, metal = 2 (all vol%).

**Geochemistry:** Diogenite: Orthopyroxenes ( $\text{Fs}_{22-27.6}\text{Wo}_{2.9}$ ;  $\text{FeO/MnO} = 28-31$ ). Unique olivine microbasalt clasts: olivine ( $\text{Fa}_{33.6}$ ;  $\text{FeO/MnO} = 54$ ; these are microphenocrysts <0.48 mm in diameter), orthopyroxene (<0.2 mm) ( $\text{Fs}_{27.7}\text{Wo}_{2.8}$ ;  $\text{FeO/MnO} = 27$ ) with <0.05 mm diameter grains of vermicular metal ( $\text{Ni} = 6.2$  wt%), and FeS ( $\text{Cr} = 6.2$  wt%); minor diopside ( $\text{Fs}_{10.0}\text{Wo}_{45.7}$ ).

**Classification:** Achondrite (diogenite, polymict breccia); low to high (recrystallized melts) shock, minimal weathering.

**Specimens:** A 50 g type specimen is on deposit at *NAU*. Farmer holds the main mass.

#### Northwest Africa 2924

Algeria

Find: 2005

Mesosiderite

**History:** Over 180 kg (several thousand pieces) have been recovered from a small strewn field in Algeria. The largest known piece weighs about 5000 g; most are small pieces that weigh less than 200 g.

**Petrography and Geochemistry:** (T. Bunch and J. Wittke, *NAU*) Fine-grained (<0.5 mm) matrix consists of brecciated and partially recrystallized subophitic to granular basalts that enclose larger (up to 2.5 mm) subrounded to subangular olivine ( $\text{Fa}_{28}$ ), pigeonite ( $\text{Fs}_{31}\text{Wo}_{5.4}$ ;  $\text{FeO/MnO} = 25.7$ ;  $\text{Cr}_2\text{O}_3 = 0.87$ ,  $\text{Al}_2\text{O}_3 = 1.28$  [both wt%]) that rarely shows twinning. These cataclastic basalts contain unusually calcic plagioclase ( $\text{An}_{99.3}$ ) with little FeO (<0.04 wt%); highly twinned pigeonite is compositionally similar to the much larger clast pigeonite ( $\text{Fs}_{32}\text{Wo}_{5.7}$ ), but with very little  $\text{Cr}_2\text{O}_3$  and  $\text{Al}_2\text{O}_3$ .

Merrillite is also present. Metal blebs ( $\text{Ni} = 6.6$  wt%) are irregular in shape and typically <2 mm in size with the exception of metal aggregates and rare, centimeter-size metal nuggets with included pigeonite.

**Classification:** Mesosiderite; low shock, variable degree of weathering dependent on the specimen size.

**Specimens:** A 85 g type specimen is on deposit at *NAU*. Farmer holds a 16 kg of sample.

#### Northwest Africa 2975

Algeria

Find: 2005

Achondrite (Martian, basaltic shergottite)

**History:** A minimally weathered fully encrusted whole stone of 70.1 g was purchased in Erfoud, Morocco, by M. Farmer in November 2005.

**Petrography:** (T. Bunch and J. Wittke, *NAU*; A. Irving, *UWS*) A medium-grained (<3.1 mm greatest dimension) basaltic shergottite that consists of ~57.3 vol% augite and pigeonite pyroxenes, and 38.3 vol% plagioclase (present as shock-formed maskelynite and glasses) with minor opaques (2.7 vol%) and phosphates (1.7 vol%) arranged in a weakly foliated subophitic to granular texture. Accessory phases include ulvöspinel, ilmenite, chlorapatite, merrillite, pyrrhotite, Si-Al-Na-K-rich glasses, and baddeleyite. Vesicular black glass veins (<3 mm in width) and pockets (up to 6 mm) are prominent.

**Geochemistry:** Pigeonite ( $\text{Fs}_{35.2-57.6}\text{Wo}_{12.6-16.5}$ ;  $\text{FeO/MnO} = 28-38$ ) and augite ( $\text{Fs}_{27.2-41.5}\text{Wo}_{30.8-35.2}$ ) show mottled compositional zoning. Fe-rich margins of pigeonite contain very thin (0.2–0.5  $\mu\text{m}$ ) orthopyroxene exsolution lamellae. Maskelynite is compositionally homogeneous  $\text{An}_{55}\text{Or}_{1.8}$  in contrast to mesostasis maskelynite ( $\text{An}_{48-60}\text{Or}_{9.2}$ ) and glasses. Melt inclusions (15–60  $\mu\text{m}$ , longest dimension) in ulvöspinel have rims of Fe-rich pigeonite ( $\text{Fs}_{73.5}\text{Wo}_{5.8}$ ), merrillite, and pyrrhotite; cores are Si-Al-K-Na-rich glass. Bulk composition: Calculated average bulk composition based on analyzed phases and their modes of three melt inclusions is:  $\text{SiO}_2 = 70.2$ ,  $\text{Al}_2\text{O}_3 = 8.5$ ,  $\text{TiO}_2 = 0.8$ ,  $\text{FeO} = 2.1$ ,  $\text{MnO} = 0.2$ ,  $\text{MgO} = 2.35$ ,  $\text{CaO} = 3.5$ ,  $\text{Na}_2\text{O} = 0.95$ ,  $\text{K}_2\text{O} = 2.1$ , and  $\text{P}_2\text{O}_5 = 2.8$  (all wt%) with trace amounts of NiO, CoO, and S. One melt inclusion contains only fayalite ( $\text{Fa}_{84}$ ) mantled around a core of glass ( $\text{K}_2\text{O} = 6.4$ ,  $\text{SiO}_2 = 78$ ; both wt%).

**Classification:** Achondrite (Martian basaltic shergottite); moderate to high shock, minimal weathering.

**Specimens:** A 20.2 g type specimen and one thin section are on deposit at *NAU*. An anonymous owner holds the main mass.

#### Northwest Africa 2977

Morocco or Algeria

Find: 2005 November

Achondrite (lunar, gabbro)

**History:** A single minimally weathered fusion-encrusted stone of 233 g was purchased from a Moroccan dealer in Tagounite, Morocco, by M. Farmer in November 2005.

**Petrography and Geochemistry:** (J. Wittke and T. Bunch, *NAU*; A. Irving, *UWS*) The specimen consists of a single yellow-green, relatively coarse-grained rock traversed by thin, black glass-rich veins. It is an olivine-rich, two-pyroxene cumulate gabbro composed of olivine ( $\text{Fa}_{31.7}$ ;  $\text{FeO}/\text{MnO} = 96$ ; 52 vol%), ( $\text{Fs}_{26.6}\text{Wo}_{6.7}$ ; 23 vol%), augite ( $\text{Fs}_{16.2}\text{Wo}_{29}$ ; 9 vol%), and plagioclase ( $\text{An}_{56}$ ; 14 vol%) with minor amounts of Ba-K feldspar, chromite, ilmenite, and merrillite. Larger pigeonite grains commonly enclose equant olivine grains, which contain abundant melt inclusions (0.025–0.125 mm). Plagioclase is partially converted to maskelynite, and pyroxenes and olivine exhibit shock lamellae and undulatory extinction. Note: This specimen is identical in texture and mineral composition to the gabbro clasts in NWA 773 and NWA 2700 and thus appears to be paired with those breccia specimens.

**Classification:** Achondrite (lunar, gabbro); minimal weathering.

**Specimens:** A 20.1 g type specimen and one polished thin section are on deposit at *NAU*. A 0.5 g specimen is on deposit at *WUSL*. An anonymous owner holds the main mass.

#### Northwest Africa 2995

Algeria

Find: 2005

Achondrite (lunar feldspathic breccia)

**History:** A 538 g fully crusted and minimally weathered stone was purchased in Morocco by A. Aaronson in November 2005.

**Petrography and Geochemistry:** (T. Bunch and J. Wittke, *NAU*) The feldspathic fragmental breccia contains many highlands fine-grained lithologies. Norite: Orthopyroxene ( $\text{Fs}_{26.4}\text{Wo}_4$ ;  $\text{FeO}/\text{MnO} = 66$ ). Olivine basalt: Olivine ( $\text{Fa}_{87.2}$ ;  $\text{FeO}/\text{MnO} = 95$ ), plagioclase ( $\text{An}_{84.7}$ ). Subophitic basalt: Ca pyroxene ( $\text{Fs}_{25-48}\text{Wo}_{37.1-25.9}$ ), pigeonite ( $\text{Fs}_{27.8-31.7}\text{Wo}_{15.4-9.3}$ ;  $\text{FeO}/\text{MnO} = 53$ ), olivine ( $\text{Fa}_{36.3}$ ;  $\text{FeO}/\text{MnO} = 90$ ), plagioclase ( $\text{An}_{97}$ ). Gabbro: Olivine ( $\text{Fa}_{34.7}$ ;  $\text{FeO}/\text{MnO} = 95$ ), pigeonite ( $\text{Fs}_{28.2}\text{Wo}_{8.9}$ ;  $\text{FeO}/\text{MnO} = 67$ ), plagioclase ( $\text{An}_{94}$ ). KREEPy-like basalt: Plagioclase ( $\text{Ab}_{50}\text{Or}_{17.4}$ ), K feldspar ( $\text{Ab}_{14.3}\text{Or}_{83}$ ) in addition to silica, phosphate, and Fe-rich pyroxenes. Troctolite: olivine ( $\text{Fa}_{30.8}$ ;  $\text{FeO}/\text{MnO} = 94$ ), plagioclase ( $\text{An}_{94.7}$ ). Granulitic impact melts: Olivine ( $\text{Fa}_{31}$ ), orthopyroxene ( $\text{Fs}_{25.2}\text{Wo}_{3.4}$ ), plagioclase ( $\text{An}_{95}$ ). Anorthosite: ( $\text{An}_{92.7-96.8}$ ), glassy impact melts, coarse-grained mineral fragments, and a 0.350 mm-size grain of meteoritic Ni, Fe-metal ( $\text{Ni} = 6.3$ ,  $\text{Co} = 1.0$  [both wt%]). In addition, the assemblage appears to be characterized by large amounts of breccias within breccias with at least four generations of brecciation observed in one centimeter-size breccia clast. Numerous shock-induced melt veins are present along large breccia clast margins as well as isolated melt pockets within clasts. Interior weathering grade is very low, all glasses are fresh, and no apparent terrestrial alteration veins were noted.

**Classification:** Achondrite (lunar, feldspathic breccia).

**Specimens:** A 21.2 g type specimen is on deposit at *NAU*. Aaronson holds the main mass.

#### Northwest Africa 2999

Morocco or Algeria

Find: 2004

Achondrite (angrite)

**History:** Twelve individual dark brown stones totaling 392 g, each with a thin fusion crust, were purchased from a Moroccan dealer in Tagounite by G. Hupé in August 2004.

**Physical Characteristics:** Grain size is predominantly from 0.1 to 0.5 mm, but all stones have irregularly distributed, larger yellowish plagioclase grains (up to 6 mm across) exhibiting an iridescent luster.

**Petrography:** (A. Irving and S. Kuehner, *UWS*; T. Bunch and J. Wittke, *NAU*) Based upon examination of thin sections of all separate stones, this meteorite is texturally heterogeneous. Terrestrial weathering has resulted in partial replacement of metal and minor grain boundary staining by iron hydroxides. The overall texture is protogranular, but there are large porphyroclasts of anorthite, spinel, and polygranular olivine. Anorthite also occurs as narrow (10–20  $\mu\text{m}$  wide) coronas around spinel grains adjacent to clinopyroxene and both spinel and diopside are compositionally zoned away from the coronas. Texturally, this meteorite is very different from most angrites.

**Geochemistry:** The major minerals are Ca-rich olivine ( $\text{Fa}_{39.8-41.0}$ ;  $\text{FeO}/\text{MnO} = 77-97$ ;  $\text{CaO} = 0.6-1.3$  wt%), Al,Ti-bearing diopside ( $\text{Fs}_{9.6-11.3}\text{Wo}_{53-54}$ ;  $\text{FeO}/\text{MnO} = 55-130$ ;  $\text{Al}_2\text{O}_3 = 5-9$ ,  $\text{TiO}_2 = 0.5-2.4$  [both wt%]), minor Cr-pleonaste spinel ( $\text{Mg}/(\text{Mg}+\text{Fe}) = 0.44-0.47$ ,  $\text{Al}_2\text{O}_3 = 55-60$ ,  $\text{Cr}_2\text{O}_3 = 4.7-8.7$  [both wt%]), pure anorthite (containing  $\text{Na}_2\text{O} < 0.02$  wt%), and kamacite, troilite, and S-bearing calcium silicophosphate. Oxygen isotopes: (D. Rumble, *CIW*) Triplicate analyses of acid-washed whole rock samples by laser fluorination gave, respectively,  $\delta^{18}\text{O} = 3.839$ , 4.093, 4.154;  $\delta^{17}\text{O} = 1.974$ , 2.054, 2.095;  $\Delta^{17}\text{O} = -0.041$ ,  $-0.095$ ,  $-0.086$  (all ‰).

**Classification:** Achondrite (angrite).

**Specimens:** A 22 g type specimen and one polished thin section are on deposit at *NAU*. Three polished thin sections are on deposit at *UWS*. G. Hupé holds the main mass.

#### Northwest Africa 3151

Morocco or Algeria

Find: April 2005

Achondrite (brachinite)

**History:** A 1500 g complete stone with a thin, translucent crust was purchased by G. Hupé in Tagounite, Morocco, in April 2005.

**Petrography and Geochemistry:** (A. Irving and S. Kuehner, *UWS*) Coarse-grained dunitic rock (grain size 0.7–1.6 mm) with protogranular texture, composed predominantly of olivine ( $\text{Fa}_{35.7}$ ;  $\text{FeO}/\text{MnO} = 81$ ; 95 vol%) with minor clinopyroxene ( $\text{Fs}_{10.5-10.6}\text{Wo}_{44.7-45.2}$ ;  $\text{FeO}/\text{MnO} = 44$ ;  $\text{Cr}_2\text{O}_3 = 0.65$ ,  $\text{Al}_2\text{O}_3 = 0.95$  [both wt%]), altered metal (with some relict taenite), troilite, chromite ( $\text{Cr}/(\text{Cr}+\text{Al}) = 0.727-0.731$ ) with very rare K-poor, sodic plagioclase ( $\text{An}_{36.2-39.9}\text{Or}_{0.2}$ ),

and orthopyroxene. Although the silicate minerals are fresh, terrestrial weathering has altered some of the primary metal to hydroxides, which also form thin coatings along grain boundaries. Oxygen isotopes: (D. Rumble, *CIW*) Replicate analyses of acid-washed whole rock samples by laser fluorination gave  $\delta^{18}\text{O} = +4.86 \pm 0.03$ ,  $\delta^{17}\text{O} = +2.42 \pm 0.02$ ,  $\Delta^{17}\text{O} = -0.15 \pm 0.02$  (all ‰).

**Classification:** Achondrite (brachinite).

**Specimens:** A 20 g type specimen and one polished thin section are on deposit at *UWS*. G. Hupé holds the main mass.

#### Northwest Africa 3160

Morocco

Find: July 2005

Achondrite (lunar, mare basalt breccia)

**History:** In July 2005, A. and G. Hupé purchased three broken stones with a total weight of 34 g from a Moroccan dealer in Erfoud, Morocco.

**Physical Characteristics:** The largest stone (28 g) has a partial thin weathered fusion crust.

**Petrography:** (R. Zeigler and R. Korotev, *WUSL*; A. Irving and S. Kuehner, *UWS*) The large specimen consists almost entirely of a fine-grained, olivine-phyric basalt clast with minor attached breccia matrix and appears to be part of a larger, coarse-grained, polygenic breccia. The two small stones are pieces of the breccia. The basalt contains phenocrysts of euhedral to subhedral olivine (~0.1–0.9 mm) and minor chromite (<0.1 mm).

**Geochemistry:** Olivine phenocrysts are zoned, with cores typically  $\text{Fo}_{55-70}$  and rims extending to  $\sim\text{Fo}_{40}$  with  $\text{FeO}/\text{MnO}$  ratios of 91–105. The groundmass has spinifex olivine ( $\text{Fo}_{29}$ ) and skeletal pyroxene ( $\text{En}_{37-39}\text{Wo}_{11-13}$ ;  $\text{FeO}/\text{MnO} = 71-75$ ) set in a fine-grained matrix of pyroxene ( $\text{En}_{35-39}\text{Wo}_{20-23}$ ), olivine ( $\sim\text{Fo}_{22}$ ), and glass. The breccia lithology is a fragmental breccia consisting primarily of olivine ( $\text{Fo}_{6-82}$ ) and pyroxene ( $\text{En}_{1-68}\text{Wo}_{9-39}\text{Fs}_{16-83}$ ), with minor amounts of plagioclase ( $\text{An}_{82-97}$ ) and trace silica; hedenbergite-fayalite-silica symplectite (after former pyroxferroite), and Fe-Ti-Cr oxides.

**Classification:** Achondrite (lunar, mare basalt breccia). Note: These samples may be paired with NWA 2727.

**Specimens:** A 4.8 g type specimen and one polished thin section are on deposit at *UWS*. A 2.1 g specimen is on deposit at *WUSL*. A. Hupé holds the main mass.

#### Northwest Africa 3163

Mauritania or Algeria

Find: August 2005

Achondrite (lunar, feldspathic granulitic impactite)

**History:** In August 2005, G. Hupé purchased a 1634 g stone from a Moroccan dealer in Ouarzazate.

**Petrography and Geochemistry:** (A. Irving and S. Kuehner, *UWS*) The exterior is almost completely coated by a thin, transparent, greenish fusion crust. The pale gray interior has multiple shock fractures (with very minor calcite coatings) and some thin glass veins. Poikiloblastic recrystallized

breccia, with larger grains of plagioclase (~70 vol%) enclosing much smaller grains (less than 100  $\mu\text{m}$  across) of pyroxenes (~20 vol%), olivine (~10 vol%), and accessory Ti-chromite ( $\text{Cr}/(\text{Cr}+\text{Al}) = 0.714-0.736$ ;  $\text{Mg}/(\text{Mg}+\text{Fe}) = 0.121-0.143$ ;  $\text{TiO}_2 = 9.1-18.4$  wt%), ilmenite, troilite, and metal ( $\text{Ni} = \sim 15$  wt%). Anorthitic plagioclase ( $\text{An}_{97.4-98.2}$ ) has been converted by shock almost entirely to maskelynite (although domains of birefringent, less-shocked feldspar remain). Pigeonitic pyroxene grains have very fine-scale exsolution lamellae of augite ( $\text{Fs}_{14.5-16.1}\text{Wo}_{40.2-40.5}$ ;  $\text{FeO}/\text{MnO} = 41.7-43.8$ ) within orthopyroxene ( $\text{Fs}_{32.0-33.9}\text{Wo}_{4.4-5.8}$ ;  $\text{FeO}/\text{MnO} = 55.5-61.2$ ). Olivine ( $\text{Fa}_{38.0-40.9}$ ;  $\text{FeO}/\text{MnO} = 91.7-110$ ).

**Classification:** Achondrite (lunar, feldspathic granulitic impactite).

**Specimens:** A 20.1 g type specimen and one polished thin section are on deposit at *UWS*. G. Hupé holds the main mass.

#### Northwest Africa 3164

Morocco or Algeria

Find: August 2004

Achondrite (angrite)

**History:** In August 2004, A. Aaronson purchased many individual dark brown stones totaling 928 g from nomads in Rabat.

**Petrography:** (T. Bunch and J. Wittke, *NAU*; A. Irving and S. Kuehner, *UWS*) All stones have irregularly distributed, larger yellowish plagioclase grains exhibiting a luster. The overall texture is granular, but there are large porphyroclasts of anorthite, spinel, and polygranular olivine. There are clinopyroxene-spinel symplectites around anorthite porphyroclasts in contact with olivine, and anorthite also occurs as narrow (10–20  $\mu\text{m}$  wide) coronas around spinel grains adjacent to clinopyroxene.

**Geochemistry:** The major minerals are Ca-rich olivine ( $\text{Fa}_{39.1-41.2}$ ;  $\text{FeO}/\text{MnO} = 62-84$ ;  $\text{CaO} = 1.2-1.8$  wt%), Al,Ti-bearing diopside ( $\text{Fs}_{10.3}\text{Wo}_{52}$ ;  $\text{FeO}/\text{MnO} = 130-142$ ;  $\text{Al}_2\text{O}_3 = 6-7$ ,  $\text{TiO}_2 = 1-1.6$  [both wt%]), minor Cr-bearing pleonaste spinel ( $\text{Mg}/(\text{Mg}+\text{Fe}) = 45.7$ ,  $\text{Al}_2\text{O}_3 = 59.7$ ,  $\text{Cr}_2\text{O}_3 = 4.7$  [both wt%]), almost pure anorthite (<0.02 wt%  $\text{Na}_2\text{O}$ ), and accessory kamacite and troilite. No kirschsteinite or orthopyroxene was found. Primary metal is partly replaced by limonite, which also occurs along grain boundaries.

**Classification:** Achondrite (angrite). Note: The sample may be paired with NWA 2999.

**Specimens:** A 21 g type specimen and one polished thin section are on deposit at *NAU*. Boswell holds the main mass.

#### Northwest Africa 3368

Northwest Africa

Find: 2005

Achondrite (eucrite)

**History:** John Birdsell and Marvin Killgore purchased a 1600 g stone.

**Petrography:** (D. Hill and K. Gardner, *UAz*) The stone was 100% covered with black fusion crust. There are a variety of

light and dark angular clasts ranging in size from several mm to ~2 cm and rounded grains several millimeters in diameter. They are set in a light matrix typical of eucrites except for its distinct pink hue. Some systematic variations in the color exist. Approximately half of the slice is slightly darker pink than the other with an indistinct, irregular boundary.

**Geochemistry:** Pyroxene ( $\text{Wo}_6\text{En}_{36}\text{Fs}_{59}$  and  $\text{Wo}_{40}\text{En}_{30}\text{Fs}_{30}$ ), plagioclase ( $\text{An}_{90}\text{Ab}_{10}\text{Or}_{0.4}$ ), chromite ( $\text{TiO}_2 = 5\text{--}27$ ,  $\text{Al}_2\text{O}_3 = 5$  [both wt%]), with minor ilmenite present. Bulk INAA (*UAz*) indicate  $\sim 10\times$  CI REE abundances.

**Classification:** Achondrite (eucrite); minimal weathering.

**Specimens:** One thick section type specimen of 20.5 g and three small chips totaling 2.5 g are on deposit at *UAz*. M. Killgore holds the main mass.

#### Northwest Africa 4017

Northwest Africa

Find: 2005

Mesosiderite

**History:** An anonymous finder recovered a single stone of 216.9 g in 2005 from the western part of the Sahara in Morocco.

**Petrography:** (A. Greshake, *MNB*) A stony portion consists of dominant low-Ca pyroxene, plagioclase, and phosphates. Minor phases include silica, troilite, and small Fe,Ni metal grains. Pyroxene grains often contain numerous small Fe,Ni metal, troilite, and silica inclusions.

**Geochemistry:** Pyroxene,  $\text{Fs}_{25.5\text{--}40.8}$ ; plagioclase,  $\text{An}_{92.4}$  (total range  $\text{An}_{90.5\text{--}93.7}$ ). The metal is mostly kamacite.

**Classification:** Achondrite (mesosiderite) with low degree of shock and low to moderate degree of weathering.

**Specimens:** A 23.4 g type specimen and one polished thin section are on deposit at *MNB*. HSSH-Pirmasens of Germany holds the main mass.

#### Northwest Africa 4018

Northwest Africa

Find: 2005

Achondrite (eucrite, polymict)

**History:** A single stone of 158.6 g was found in 2005 in the western part of the Sahara in Morocco.

**Petrography:** (A. Greshake, *MNB*) A polymict breccia composed of lithic and mineral clasts set into a fine-grained brecciated groundmass. The lithic clasts are mostly fine- or coarse-grained basaltic fragments with rare impact melt clasts. Mineral fragments are dominantly plagioclase and exsolved Ca pyroxene. Minor phases include silica, Al-Ti-chromite, and ilmenite.

**Geochemistry:** Plagioclase ( $\text{An}_{87\text{--}92.2}$ ), pyroxene ( $\text{Fs}_{42.9\text{--}54.0}\text{Wo}_{2.7\text{--}19.0}$ ).

**Classification:** Achondrite (eucrite, polymict); moderate shock, moderate weathering.

**Specimens:** A 23.1 g type specimen and one polished thin section are on deposit at *MNB*. HSSH-Pirmasens of Germany holds the main mass.

#### Northwest Africa 4019

Northwest Africa

Find: 2005

Achondrite (eucrite, polymict)

**History:** A single stone of 504.7 g partly covered with fusion crust was found in 2005 in the western part of the Sahara in Morocco.

**Petrography:** (A. Greshake, *MNB*) A polymict breccia dominated by large basaltic clasts set into a fine-grained groundmass with rare melt clasts and large mineral fragments, the latter being mostly plagioclase and exsolved Ca pyroxene. Minor phases include silica and ilmenite.

**Geochemistry:** Plagioclase ( $\text{An}_{74.1\text{--}90.7}$ ), pyroxene ( $\text{Fs}_{41.6\text{--}65.4}\text{Wo}_{2.9\text{--}29.3}$ ).

**Classification:** Achondrite (eucrite, polymict); low shock, minimal weathering.

**Specimens:** A 22 g type specimen and one polished thin section are on deposit at *MNB*. HSSH-Pirmasens of Germany holds the main mass.

#### Northwest Africa 4023

Northwest Africa

Find: 2005

Achondrite (eucrite, polymict)

**History:** A single stone of 17.2 g partly covered with fusion crust was found in the western part of the Sahara in Morocco in 2005.

**Petrography:** (A. Greshake, *MNB*) A polymict breccia consisting of basaltic, melt clasts, and mineral clasts set into a fine-grained brecciated groundmass. Mineral fragments are dominantly plagioclase and exsolved Ca pyroxene. Minor phases include silica, Al-Ti-chromite, and ilmenite.

**Geochemistry:** Plagioclase ( $\text{An}_{78.6\text{--}94.2}$ ), pyroxene ( $\text{Fs}_{27.7\text{--}56.5}\text{Wo}_{2.7\text{--}40.2}$ ).

**Classification:** Achondrite (eucrite, polymict) with a high degree of shock and moderate degree of weathering.

**Specimens:** A 3.45 g type specimen and one polished thin section are on deposit at *MNB*. HSSH-Pirmasens of Germany holds the main mass.

#### Northwest Africa 4032

Northwest Africa

Find: 2004

Achondrite (eucrite, polymict)

**History:** A single stone of 10.5 g partly covered with fusion crust was found in the western Sahara in Morocco in 2004.

**Petrography:** (A. Greshake, *MNB*) A polymict breccia consisting of basaltic and mineral clasts set into a fine-grained groundmass with mineral fragments that are dominantly large plagioclase and exsolved Ca pyroxene. Minor phases include silica, troilite, and ilmenite.

**Geochemistry:** Plagioclase ( $\text{An}_{79\text{--}91.4}$ ), pyroxene ( $\text{Fs}_{26.6\text{--}61.2}\text{Wo}_{2.6\text{--}42.5}$ ).

**Classification:** Achondrite (eucrite, polymict); high shock, minimal weathering.



**Specimens:** A 2.2 g type specimen and one polished thin section are on deposit at *MNB*. JNMC-Zürich, Switzerland, holds the main mass.

#### Northwest Africa 4034

Northwest Africa

Find: 2005

Achondrite (diogenite)

**History:** A single stone of 1513 g partly covered with fusion crust was found in 2005 by an anonymous finder in the western part of the Sahara in Morocco.

**Petrography:** (A. Greshake, *MNB*) The specimen is dominated by large blocky orthopyroxenes; locally, these crystals are brecciated into smaller grains. Minor phases include Mg-Al-Cr-spinel and plagioclase.

**Geochemistry:** Orthopyroxene ( $\text{Fs}_{25.2}\text{Wo}_{3.3}$ ), plagioclase ( $\text{An}_{81.3-83}$ ).

**Classification:** Achondrite (diogenite); moderate shock, moderate to extensive weathering.

**Specimens:** A 24.5 g type specimen and one polished thin section are on deposit at *MNB*. JNMC-Zurich of Zurich, Switzerland, holds the main mass.

#### Northwest Africa 4039

Northwest Africa

Find: 2005

Achondrite (eucrite, monomict breccia)

**History:** Several stones totaling 950 g were found in 2005 by an anonymous finder in 2004 in the western part of the Sahara in Morocco.

**Petrography:** (A. Greshake, *MNB*) An unusual eucrite with dominantly coarse-grained basaltic texture of exsolved Ca pyroxene and plagioclase cross-cut by centimeter-size granular bands or crushed zones. Locally these zones display an equigranular texture with  $120^\circ$  grain boundaries; minor phases include orthopyroxene, silica, ilmenite, troilite, and Al-Ti chromite.

**Geochemistry:** Plagioclase ( $\text{An}_{90.1}$ , range  $\text{An}_{88.3-92.6}$ ), orthopyroxene ( $\text{Fs}_{58.1-62.2}\text{Wo}_{2.7-4.6}$ ), pigeonite ( $\text{Fs}_{50.7-58.6}\text{Wo}_{5.2-15.7}$ ), augite ( $\text{Fs}_{26.3-34.9}\text{Wo}_{32.1-44.1}$ ).

**Classification:** Achondrite (eucrite, monomict breccia); high shock, moderate weathering.

**Specimens:** A 20.1 g type specimen and one polished thin section are on deposit at *MNB*. An anonymous finder holds the main mass.

#### Northwest Africa 4040

Find: 2004

Ordinary chondrite (L3)

**History:** A complete stone weighing 226.3 g was found by an anonymous finder in the western Sahara in Morocco.

**Petrography:** (A. Greshake, *MNB*) The sample is a brecciated ordinary chondrite with centimeter-size dark and light inclusions. The dark inclusion is composed of small anhedral, mostly skeletal olivine displaying compositional

zoning from an Mg-rich core to a more Fe-rich rim embedded into a mostly glassy groundmass containing small dendrites of olivine and pyroxene, with sulfide spherules also present. A light inclusion is a brecciated intergrowth of compositionally zoned olivine and low-Ca pyroxene. Si, Al-rich glass occurs in the interstitial areas of the rock and commonly contains sulfide spherules. The sample contains abundant chondrules with a chondrule/matrix ratio typical of L chondrites. The mean chondrule size is  $\sim 0.7$  mm. The petrologic type is probably very low.

**Composition:** Olivine ( $\text{Fa}_{0.4-29.5}$ ), pyroxene ( $\text{Fs}_{2.8-23.9}$ ).

**Classification:** Ordinary chondrite (L3); S2, W2.

**Specimens:** One 24.5 g type specimen and one polished thin section are on deposit at *MNB*. Stefan Ralew of Germany holds the main mass.

#### Northwest Africa 4042

Find: 2004

Achondrite (ungrouped)

**History:** A single stone weighing 56.2 g was found by an anonymous finder in the western part of the Sahara in Morocco.

**Petrography, Composition, and Classification:** (A. Greshake, *MNB*) The rock is composed of mostly equigranular olivine ( $\sim 93.3$ ), somewhat larger low-Ca pyroxene ( $\sim 4.6$ ), Fe,Ni metal (0.7), pyrrhotite ( $\sim 0.4$ ), and Mg-Al-Ti chromite ( $\sim 1.1$ ; all vol%).

**Geochemistry:** Olivine ( $\text{Fa}_{20.3}$ ;  $\text{Cr}_2\text{O}_3$  and CaO between 0.03–0.07 wt%), pyroxene ( $\text{Fs}_{16.2}\text{Wo}_{1.0}$ ). Total C = 834 ppm (I. Franchi and R. C. Greenwood, *OU*). Oxygen isotopes: (I. Franchi and R. C. Greenwood, *OU*)  $\delta^{17}\text{O} = 2.539$ ,  $\delta^{18}\text{O} = 5.178$ ,  $\Delta^{17}\text{O} = -0.154$  (all ‰;  $n = 5$ ) plot close to aubrites, brachinites, and winonaites; however, no unambiguous assignment to a particular meteorite class can be made.

**Classification:** Achondrite (ungrouped); low shock.

**Specimens:** One 11.6 g type specimen and one thin section are on deposit at *MNB*. Stefan Ralew of Kunibertstraße 29, 12524 Berlin, Germany, holds the main mass.

#### Northwest Africa 4123

Northwest Africa

Find: 2004

Achondrite (eucrite, polymict)

**History:** A single stone of 46.9 g was found by an anonymous finder in the western part of the Sahara in Morocco and purchased in 2004 in Erfoud.

**Petrography:** (A. Greshake, *MNB*) Polymict breccia with basaltic, impact melt, and mineral clasts set into a fine-grained matrix of exsolved Ca pyroxene, plagioclase, and opaques. Large mineral clasts are present and are predominantly plagioclase and exsolved low-Ca pyroxene. Exsolution lamellae in pyroxene are generally very fine. Minor phases include silica and ilmenite.

**Geochemistry:** Plagioclase ( $\text{An}_{90.3}$ ), pyroxene ( $\text{Fs}_{39.1-55.0}\text{Wo}_{2.3-17.4}$ ).

**Classification:** Achondrite (eucrite, polymict); moderate shock, minimal weathering.

**Specimens:** A 10 g type specimen is on deposit at *MNB*. Mr. Christian Anger of Austria holds the main mass.

#### Northwest Africa 4215

Mhamid, Morocco

Find: April 2002

Achondrite (diogenite, unbrecciated)

**History:** This single stone was bought by Bruno Fectay and Carine Bidaut in Mhamid, Morocco, in April 2002.

**Physical Characteristics:** A single brown stone weighing 46.4 g and displaying limited patches of fusion crust.

**Petrography:** (J. A. Barrat, M. Bohn, *UBO-IUEM*; P. Beck, Ph. Gillet, *ENSL*) The sample displays a well-preserved cumulative texture, consisting of zoned xenomorphic orthopyroxene grains on the order of 500  $\mu\text{m}$  in size, with a few large chromite crystals (<5 vol%, up to 3 mm). Accessory olivine and scarce diopside grains occur within the groundmass, usually around the chromite crystals. Minor phases are cristobalite (determined by Raman spectrometry), troilite, and metal. This meteorite is weathered and its fractures are filled by calcite, limonite, and gypsum typical of hot desert alteration.

**Geochemistry:** (J. A. Barrat, M. Bohn, J. Cotten, *UBO-IUEM*; R. Greenwood, I. Franchi, *OU*) Orthopyroxenes ( $\text{En}_{76.2}\text{Wo}_{1.1}\text{Fs}_{22.7}$  to  $\text{En}_{68.6}\text{Wo}_{5.5}\text{Fs}_{25.9}$ ); olivines ( $\text{Fo}_{76}$  to  $\text{Fo}_{71}$ ); chromites ( $\text{Mg\#} = 14.3\text{--}44.0$ ,  $\text{Cr\#} = 42.2\text{--}86.5$ ) are chemically zoned (EMP). The bulk composition of this stone has been determined for major and trace elements (ICP-AES, ICP-MS). Its FeO, CaO abundances and most of the trace element concentrations (Sr, Ba, Pb, and REE among others) are high and indicate a significant contribution of the secondary minerals (limonite+calcite). In order to remove the terrestrial contribution, a subsample has been leached with hot HCl. The residue, made essentially of orthopyroxene and chromite, is similar in major and trace element abundances to diogenites as shown by the shape of its REE pattern and by its high Al/Ga ratio. Oxygen isotopic: (R. Greenwood, I. Franchi, *OU*) Were determined on a fraction of the leached powder,  $\delta^{17}\text{O} = 1.431 \pm 0.102$ ,  $\delta^{18}\text{O} = 3.203 \pm 0.205$  relative to V-SMOW, and  $\Delta^{17}\text{O} = -0.248 \pm 0.005$  (all ‰).

**Classification:** Achondrite (diogenite, unbrecciated).

**Specimens:** A total of 10 g type specimen and two polished thick sections are on deposit at *ENSL*. Bruno Fectay and Carine Bidaut of La mémoire de la Terre hold the main mass.

### THE AMERICAS

#### North America

##### Lucerne Valley 028

34°29'25"N, 116°57'34"W

San Bernardino County, California, USA

Find: 11 October 2003

Carbonaceous chondrite (CK4)

**History:** Two small dark gray stones weighing 3.0 g (LV 028) and 10.1 g (LV 029) were found within 100 meters of each other by R. Matson while searching for meteorites on Lucerne Dry Lake. Six additional fragments (LV 030, 031, 032, 035, 036, and 037) were found during detailed searches in March 2004, scattered over a linear kilometer. Total mass is 36.61 g.

**Petrography:** (A. E. Rubin, *UCLA*) The meteorites contain ~15 vol% chondrules with a mean apparent diameter of ~500  $\mu\text{m}$ . Chondrule types include BO, PO, and POP. The principal opaque phases are magnetite and pyrrhotite. The magnetite occurs both as small grains and as clumps as large as 250  $\mu\text{m}$ . The pyrrhotite is moderately abundant. Olivine ( $\text{Fa}_{33.2 \pm 0.4}$ ), groundmass of 25–50  $\mu\text{m}$ .

**Classification:** The stones are carbonaceous (CK4); moderate shock.

**Specimens:** A 7.33 g type specimen is on deposit at *UCLA*. Matson holds the main mass.

##### Mohawk

32°43'8"N, 113°42'6"W

Yuma County, Arizona, USA

Find: October 2000

Iron (IAB complex)

**History:** One mass of 586 g found in the desert near Mohawk, Arizona, by Don Armijo.

**Geochemistry:** (D. Hill and D. A. Kring, *UAz*) Bulk composition: Fe =  $88.9 \pm 0.7$ , Co =  $0.443 \pm 0.003$ , Ni =  $7.41 \pm 0.7$  (all wt%); Ir = 2.42, Au 1.6, Ga = 76, Ge =  $292 \pm 28$  (all ppm) as analyzed by INAA.

**Classification:** Iron (IAB complex) with kamacite bandwidth of  $2.8 \pm 1.1$  mm (coarse octahedrite).

**Specimens:** A 37.4 g type specimen is on deposit at *LPL*. The main mass is held by Don Armijo.

##### Nova 005

Location is unconfirmed within the Americas

Find: 2002

Ordinary chondrite (L5)

**History:** One mass of 242 g was found. There have been a variety of explanations of how and where this meteorite was found. The current version involves an 80-year-old man as having found this stone while hunting for obsidian artifacts in the Cactus Flat-Coso Junction-Red Hill-Fossil Falls area (the east side of Rose Valley, California). The find location has been alternatively represented as Owens Lake, then Rose Valley, and most recently Fossil Falls.

**Physical Characteristics:** (A. Rubin, *UCLA*) The stone was found with 100% black fusion crust with a few spots of a rust-like color. The interior is cream-colored and sparsely speckled with orange spots.

**Petrography:** The sample is ~30 vol% chondrules and 70 vol% matrix. Most sulfide and metal grains appear slightly oxidized. Olivine ( $\text{Fa}_{24.2}$ ). Note: The terrestrial age (Battelle Labs) is  $25 \pm 6$  yr.

**Classification:** Ordinary chondrite (L5); S1, W1.

**Specimens:** A 41.2 g sample is on deposit at *UCLA*. A total of ~200 g is held by Meteorite Recovery Lab.

**Orlando** **28°32'51"N, 81°21'44"W**

Orange County, Florida, USA

Fall: 8 November 2004

Achondrite (eucrite, monomict)

**History:** On Monday, November 8, 2004, around 6:15 P.M., Ms. Donna Shuford was startled by the noise of something hitting the side of her house. She discovered that something had hit the top of her car and ricocheted onto the side of her house. A single ~180 g stone that had fragmented on impact was found.

**Petrography and Geochemistry:** (D. Mittlefehldt and M. Zolensky, *NASA JSC*) Major phases are low-Ca pyroxene ( $\text{Wo}_3\text{En}_{35}\text{Fs}_{62}$ ; Fe/Mn ~30) with lamellae of high-Ca pyroxene ( $\text{Wo}_{45}\text{En}_{29}\text{Fs}_{26}$ ), and calcic plagioclase ( $\text{An}_{71-83}\text{Ab}_{16-28}\text{Or}_{-1}$ ). Minor phases include titanian chromite ( $\text{TiO}_2 = 16-20$ ,  $\text{Al}_2\text{O}_3 = 2-3$ ,  $\text{MgO} = 0.4$ ,  $\text{MnO} = 0.8$ ; [all wt%]), ilmenite ( $\text{MgO} = 0.5$ ,  $\text{MnO} = 0.9$  [both wt%]), with silica, iron sulfide, and Fe,Ni metal. The rock is largely unbrecciated, but has shock veins with granular texture and containing some glass. Remnant ophitic/subophitic igneous texture is preserved with plagioclase laths ~1 mm by ~30  $\mu\text{m}$ , and ~2 mm blocky pyroxene grains. In much of the rock, pyroxene has been recrystallized to ~20–50  $\mu\text{m}$  equant grains while plagioclase retains its original shape.

**Classification:** Achondrite (eucrite, monomict).

**Specimens:** A 20 g type specimen is on deposit at *SI*. The finder holds the main mass.

**Pedernales** **30°20'N, 98°57'W**

Gillespie County, Texas, USA

Find: 1 December 1980

Iron (ungrouped and related to IAB clan)

**History:** Mr. John Stitt excavated a mass of 691 g from a depth of 4 feet from an Indian rock midden on ranch land close to Fredericksburg.

**Geochemistry:** (J. T. Wasson, *UCLA*) Bulk composition: Co = 4.89, Ni = 75.0 (both mg/g); Ga = 65, Ge = 190, As = 16.4, Ir = 1.17, Au = 1.455 (all  $\mu\text{g/g}$ ).

**Classification:** The stone is an iron of the IAB complex, but it is not a member of the IAB main group or the subgroups. It is closely related to Algarrabo and slightly further away in composition from Livingston (Tennessee). The kamacite bandwidth is  $0.9 \pm 0.2$  mm (medium octahedrite). In some areas such bands have merged to create bands twice as wide.

**Classification:** Iron (ungrouped and related to IAB clan).

**Specimens:** A 23 g type specimen is on deposit at *UCLA* and one 86 g type specimen is on deposit at *ASU*. The main mass is held by B. Barnett.

**Purmela** **29°30'N, 98°3'W**

Coryell County, Texas, USA

Find: 10 January 1977

Iron (IIF)

**History:** Calvin Perryman recovered one 4.5 kg mass on a farm road.

**Petrography and Geochemistry:** (M. McGehee, *ASU*; J. Wasson, *UCLA*) Metal composition determined by INAA, Ni = 107, Co = 6.2 (both mg/g), Ga = 14, As = 4.3, Ir = 11 (all  $\mu\text{g/g}$ ). Discontinuous kamacite spindles (<2 mm wide) are visible on an etched surface. The metal also contains troilite nodules (<1 mm to ~18 mm wide).

**Classification:** Iron IIF, plessitic octahedrite.

**Specimens:** A 406.3 g type specimen is on deposit at *ASU*. M. Jones holds the main mass.

**San Pedro Jacuaro** **19°46'N, 100°39'W**

Estado de Michoacan de Ocampo, Mexico

Fall: 1 December 1968

Ordinary chondrite (LL6)

**History:** In December 1968, Jose Dimas Bautista of San Luis Potosi, Mexico, heard a loud "airplane type" sound while working outside the mines in the Sierra Madres mountains near San Pedro Jacuaro. A search recovered a 460 g mass from a fresh hole in the sand. He kept it in his possession until July 2003, when he brought it to the attention of Robert Cucchiara of Alameda, California, who recognized it as a meteorite.

**Geochemistry:** (P. Sipiera, Harper, K. Cole, *Univ. Illinois*) Olivine ( $\text{Fa}_{27.7}$ ), pyroxene ( $\text{Fs}_{23.2}\text{Wo}_{2.0}$ ).

**Classification:** Ordinary chondrite (LL6); S2, W0.

**Specimens:** A 20 g type specimen is on deposit at *PSF*. Robert Cucchiara holds the main mass.

**Superior Valley 014** **35°14'160"N, 117°02'527"W**

San Bernardino County, CA, USA

Find: 25 August 2002

Achondrite (acapulcoite)

**History:** A 1.77 g specimen was found by Jason Utas.

**Petrography:** (A. Rubin, *UCLA*) The rock has a granular texture consisting mainly of quasi-equant olivine and low-Ca pyroxene grains from 60 to 500  $\mu\text{m}$  in size and averaging ~160  $\mu\text{m}$ . Accessory plagioclase grains ~200  $\mu\text{m}$  with polysynthetic twinning are also present. In some areas the silicate grains are densely packed and many grain boundaries meet at 120° triple junctions. In other areas the silicate grains are separated and surrounded by limonite. The separated silicate grains range in shape from fragmental to subrounded to subhedral. No relict chondrules were observed. Some silicate grains contain 100–300  $\mu\text{m}$  long curvilinear trails of 2–4  $\mu\text{m}$  chromite blebs. The rock is very weathered and contains abundant limonite and patches of clay phases. Nevertheless, a few patches of troilite and rare grains of metallic Fe,Ni are present.

**Geochemistry:** Olivine ( $\text{Fa}_{4.6}$ ), pyroxene ( $\text{Fs}_{6.9 \pm 0.1}\text{Wo}_{2.9 \pm 1.6}$ ).

**Classification:** Achondrite (acapulcoite); moderate shock.

**Specimens:** A 0.49 g type specimen is on deposit at *UCLA*. The main mass is held by Jason Utas.

## ANTARCTICA

## Grove Mountains

**Grove Mountains (GRV) 024516**      **73°00'S, 75°12'E**

Antarctica

Find: 2003

Achondrite (ureilite)

**History:** A stone of 24.7 g was found within a moraine during the 19th Chinese Antarctic Research Exploration in Grove Mountains.

**Physical Characteristics:** (Miao B., Xu L., Lin Y., *IGGCAS*) It is brownish grey, with some yellow and dark grey spots and no fusion crust.

**Petrography:** It shows typical ureilite texture consisting mainly of olivine and pigeonite with minor carbonaceous matrix (graphite and diamond were identified by Raman). Triple junctions with an angle of 120° are common among coarse-grained silicates.

**Geochemistry:** Olivine ( $\text{Fo}_{84.0 \pm 0.4}$  with homogeneous cores), pigeonite ( $\text{En}_{77.1 \pm 0.4}\text{Wo}_{9.3 \pm 0.1}\text{Fs}_{13.6 \pm 0.4}$ ). Olivine grains have reduced rims ( $\text{Fa} \leq 5.4$  mol%) that contain abundant fine-grained inclusions of Ni-poor metal and sulfide. Limonite veins are common.

**Classification:** Achondrite (ureilite); moderate shock, extensive weathering.

**Specimens:** The entire specimen is on deposit at the *GUT*.

## ASIA

## People's Republic of China

**Fukang**      **44°26'N, 87°38'E**

Fukang, Xinjiang Province

Find: 2000

Pallasite (main group)

**History:** An anonymous finder recovered a 1003 kg specimen near Fukang, China, in 2000. The sample was at the Tucson Gem and Mineral Show in February 2005, and seen by D. S. Lauretta of *UAz*. Approximately 20 kg had been removed from the main mass by the finder before the Tucson show and the mass investigated at *UAz* was 983 kg. (D. S. Lauretta, D. Hill, M. Killgore, D. Della-Giustina, Y. Goreva, *UAz*; I. Franchi, *Open U*).

**Petrography and Geochemistry:** Olivine: Throughout the large mass, olivines vary in shape from rounded to angular; many are fractured. They range in size from <5 mm to several cm. The main pallasite contains several regions of "massive" olivine clusters up to 11 cm in diameter with thin metal veins only a few millimeters in width.  $\text{Fo}_{86.4}$  with molar  $\text{Fe/Mg} = 0.1367$ ,  $\text{Fe/Mn} = 40.37$ , and  $\text{Ni} = 0.03$  wt%. Zoning was not observed for Al, Cr, Ca, Mn, or Fe typical of olivines in main-group pallasites. Metal and sulfides: Groundmass is mostly kamacite with some occurrences of kamacite mantles surrounding taenite cores, rounded taenite adjacent to

kamacite, and regions of "comb plessite." Kamacite contains an average  $\text{Ni} = 6.98$  wt%. Schreibersite is enclosed by wide kamacite bands and as mantles adjacent to olivines. Two populations of schreibersite are present with  $\text{Ni} = 26$  and 35 wt% near chromite. Vermicular sulfide (troilite) is present in some olivine. Thin veins of kamacite and troilite occur inside many olivines as well. Minor phases: Euhedral chromites up to 0.5 cm, rounded whitlockite adjacent to olivine, and troilite heterogeneously distributed in thin veins. Several regions, ranging from <100  $\mu\text{m}$  to several millimeters, that contain a complex mixture of olivine, low-Ca pyroxene, troilite, and whitlockite were observed adjacent to chromite. Bulk composition:  $\text{Fe} = 89.9 \pm 0.3$ ,  $\text{Ni} = 9.0 \pm 0.2$ ,  $\text{P} = 0.62 \pm 0.02$ ,  $\text{Co} = 0.51 \pm 0.01$  (all wt%);  $\text{Ge} = 41 \pm 4$ ,  $\text{As} = 26 \pm 5$ ,  $\text{Ga} = 19.1 \pm 0.5$ ,  $\text{Pd} = 5.1 \pm 0.2$ ,  $\text{Au} = 2.6 \pm 0.2$ , (all  $\mu\text{g/g}$ );  $\text{Ir} = 43 \pm 4$  ng/g. Oxygen isotopes:  $\delta_{18}\text{O} = 2.569$ ,  $\delta^{17}\text{O} = 1.179$ ,  $\Delta^{17}\text{O} = -0.157$  (all ‰).

**Classification:** Pallasite (main group).

**Specimens:** A total of 31 kg of type specimen is on deposit at *UAz*. M. Killgore holds a total of 31 kg. An anonymous collector holds the main mass.



Fig. 1. Dante Lauretta with the main mass of Fukang at the University of Arizona.

**Ulasitai**      **44°57'24"N, 91°24'09"E**

Mulei County, Xinjiang Province

Find: 28 April 2004

Iron (IIIIE)

**History:** A single iron meteorite weighing 430 kg was found during fieldwork by Mr. Xiaodong Li, a geologist. The meteorite was discovered on the hillside in the Mountain Beita area, Mulei county, Xinjiang Province.

**Physical Characteristics:** (Miao B., Xu L., Lin Y., *IGGCAS*) The meteorite has an angular shape with cm-scale concaves on its surface and is dark brown with a sub-cm layer of limonite on its bottom, but with no fusion crust remaining.

**Petrography and Geochemistry:** Kamacite ( $\text{Ni} = 6.87\text{--}7.39$  Co = 0.47–0.75 [both wt%]), taenite ( $\text{Ni} = 11.6\text{--}36.7$  Co = 0.22–0.65 [both wt%]), Widmanstätten pattern with 0.6–

2.0 mm (average 1.14 mm) kamacite bandwidths. Bulk composition: Ni = 10.03 wt% (by ICP-AES), and trace elements (by ICP-MS, in  $\mu\text{g/g}$ ) Ge = 31.28, Ga = 16.09, Ir = 0.22. Minor phases include schreibersite, cohenite, troilite, and chromite.

**Classification:** Iron (IIIE). Note: This meteorite may be paired with Armanty.

**Specimens:** A 530 g type specimen is on deposit at *GUT* and 730 g at *IGGCAS*. The finder holds the main mass.

## EUROPE

### Germany

#### Königsbrück 51°16'0"N, 13°54'0"E

Königsbrück, Saxony, Germany

Find: May 2004

Ordinary chondrite (H/L4)

**History:** A complete oriented stone weighing 51.8 g, partly covered with fusion crust, was found by an anonymous finder on a field close to the village of Königsbrück, Saxony, Germany, during a search for moldavites.

**Petrography:** (A. Greshake and M. Kurz, *NHB*) The sample is a fresh, unbrecciated stone with equilibrated olivine and unequilibrated pyroxene.

**Geochemistry:** Olivine ( $\text{Fa}_{22.6}$ ), pyroxene ( $\text{Fs}_{8.2-20.1}$ ).

**Classification:** Ordinary chondrite (H/L4); S4, W1.

**Specimens:** A 11.7 g type specimen plus one thin section are on deposit at *MNB*. An anonymous finder holds the main mass.

#### Pallasovka 49°52'0"N, 46°36'7"E

Volgograd district, Russia

Find: July 1990

Pallasite

**History:** One stone weighing 198 kg was found by N. F. Kharitonov at a shore of an artificial water reservoir, 27.5 km from the town of Pallasovka. Interestingly, the town was named after Peter Pallas (1741–1811), a famous naturalist who took part in the discovery and the first study of the Pallas Iron Mass, which was found near Krasnojarsk in 1749 and gave the name of the pallasite meteorite group. A. E. Milanovsky transferred a sample of the meteorite to the Vernadsky Institute, Moscow.

**Petrography and Geochemistry:** (M. A. Ivanova, N. N. Kononkova, *Vernad*; S. E. Borisovsky, Institute of Geology of Ore Deposits, Mineralogy, Moscow)

**Petrography:** The stone consists of approximately equal parts of olivine and metal, and has abundant brown, rusty fusion crust with regmaglypts.

**Geochemistry:** Olivine (mg# 87.7, Fe/Mn = 45.2, Fe/Mg = 0.14; similar to main group pallasites). Metal (bulk ICP AES) Ni = 13.1 wt%; Ir = 0.12, Au = 2.8, Pt = 3.2, Ga = 22.5, Ge = 24.9 (all ppm). Kamacite (Co = 0.61, Ni = 7.21; both wt%) and taenite (Co = 0.35, Ni = 26.5 both wt%). Additional phases: Troilite (Ni = 0.41 wt%), schreibersite, and chromite

(mg# 36; Fe/Mn = 48.9; Cr/(Cr+Al) = 77.2). Note: Chromites in this sample differ in composition from that of the main and Eagle Station groups.

**Classification:** Pallasite (Main group).

**Specimens:** A 9336 g sample and one polished section are on deposit at *Vernad*. The main mass is held by an anonymous purchaser.

## MIDDLE EAST

### Oman, Dhofar

#### Dhofar 1277 19°10'84"N, 54°25'13"E

Oman

Find: 8 March 2005

Carbonaceous chondrite (CM2)

**History:** One mass of 10 g was found in March 2005 in the Dhofar region of Oman.

**Physical Characteristics:** (M. Ivanova, *Vernad*; F. Brandstaetter, *NHWW*) The sample has some dark brown fusion crust.

**Petrography:** The meteorite consists of altered POP, rare BO chondrules and chondrule fragments, olivine aggregates with halos around, and fragments of a nonhydrated matrix material set within a phyllosilicate matrix. Scarce isolated olivine and pyroxene grains and refractory inclusions consisting of altered silicates, perovskite, and spinel are present.

**Geochemistry:** Olivine ( $\text{Fa}_{0.4-61}$ ; CaO = 0.26,  $\text{Cr}_2\text{O}_3$  = 0.36 [both wt%]), low-Ca pyroxene ( $\text{Fs}_{1.3-7.2}\text{Wo}_{0.5-3.2}$ ), high-Ca ( $\text{Fs}_{21}\text{Wo}_{27.5}$ ). Minor phases are tochilinite, kamacite, pyrrhotite, pentlandite, Cr, P-rich sulfides, chromite, and Ca carbonates.

**Classification:** Carbonaceous chondrite (CM2); S1.

**Specimens:** A 3.2 g type specimen and one thin section are on deposit at *Vernad*. The main mass is with anonymous finder.

#### Dhofar 1285 19°18'15"N, 54°33'3"E

Oman

Find: 11 December 2002

Achondrite (ureilite)

**History:** Three dark brown stones, broken into fragments of total weight 406 g, were found on 11 December 2002 in the desert of Oman. The fragments were separated by ~2.5 km.

**Petrography:** (Lorenz, *Vernad*; Brandstätter, *NHWW*) A coarse-grained rock that consists mainly of olivine with minor amounts of pyroxene. The veins of troilite and Fe,Ni metal crosscut the whole rock. Accessory phases, graphite, and daubreelite are present.

**Geochemistry:** The cores of olivine grains ( $\text{Fo}_{76.6}$ ; Fe/Mn = 59) are surrounded by reduction rims (~50  $\mu\text{m}$  in width), which contain small Fe-rich metal grains. The composition of olivine within the reduction rims gradually changes from  $\text{Fo}_{78}$ ; Fe/Mn = 51 on the inside towards  $\text{Fo}_{96}$ ; Fe/Mn = 8.6 at the outer edge. The outer parts of the olivine grain rims are

completely replaced by a fine-grained aggregate of olivine (Fo<sub>97-99</sub>), troilite, and metal. Reduction zones are developed along cracks and veins that cross-cut olivine grains. Pyroxene (En<sub>78.7</sub>Wo<sub>9.8</sub>; Fe/Mn = 20) contains numerous, tiny, co-oriented inclusions of Ca-rich pyroxene (En<sub>56.2</sub>Wo<sub>37.3</sub>; Fe/M = 13), SiO<sub>2</sub>-rich feldspathic (Ab<sub>50</sub>An<sub>48.5</sub>) glass, low-Ni Fe,Ni metal grains, and extremely fine, poorly resolvable exsolution lamellae of Ca pyroxene. In places, pyroxene grains also have reduction rims with composition En<sub>83</sub>Wo<sub>6.7</sub>. Cr<sub>2</sub>O<sub>3</sub> contents of olivine and pyroxene (0.7 and 1.1 wt% respectively) and CaO content of olivine = 0.4 wt% are in the ranges of those established for ureilites.

**Classification:** Achondrite (ureilite); moderate shock, extensive weathering.

**Specimens:** A 92.48 g type specimen and one thin section are on deposit at *Vernad*. An anonymous finder holds the main mass.

#### Dhofar 1286

18°25'579"N, 54°25'719"E

Oman

Find: December 2005

Achondrite (eucrite, polymict)

**History:** Two pieces of meteorite weighing 898 g in total were found on a sandy surface in the desert of Oman. The distance between the fragments was about 30 m. Joined together, the fragments form an almost complete individual sample with ~10% missing.

**Petrography:** (Lorenz, *Vernad* and Brandstätter, *NHMW*) On the unbroken surfaces, black fusion crust is partly preserved. The meteorite is a polymict breccia consisting of fragments of medium- to coarse-grained metamorphosed gabbroic rocks and fine- to medium-grained subophitic basaltic rocks. Minor equigranular metamorphic pyroxene-feldspar rocks, melts, and melt matrix breccias are present. The rock fragments are situated in a fine-grained clastic matrix and comprise 60 vol% of the whole meteorite. Accessory minerals are troilite, chromite, ilmenite, silica, Ca phosphate, and metal Fe,Ni.

**Geochemistry:** Pyroxene (En<sub>35.1-62.8</sub>Wo<sub>1.8-5.7</sub>; Fe/Mn = 30) with exsolved augite (En<sub>21.1-54.2</sub>Wo<sub>21.1-38.9</sub>).

**Classification:** Achondrite (eucrite, polymict).

**Specimens:** A 20.6 g type specimen and one polished section are on deposit at *Vernad*. An anonymous finder holds the main mass.

#### Sayh al Uhaymir 402

21°4'37"N, 57°16'11"E

Oman

Find: 2004

Achondrite (ungrouped enstatite)

**History:** A meteorite collector recovered a single stone was found in the desert of Oman.

**Physical Characteristics:** The stone, weighing 78 g, has a brown surface color and is free of fusion crust.

**Petrography:** (D. D. Badjukov, *Vernad*; F. Brandstaetter, *NHMF*) The meteorite consists of <2 millimeter-size silicate

clasts embedded in an iron oxide groundmass; the silicate clasts are composed of fine (0.002–0.5 mm, average 0.05 mm) fragments of enstatite crystals in a matrix consisting of aggregates of a few millimeters wide plagioclase laths in a transparent glass. Fine droplets and blebs of metal, ninigerite, and troilite are dispersed in the glass. Silicate mode (vol%) for enstatite and matrix is 63% and 37%, respectively. Mineral chemistry: Enstatite (Fs<sub>1</sub>Wo<sub>1</sub>), glass (SiO<sub>2</sub> = 70–75, Al<sub>2</sub>O<sub>3</sub> = 17–18, MgO = 4–5, Na<sub>2</sub>O = 5–6, FeO and K<sub>2</sub>O < 1 [all wt%]), metal (Si = 2 wt%), ninigerite (Ca = 1.3, Cr = 1.8 [both wt%]); troilite contains Cr, Ti, and Mn.

**Classification:** Achondrite (ungrouped enstatite). The texture and mineralogy imply that it could be an impact melt rock; relic fragments of enstatite show strong mosaicism and planar features. The meteorite is permeated by iron oxides (45 vol%) but the silicate clasts are unaltered.

**Specimens:** A 17.0 g type specimen and one thin section are on deposit at *Vernad*. An anonymous collector holds the main mass.

### METEORITES FROM OTHER PLANETARY BODIES

#### Meridiani Planum, Mars

1°56'46"S, 354°28'24"E

MER landing site, Meridiani Planum, Mars

Find: 5 January 2005

Iron (IAB complex)

**History:** While exploring the remnants of the heat shield on Meridiani Planum, the Mars Exploration Rover (MER) Opportunity imaged the surroundings with its panoramic camera (PanCam). On sol 324, a close-by object with a maximum dimension of 31 cm across became for the first time clearly visible in one of these images. Subsequent PanCam images reveal a smooth rock surface covered by depressions partly reminiscent of regmaglypts.

**Classification and Description:** (Athena Science Team) Spectra obtained by the Miniature Thermal Emission spectrometer on sol 339 and 342 show a thermal emissivity of 0.35, which is only consistent with metal indicating that the object was an iron meteorite. Detailed investigations from sol 349 to 352 by the in situ instruments on the rover's arm, the Rock Abrasion Tool (RAT), Microscopic Imager (MI), Mössbauer Spectrometer (MB), and Alpha Particle X-Ray Spectrometer (APXS), confirmed the metallic nature and allowed classification. Classification based on analyses of a brushed surface: MB, ~94% of the Fe is bound in kamacite, and APXS, dust corrected composition of iron is Ni = ~7 wt%; Ge = ~300, Ga = <70 (both ppm), consistent with its classification as an iron (IAB complex).

**Specimens:** Type specimen and main mass, Mars.

### ERRATA

1. In Meteoritical Bulletin No. 87, the recovery date of Tanezrouft 059, Tanezrouft 060, and Tanezrouft 061 was

incorrectly reported. These meteorites were recovered on 16 May 2002 with Tanezrouft 056 and Tanezrouft 057.

2. In Meteoritical Bulletin No. 88, the meteorite Cheder is classified as an iron (IIIAB), but it is actually an iron (IID).
3. In Meteoritical Bulletin No. 89, Table 7, two meteorites are listed under the name NWA 1392. The first one, with a mass of 1091 g, should have been named NWA 1391. The second one, with a mass of 64 g, is the assigned NWA 1392.

*Acknowledgments*—The expertise of R. Korotev has been most helpful, and we are all grateful to him for his generosity. The editor is especially grateful to M. Weisberg, C. Goodrich, and D. Ebel for their constant support.

#### ABBREVIATIONS FOR CLASSIFIERS, ANALYSTS, AND TYPE SPECIMEN LOCATIONS

A list of type specimen locations, classifiers, and analysts referenced from the tables of approved meteorites. The abbreviations appear in the location column of the tables. Unless specifically noted, all type specimens are at the home institution of the first listed analyst and main masses are with anonymous finders. A PDF file with a key to the abbreviations used for each institute listed in italics below and throughout the Meteoritical Bulletin is available for download at our home page: <http://www.meteoriticalsociety.org/bulletin/institutions.pdf>. Unless specifically noted, all main masses are located with an anonymous finder.

ASU1: Type specimen on deposit at *ASU*, L. Bleacher, classifier. M. Miller holds the main mass.

ASU1-1: Type specimen on deposit at *ASU*, L. Bleacher, classifier. J. Gwilliam holds the main mass.

ASU1-2: Type specimen on deposit at *ASU*, L. Bleacher, classifier. B. Southern holds the main mass.

Bart1: Type specimen on deposit at *Vernad*. R. Bartoschewitz, classifier. Purchased U. Eger. *Vernad*; specimen, Bart; main mass, U. Eger.

Bart2: Type specimen on deposit at *MPI*. F. Wlotzka (*MPI*) and R. Bartoschewitz, classifiers. Purchased Fectay, the main mass with Bart.

Bart3: Type specimen on deposit at *Vernad*. R. Bartoschewitz, classifier. Purchased Fectay, the main mass with Bart.

CML1-1: Type specimen on deposit at *CML*. K. R. Carroll classifier under the mentorship of M. Hutson.

CML1-2: Type specimen on deposit at *CML*. M. Hutson (*CML*), classifier. L. Sloan holds the main mass.

IfP1: Type specimen on deposit at *Mun*, A. Sokol, Meimeier, and A. Bischoff, classifiers.

IfP2: Type specimen on deposit at *Mun*, Niemeier, Jording, and A. Bischoff, classifiers.

LPL1: Type specimen on deposit at *LPL*, D. Hill, classifier. E. Ghior holds the main mass.

MIN-KB1: Type specimen on deposit at *MIN-KB*, R. Bartoschewitz and P. Appel, classifiers. R. Bartoschewitz holds the main mass.

MNB1-1: Type specimen on deposit at *MNB*, A. Greshake and M. Kurz, classifiers.

MNB1-2: Type specimen on deposit at *MNB*, A. Greshake, classifier.

MNB1-3: Type specimen on deposit at *MNB*, A. Greshake, classifier. The main mass held by JNMC-Zürich, Switzerland.

MNB1-4: Type specimen on deposit at *MNB*, A. Greshake, classifier. The main mass is held by HSSH-Primasens, Germany.

NAU1: Type specimen on deposit at *NAU*, J. Wittke and T. Bunch, classifiers.

NHNV1: Type specimen on deposit at *NHNV* (inventory # M6694). C. Koeberl (*NHNV*) and J. Wasson (*UCLA*), classifiers.

MNB1-5: Type specimen on deposit at *MNB*, A. Greshake and M. Kurz (*MNB*), classifiers.

MNHNP-1: Type specimen on deposit at *MNHNP* and main mass with anonymous finder. M. Benise and B. Zanda (*MNHNP*), classifiers.

MSP1: Type specimen on deposit at *MSP-PO* and the main mass with an anonymous finder. V. Moggie Cecchi, A. Salvadori, and G. Pratesi (*MSP*), classifiers.

NUM1: Type specimen on deposit at *NUM*, M. Kimura, classifier.

SHIM1: Type specimen on deposit at *Shim*, C. Fukuda, classifier.

SHIM2: Type specimen on deposit at *Shim*, C. Nishina, classifier.

SI1-MZ: Type specimen on deposit at the *SI*, D. Mittlefehldt and M. Zolensky, classifiers. D. Shuford holds the main mass.

UCLA1: Type specimen on deposit at *UCLA*, A. Rubin, classifier. M. Matson holds the main mass.

UCLA1-1: Type specimen on deposit at *UCLA*, A. Rubin, classifier. P. Utas holds the main mass.

UCLA1-2: Type specimen on deposit at *UCLA*, A. Rubin, classifier. J. Utas holds the main mass.

ULCA2/LPL: Type specimens are on deposit at *UCLA* and *LPL*, J. Wasson, classifier.

UNM1: Type specimen on deposit at University of New Mexico, R. Jones, classifier. J. Pringle holds the main mass.

UWS1: Type specimen on deposit at *UWS*, A. Irving and S. Kuehner, classifiers.

Vernad1-1: Type specimen on deposit at *Vernad*, M. Ivanova, classifier.

Vernad1-2: Type specimen on deposit at *Vernad*, M. Ivanova, classifier and N. Kononkova, analyst.

Vernad1-3: Type specimen on deposit at *Vernad* and *MSU*, M. Ivanova of Vernad, classifier and A. Ulianov of *MSU*, analyst.

Vernad2-1: Type specimen on deposit at *Vernad*, C. Lorenz of *Vernad*, classifier.

Table 1. Meteorites from specific locations within Africa.

| Name                         | Abbreviation     | Find/Place | Date of recovery (dd/mm/yyyy) | Latitude (N) | Longitude (E) | Mass (g) | Pieces | Class <sup>a</sup> | Shock | WG <sup>b</sup> | Fa (mol%) | Fs (mol%)  | Wo (mol%) | Magnetic susc. | Type spec. (g) | Location of type <sup>c</sup> | Comments                            |
|------------------------------|------------------|------------|-------------------------------|--------------|---------------|----------|--------|--------------------|-------|-----------------|-----------|------------|-----------|----------------|----------------|-------------------------------|-------------------------------------|
| Afer 341                     | Afer 341         | Find       | 05/12/2002                    | 27°28'       | 3°50'         | 132      | 7      | L3                 | S5    | W1              | 20.4 ± 4  | 14.8 ± 4.8 | —         | 4.84           | 20.5           | MNHN-1                        | —                                   |
| Afer 366                     | Afer 366         | Find       | 11/2002                       | 26°36.56'    | 03°56.14'     | 1456     | 1      | CH3                | —     | —               | —         | —          | —         | —              | All            | MSP1                          | See separate entry                  |
| Afer 374                     | Afer 374         | Find       | 11/2002                       | 26°36.52'    | 04°03.18'     | 118      | Many   | CO3                | —     | —               | —         | —          | —         | —              | All            | MSP1                          | See separate entry                  |
| Dar al Gani 1041             | DaG 1041         | Find       | 1998                          | 27°05.00'    | 16°03.30'     | 169      | 1      | CO3                | S2    | —               | 0.4–60.8  | 0.8–21.7   | —         | —              | 23.8           | MNB1-5                        | Mean diameter of chondrules ~140 µm |
| Goronyo Hammada al Hamra 337 | Goronyo HamH 337 | Find       | 31/10/2001                    | 13°16'       | 5°24'         | 11,000   | 1      | H4                 | S4    | W1              | 21.8      | 19.9       | —         | —              | 161            | UNMI                          | —                                   |
| Mafuta                       | Mafuta           | Find       | 12/01/1984                    | 16°54.09'    | 30°24.26'     | 71.5 kg  | 1      | Iron (IID)         | —     | —               | —         | —          | —         | —              | 221            | NHNV                          | See separate entry                  |

<sup>a</sup>Class = classification.<sup>b</sup>WG = weathering grade.

<sup>c</sup>For location of type: MNB1-5 = classified by A. Greshake, MNB, and M. Kurz; type specimen: MNB; main mass with anonymous finder. MNHN-1 = classified by B. Devouard and J.-L. Devidal, UPB, M. Denise and B. Zanda, MNHN; type specimen, MNHN; main masses with finders, F. Beroud and C. Boucher. MSP1 = classified by V. Moggi Cecchi, A. Salvadori, and G. Pratesi, MSP; type specimen, MSP-PO; main mass with anonymous finder. NHMV = classified by C. Koeberl, NHMV, and J. Wasson, UCLA; type specimen NHMV (inventory #M6694); main mass with finder, Mr. M. Alexander of Mafuta Farm, Zimbabwe. UNMI = classified by R. Jones, Institute of Meteoritics, University of New Mexico, 200 Yale N.E., Albuquerque, New Mexico 87131, USA; type specimen and thin section, Institute of Meteoritics, main mass, Twelker.

Table 2. Meteorites from Northwest Africa.

| Name                  | Recovery area (or purchase) <sup>a</sup> | Recovery date (dd/mm/yyyy) | TKM <sup>b</sup> (g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup> (g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa (mol%)  | Fs (mol%) | Location | Comments  |
|-----------------------|--|----------------------------|----------------------|---------------------|--------------------|--------------------------|-----------------|-----------------|------------|-----------|----------|---|
| NWA 869               | Unknown                                  | 2001 or 2002               | n kg                 | n                   | L4–6               | 189.3                    | S3              | W1              | 24.2       | —         | UCLA1    | —   |
| NWA 999               | Morocco                                  | 2000                       | 330                  | 1                   | Euc                | 20                       | L-M             | —               | —          | 68        | —        | See separate entry  |
| NWA 1006              | Morocco                                  | 27/08/2001                 | 24.5                 | 1                   | Ure                | 5.86                     | L               | —               | ~10        | —         | —        | See separate entry  |
| NWA 1460              | Morocco                                  | 05/1998                    | 70.2                 | 1                   | Martian            | Various                  | —               | Min             | —          | —         | —        | See separate entry  |
| NWA 1462              | Er-Mor                                   | 03/03/2002                 | 203                  | 1                   | Ure                | 20.2                     | Low             | —               | ~22        | —         | —        | See separate entry  |
| NWA 1617 <sup>h</sup> | Ag-Mor                                   | 06/1998                    | 21                   | 1                   | Acp                | 4.5                      | —               | —               | —          | —         | UWS1     | See separate entry in MB 88   |
| NWA 1918              | Morocco                                  | 01/2003                    | 136                  | 1                   | Euc                | 20.2                     | —               | —               | —          | —         | —        | See separate entry  |
| NWA 1923              | Morocco                                  | 03/2003                    | 112                  | 1                   | Euc                | 20.4                     | High            | —               | —          | —         | —        | See separate entry  |
| NWA 1929              | Er-Mor                                   | 05/2003                    | 922.2                | 1                   | How                | 22.02                    | High            | —               | —          | —         | NAU1     | See separate entry  |
| NWA 2040              | Er-Mor                                   | 2003                       | 1200                 | 1                   | LL3.1              | 20.3                     | S2              | W3              | 1.4–52.4   | —         | NAU1     | Olv (FeO/MnO = 44–61, Cr <sub>2</sub> O <sub>3</sub> = 0.32 ± 0.19) |
| NWA 2134              | Er-Mor                                   | 2004                       | 916                  | 1                   | H6                 | 24                       | S2              | W3              | 18.4       | 16.3      | NAU1     | —   |
| NWA 2135              | Er-Mor                                   | 2004                       | 421                  | 1                   | LL4                | 26.4                     | S2              | W4              | 29.4       | 24        | NAU1     | —   |
| NWA 2136              | Er-Mor                                   | 2004                       | 1045                 | 1                   | L3.5               | 22                       | S2              | W3              | 18.9 ± 6.5 | —         | NAU1     | Olv (Cr <sub>2</sub> O <sub>3</sub> = 0.11 ± 0.07)                  |
| NWA 2137              | Er-Mor                                   | 2004                       | 6174                 | 1                   | LL3.7              | 24.6                     | S2              | W2              | 23.5 ± 5.7 | —         | NAU1     | —   |
| NWA 2200              | Morocco                                  | 08/2004                    | 552                  | 1                   | Lunar              | 20.5                     | —               | —               | —          | —         | —        | See separate entry  |
| NWA 2210              | Er-Mor                                   | 12/2004                    | 74                   | 1                   | CH3                | 14.8                     | —               | —               | —          | —         | —        | See separate entry  |
| NWA 2225              | Er-Mor                                   | 04/2004                    | 40                   | 1                   | Ure                | 20                       | —               | —               | ~18        | —         | —        | See separate entry  |
| NWA 2439              | Münster                                  | 12/2002                    | 660                  | 1                   | H5                 | 22.0                     | S2              | W1              | 18.5       | 15.5      | IfP1     | —   |



Table 2. *Continued.* Meteorites from Northwest Africa.

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location | Comments   |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|----------|--|
| NWA 2440 | Münster                                     | 04/2004                       | 182                     | 13                  | H5                 | 20.0                        | S3              | W1              | 17           | 14.5         | IfP1     | —  |
| NWA 2441 | Munich                                      | 10/2002                       | 156                     | 1                   | L3–6               | 20.0                        | S2              | W2/3            | 27.5 ± 3.5   | 20.5 ± 2.5   | IfP1     | —  |
| NWA 2442 | Munich                                      | 10/2002                       | 192                     | 1                   | H5                 | 20.0                        | S4              | W2/3            | 18.5         | 15.5         | IfP1     | —  |
| NWA 2444 | Dortmund                                    | 11/2003                       | 74                      | 1                   | H6                 | 18.0                        | S4              | W1              | 17           | 15.5         | IfP1     | —  |
| NWA 2445 | Sainte-Marie                                | 06/2002                       | 42                      | 1                   | H3–5               | 10.0                        | S1              | W1              | 16.5 ± 7     | 15 ± 3       | IfP1     | —  |
| NWA 2450 | Stuttgart                                   | 04/2004                       | 766                     | 1                   | H4–6               | 26.0                        | S1              | W2              | 17           | 14 ± 1       | IfP1     | —  |
| NWA 2457 | Sainte Marie                                | 06/2004                       | 172                     | 1                   | Imc                | 22.0                        | S2              | W2              | 10.5 ± 3     | —            | IfP1     | —  |
| NWA 2458 | Sainte Marie                                | 06/2004                       | 284                     | 1                   | L3                 | 22.0                        | S4              | W1              | 15 ± 10      | 10 ± 7       | IfP1     | —  |
| NWA 2463 | Sainte Marie                                | 06/2004                       | 62                      | 1                   | H6                 | 11.0                        | S2              | W3              | 18           | 15.5         | IfP1     | —  |
| NWA 2464 | Sainte Marie                                | 06/2004                       | 126                     | 1                   | H6                 | 22.0                        | S2              | W3              | 18.5         | 16           | IfP1     | —  |
| NWA 2465 | Sainte Marie                                | 06/2004                       | 1.167                   | 15                  | H4                 | 26.0                        | S2              | W2              | 17           | 15 ± 1.5     | IfP1     | —  |
| NWA 2468 | Sainte Marie                                | 06/2004                       | 25                      | 1                   | H6                 | 5.0                         | S2              | W1              | 17           | 15           | IfP1     | —  |
| NWA 2469 | Sainte Marie                                | 06/2004                       | 348                     | 3                   | H5/6               | 22.0                        | S4              | W1              | 20.5         | 18           | IfP1     | —  |
| NWA 2470 | Sainte Marie                                | 06/2004                       | 266                     | 1                   | L3–4               | 20.0                        | S2              | W2              | 21 ± 1.3     | 14.5 ± 3     | IfP1     | —  |
| NWA 2471 | Sainte Marie                                | 06/2004                       | 374                     | 1                   | H6                 | 24.0                        | S2              | W3              | 18           | 14.5         | IfP1     | —  |
| NWA 2472 | Sainte Marie                                | 06/2004                       | 362                     | 1                   | H4                 | 30.0                        | S2              | W2              | 16           | 14.5         | IfP1     | —  |
| NWA 2473 | Sainte Marie                                | 06/2004                       | 60                      | 1                   | L4                 | 14.0                        | S2              | W4              | 22.5         | 18.5 ± 1.5   | IfP1     | —  |
| NWA 2478 | Sainte Marie                                | 06/2004                       | 473                     | 5                   | L6                 | 22.0                        | S4              | W5              | 24           | 20.5         | IfP1     | Ringwoodite  |
| NWA 2646 | Morocco                                     | 12/2004                       | 9.3                     | 1                   | Martian            | 3.4                         | —               | —               | —            | —            | IfP1     | See separate entry   |
| NWA 2656 | Unknown                                     | 2003                          | 386                     | 1                   | Acp                | 21                          | Low             | W3              | 8.0          | 8.4          | —        | See separate entry   |
| NWA 2681 | Er-Mor                                      | 2004                          | 37                      | 1                   | CO3.5              | 6.2                         | S2              | W2              | 38.5 ± 2.4   | —            | NAUI     | Plag (An <sub>86</sub> Or <sub>3</sub> ) + rare CAIs                   |
| NWA 2690 | Er-Mor                                      | 2004                          | 15,000                  | n                   | Euc                | 35                          | M-H             | W2              | —            | —            | NAUI     | Paired with NWA 1929   |
| NWA 2697 | Er-Mor                                      | 2004                          | 9424                    | 1                   | CV3                | 23                          | S1              | W2              | 38.1–53.6    | —            | NAUI     | Matrix olivine only  |
| NWA 2698 | Er-Mor                                      | 2004                          | 134                     | 1                   | How                | 20.3                        | S3–5            | W2              | —            | —            | NAUI     | Euc (Fs <sub>97.2-0</sub> Wo <sub>7-18</sub> )<br>Paired with NWA 1929 |
| NWA 2699 | Er-Mor                                      | 2005                          | 1294                    | n                   | Acp                | 22                          | Low             | W3              | 8.2 ± 0.2    | 9.1 ± 0.2    | NAUI     | Opx (Wo <sub>30</sub> ; An <sub>23.8</sub> )<br>Paired with NWA 2656   |
| NWA 2700 | Unknown                                     | 2004                          | 31.7                    | 1                   | Lunar              | 6.8                         | —               | —               | —            | —            | NAUI     | See separate entry   |
| NWA 2701 | Er-Mor                                      | 2004                          | 1168                    | 1                   | LL5                | 21.6                        | S2–4            | W2              | 28.7         | 23.5         | NAUI     | —  |
| NWA 2702 | Er-Mor                                      | 2005                          | 215                     | 1                   | R4                 | 20.3                        | S2              | W4              | 39.5 ± 2.4   | —            | NAUI     | Plag (An <sub>44</sub> Or <sub>4.8</sub> )<br>Chromite (Cr# = 89.7)    |
| NWA 2703 | Unknown                                     | 2004                          | 121                     | 1                   | Ure                | 20.5                        | Low             | Min             | 12.3         | 10.6         | —        | See separate entry   |
| NWA 2704 | Er-Mor                                      | 2005                          | 871                     | 1                   | L3.8               | 27.3                        | S2              | W2              | 21.3–29.5    | 18.2–24.2    | NAUI     | —  |
| NWA 2705 | Unknown                                     | 2005                          | 30                      | 1                   | Ure                | 16.8                        | —               | —               | 22.3         | —            | —        | See separate entry   |
| NWA 2706 | Er-Mor                                      | 2005                          | 1913                    | 1                   | H4                 | 26.2                        | S3–S6           | W2              | 17.5         | —            | NAUI     | —  |
| NWA 2707 | Er-Mor                                      | 2005                          | 577                     | 1                   | H5                 | 23.1                        | S2              | W3              | 18.7         | 16.8         | NAUI     | —  |
| NWA 2708 | Unknown                                     | 2004                          | 528                     | 1                   | CK4                | 20.3                        | M-H             | Mod             | —            | —            | —        | See separate entry   |
| NWA 2709 | Er-Mor                                      | 2005                          | 148                     | 1                   | L4                 | 21                          | S3              | W2              | 22.8         | 20           | NAUI     | —  |
| NWA 2710 | Er-Mor                                      | 2005                          | 216                     | 1                   | H5                 | 20                          | S2              | W1              | 18.8         | 16.3         | NAUI     | —  |
| NWA 2711 | Unknown                                     | 2004                          | 433                     | 1                   | Mes                | 24                          | —               | —               | —            | —            | —        | See separate entry   |
| NWA 2712 | Er-Mor                                      | 2005                          | 2500                    | 1                   | H5                 | 25.4                        | S2              | W2              | 18.5         | 16.6         | NAUI     | —  |
| NWA 2713 | Er-Mor                                      | 2004                          | 184                     | 1                   | H5                 | 21.1                        | S2              | W2              | 18.8         | 16.8         | NAUI     | —  |
| NWA 2714 | Er-Mor                                      | 2004                          | 1656                    | Several             | Acp                | 20.2                        | low             | W3/4            | 8.1          | 8.6          | NAUI     | Plag (A 22.2); Chromite,<br>Cr# = 0.85;<br>Paired with NWA 2656        |
| NWA 2717 | Er-Mor                                      | 2004                          | 70                      | 1                   | LL3.5              | 15.1                        | S2              | W2              | 25.8 ± 12.7  | —            | NAUI     | Olv (Cr <sub>2</sub> O <sub>3</sub> = 0.13 ± 0.08)                     |
| NWA 2718 | Er-Mor                                      | 2004                          | 254                     | 1                   | CO3.1              | 26.8                        | S2              | W4              | 20.8–53.6    | —            | NAUI     | Olv (Cr <sub>2</sub> O <sub>3</sub> = 0.26 ± 0.10)                     |

Table 2. *Continued. Meteorites from Northwest Africa.*

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location | Comments  |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|----------|---|
| NWA 2719 | Er-Mor                                      | 2004                          | 7581                    | 1                   | L4                 | 30                          | S5              | W2              | 23.8         | 20.4         | NAUI     | —   |
| NWA 2720 | Er-Mor                                      | 2004                          | 155                     | 1                   | L4                 | 20.5                        | S5              | W1              | 23.4         | 19.6         | NAUI     | —   |
| NWA 2721 | Er-Mor                                      | 2004                          | 46.5                    | 1                   | L3.8               | 10.3                        | S2              | W2              | 22.2 ± 5.6   | —            | NAUI     | —   |
| NWA 2722 | Er-Mor                                      | 2004                          | 707                     | 1                   | L4                 | 33                          | S4              | W2              | 22           | 19           | NAUI     | —   |
| NWA 2723 | Er-Mor                                      | 2004                          | 169                     | 1                   | L-Imc              | 23                          | S5-6            | W1              | 22.5–24.8    | —            | NAUI     | Quench olv = $\text{Fe}_{24.7-26.9}$  |
| NWA 2724 | Er-Mor                                      | 2004                          | 3804                    | 1                   | Euc                | 21                          | L-M             | Min             | —            | —            | —        | See separate entry  |
| NWA 2725 | Er-Mor                                      | 2004                          | 200                     | 1                   | L5                 | 46                          | S2              | W2              | 25.3         | 21.3         | NAUI     | —   |
| NWA 2726 | Er-Mor                                      | 2004                          | 86                      | 1                   | LL3.6              | 36                          | S2              | W2              | 26.8 ± 8.9   | —            | NAUI     | —   |
| NWA 2727 | Er-Mor                                      | 2005                          | 191.2                   | 4                   | Lunar              | 20.2                        | —               | —               | —            | —            | —        | See separate entry  |
| NWA 2728 | Er-Mor                                      | 2004                          | 338                     | 1                   | CO3.2              | 46                          | S2              | W2              | 42.2 ± 6.3   | —            | NAUI     | Olv ( $\text{Cr}_2\text{O}_3 = 0.12 \pm 0.09$ )   |
| NWA 2737 | Morocco                                     | 2000                          | 611                     | 9                   | Martian            | 20                          | —               | —               | —            | —            | —        | See separate entry  |
| NWA 2754 | Er-Mor                                      | 2004                          | 5500                    | 1                   | LL5                | 22.2                        | S4              | W2              | 29.6         | 24.7         | NAUI     | —   |
| NWA 2758 | Er-Mor                                      | 2004                          | 678                     | 4                   | Euc                | 37.5                        | L-H             | Min             | —            | —            | —        | See separate entry  |
| NWA 2760 | Er-Mor                                      | 2004                          | 607                     | 1                   | CO3.1              | 22                          | S1              | W2              | 1.2–49.4     | —            | NAUI     | Olv ( $\text{Cr}_2\text{O}_3 = 0.22 \pm 0.14$ )   |
| NWA 2761 | Er-Mor                                      | 2003                          | 234                     | 1                   | LL3.6              | 21.6                        | S2              | W2              | 13.8–29.0    | —            | NAUI     | —   |
| NWA 2762 | Er-Mor                                      | 2004                          | 156                     | 1                   | Ure                | 21.1                        | Low             | Min             | 15.2         | —            | NAUI     | Olivine/pigeonite = 19:1 (vol%)   |
| NWA 2763 | Er-Mor                                      | 2004                          | 69                      | 1                   | EL6                | 17.2                        | S2              | W1              | —            | —            | NAUI     | Olv rim ( $\text{Fa}_{12.4}$ )  |
| NWA 2765 | Er-Mor                                      | 2004                          | 68                      | 1                   | Euc                | 14                          | Low             | Min             | —            | —            | NAUI     | Pyx ( $\text{Fe}_{50.2}\text{Wo}_{13.3}$ ); An <sub>18</sub> ; CaS, MnS, FeS, daubreelite                       |
| NWA 2766 | Er-Mor                                      | 2004                          | 214                     | 1                   | LL3.9              | 23.9                        | S2              | W2              | 29.8         | 24           | NAUI     | Paired with NWA 1849  |
| NWA 2767 | Er-Mor                                      | 2004                          | 82.3                    | 1                   | LL6                | 16.8                        | S2              | W1              | 29.7         | 24.8         | NAUI     | —   |
| NWA 2768 | Er-Mor                                      | 2004                          | 2200                    | 1                   | LL4                | 31                          | S2              | W3              | 28.5         | 24.7         | NAUI     | —   |
| NWA 2769 | Er-Mor                                      | 2004                          | 126                     | 1                   | LL3.2              | 20.1                        | S2              | W2              | 1–37.2       | —            | NAUI     | Olv ( $\text{Cr}_2\text{O}_3 = 0.31 \pm 0.24$ )   |
| NWA 2770 | Er-Mor                                      | 2004                          | 58                      | 1                   | LL4                | 10.1                        | S3              | W3              | 29.1         | 24.4         | NAUI     | —   |
| NWA 2771 | Er-Mor                                      | 2004                          | 390                     | 1                   | H5                 | 44                          | S1              | W1              | 18.3         | 16.3         | NAUI     | —   |
| NWA 2772 | Er-Mor                                      | 2004                          | 35                      | 1                   | LL6                | 8                           | S4              | W3              | 30           | 24.7         | NAUI     | —   |
| NWA 2773 | Er-Mor                                      | 2004                          | 656                     | 1                   | L3.9               | 53                          | S2              | W3              | 23.0 ± 3.2   | —            | NAUI     | —   |
| NWA 2774 | Er-Mor                                      | 2004                          | 398                     | 1                   | H4                 | 20.4                        | S2              | W1              | 18.8         | 16.3         | NAUI     | —   |
| NWA 2776 | Er-Mor                                      | 2004                          | 289                     | 1                   | H5                 | 20.3                        | S2              | W3              | 18.4         | 16           | NAUI     | —   |
| NWA 2777 | Er-Mor                                      | 2004                          | 1987                    | 2                   | H5                 | 21.6                        | S2              | W3              | 19           | 17.2         | NAUI     | —   |
| NWA 2778 | Er-Mor                                      | 2004                          | 442                     | 1                   | H4                 | 23                          | S2              | W4              | 17.5         | 16.2         | NAUI     | —   |
| NWA 2779 | Er-Mor                                      | 2004                          | 1537                    | 1                   | L5                 | 21.1                        | S3              | W3              | 24.3         | 20.6         | NAUI     | —   |
| NWA 2781 | Er-Mor                                      | 2004                          | 528                     | 2                   | L4–5               | 20.5                        | S4              | W3              | 24.5         | 20.3         | NAUI     | —   |
| NWA 2782 | Er-Mor                                      | 2005                          | 330                     | 1                   | Mes                | 23.5                        | Low             | W2              | —            | —            | NAUI     | Opx ( $\text{Fe}_{52.5}\text{Wo}_{3.1}$ )<br>Pyx ( $\text{FeO/MnO} = 27.5–30$ )                                 |
| NWA 2783 | Er-Mor                                      | 2005                          | 176                     | 1                   | H4                 | 22                          | S3              | W2              | 18.7         | 15.8         | NAUI     | —   |
| NWA 2784 | Unknown                                     | 2005                          | 141                     | 1                   | Euc                | 27                          | Low             | W1              | —            | —            | NAUI     | See separate entry  |
| NWA 2785 | Er-Mor                                      | 2005                          | 264                     | 1                   | L3.5               | 25.2                        | S2              | W2              | 13–28        | —            | NAUI     | Olv ( $\text{Cr}_2\text{O}_3 = 0.05–0.15$ )   |
| NWA 2786 | Er-Mor                                      | 2005                          | 30                      | 1                   | Mes                | 6                           | Mod             | W1              | —            | —            | NAUI     | Pyx ( $\text{Fe}_{58.1}\text{Wo}_{3.0}$ ;<br>$\text{FeO/MnO} = 27$ );<br>An <sub>93</sub> ; kamacite (Ni = 5.4) |
| NWA 2787 | Er-Mor                                      | 2005                          | 32                      | 1                   | CK4                | 6.5                         | S2              | W3              | 30.5         | —            | NAUI     | Mag ( $\text{Cr}_2\text{O}_3 = 3.9$ ,<br>$\text{Al}_2\text{O}_3 = 1.2$ )  |
| NWA 2792 | Er-Mor                                      | 2005                          | 18.6                    | 1                   | Ure                | 4.4                         | Mod             | W2              | 20.3 (core)  | —            | NAUI     | Olv ( $\text{FeO/MnO} = 46$ )<br>Pyx ( $\text{Fe}_{517.8}\text{Wo}_{7.5}$ ,<br>$\text{FeO/MnO} = 26$ )          |

Table 2. *Continued.* Meteorites from Northwest Africa.

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location | Comments  |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|----------|---|
| NWA 2793 | Er-Mor                                      | 2005                          | 768                     | 1                   | L4                 | 24.5                        | S4              | W2              | 24           | 19.9         | NAUI     | —   |
| NWA 2794 | Er-Mor                                      | 2005                          | 145                     | 1                   | How                | 21.4                        | L-M             | Mod             | —            | —            | —        | See separate entry  |
| NWA 2795 | Er-Mor                                      | 2005                          | 329                     | 1                   | Dio                | 20.4                        | Low             | Mod             | —            | —            | —        | See separate entry  |
| NWA 2796 | Er-Mor                                      | 2005                          | 34                      | 1                   | LL3.1              | 6.4                         | S2              | W3              | 3.2–56.1     | —            | NAUI     | Olv (Cr <sub>2</sub> O <sub>3</sub> = 0.44 ± 0.19,<br>FeO/MnO = 38–63)                        |
| NWA 2797 | Er-Mor                                      | 2005                          | 1211                    | 1                   | L3.8               | 21.9                        | S2              | W2              | 19.2–24.2    | —            | NAUI     | (FeO/MnO)<br>in olivine = 34–50   |
| NWA 2798 | Er-Mor                                      | 2005                          | 1805                    | 3                   | L3.2               | 23.6                        | S2              | W1              | 5–39         | —            | NAUI     | Olv-Cr <sub>2</sub> O <sub>3</sub> = 0.21 ± 0.09  |
| NWA 2799 | Er-Mor                                      | 2005                          | 1311                    | 1                   | LL6/7              | 21.3                        | S1              | W2              | 29.7 ± 0.02  | 25.1 ± 0.02  | NAUI     | Plag (An <sub>12</sub> Or <sub>5.1</sub> )<br>1 chondrule fragment<br>per 1.4 cm <sup>2</sup> |
| NWA 2801 | Er-Mor                                      | 2005                          | 7700                    | 1                   | Dio                | 20                          | Low             | W3              | 27.4         | —            | NAUI     | Olv (FeO/MnO = 27);<br>Fs <sub>22.6</sub> Wo <sub>1.8</sub> ;<br>Paired with NWA 1877         |
| NWA 2802 | Er-Mor                                      | 2005                          | 30                      | 2                   | CV3                | 6                           | S2              | W3              | 36.4–52.1    | —            | NAUI     | —   |
| NWA 2803 | Er-Mor                                      | 2005                          | 4500                    | 1                   | L5                 | 45                          | S2              | W2              | 24.6         | 23.8         | NAUI     | —   |
| NWA 2822 | Er-Mor                                      | 2005                          | 1675                    | 3                   | R4                 | 26.2                        | S2              | W2–4            | 39.5 ± 0.4   | —            | NAUI     | Olv (FeO/MnO = 83);<br>Pyx (Fs <sub>12.1</sub> Wo <sub>45.8</sub> );<br>Polymict              |
| NWA 2823 | Er-Mor                                      | 2005                          | 144                     | 1                   | LL6                | 20.8                        | S3              | W1              | 28.8         | 24.1         | NAUI     | —   |
| NWA 2825 | Er-Mor                                      | 2004                          | 664                     | 1                   | Pac                | 21.1                        | low             | W3              | 22           | 18.9         | NAUI     | Olv (FeO/MnO = 63.5)<br>Paired with NWA 3133  |
| NWA 2826 | Er-Mor                                      | 2004                          | 685                     | 1                   | LL5                | 22                          | S2              | W1              | 28.2         | 22.1         | NAUI     | Paired with NWA 2053  |
| NWA 2827 | Er-Mor                                      | 2004                          | 1000                    | 1                   | LL4                | 21.3                        | S2              | W2              | 27.8         | 22.4         | NAUI     | —   |
| NWA 2829 | Er-Mor                                      | 2004                          | 687                     | 1                   | Dio                | 21.7                        | Mod             | W2              | 28           | 23.1         | NAUI     | Olv (FeO/MnO = 46)<br>Paired with NWA 1877  |
| NWA 2830 | Er-Mor                                      | 2005                          | 1130                    | 1                   | H4                 | 23.3                        | S2              | W3              | 19.1         | 17           | NAUI     | —   |
| NWA 2831 | Er-Mor                                      | 2005                          | 429                     | 1                   | L4                 | 20                          | S2              | W4              | 27.9         | 23           | NAUI     | —   |
| NWA 2832 | Er-Mor                                      | 2005                          | 359                     | 1                   | LL4                | 21.5                        | S3              | W3              | 30.7         | 24.4         | NAUI     | —   |
| NWA 2833 | Er-Mor                                      | 2005                          | 2112                    | 1                   | L3.8               | 21.6                        | S2              | W3              | 23.4 ± 5.3   | 20 ± 3.3     | NAUI     | —   |
| NWA 2834 | Er-Mor                                      | 2005                          | 792                     | 1                   | L5                 | 23.5                        | S2              | W3              | 24.3         | 20.3         | NAUI     | —   |
| NWA 2835 | Er-Mor                                      | 2005                          | 1233                    | 1                   | H7                 | 23.3                        | S2              | W4              | 17.8         | 16.4         | NAUI     | Plag (An <sub>80</sub> )<br>Paired with NWA 2353  |
| NWA 2836 | Er-Mor                                      | 2005                          | 1300                    | 1                   | LL3.7              | 30                          | S2              | W3              | 29.7 ± 5.3   | —            | NAUI     | —   |
| NWA 2837 | Er-Mor                                      | 2005                          | 41                      | 1                   | CV3                | 7.4                         | S2              | W0/1            | 36–54        | —            | NAUI     | —   |
| NWA 2838 | Er-Mor                                      | 2005                          | 253                     | 1                   | L4                 | 28                          | S2              | W2              | 22           | 19.8         | NAUI     | —   |
| NWA 2839 | Er-Mor                                      | 2004                          | 144                     | 1                   | L4                 | 37                          | S4              | W3              | 24.7         | 20.8         | NAUI     | —   |
| NWA 2840 | Er-Mor                                      | 2004                          | 129                     | 1                   | L4                 | 25                          | S3–5            | W1              | 23.6         | 20.4         | NAUI     | —   |
| NWA 2841 | Er-Mor                                      | 2004                          | 190                     | 1                   | H3.9               | 25                          | S2              | W1              | 18.0 ± 1.4   | 16.2 ± 0.8   | NAUI     | —   |
| NWA 2842 | Er-Mor                                      | 2004                          | 423                     | 1                   | L4                 | 31                          | S2              | W2              | 24.2         | 20.6         | NAUI     | —   |
| NWA 2843 | Er-Mor                                      | 2004                          | 404                     | 1                   | H3.9               | 57                          | S2              | W0/1            | 17.5 ± 2.0   | 15.6 ± 1.6   | NAUI     | —   |
| NWA 2844 | Er-Mor                                      | 2004                          | 837                     | 1                   | L5                 | 52                          | S2              | W3              | 24.4         | 20.6         | NAUI     | —   |
| NWA 2845 | Er-Mor                                      | 2004                          | 546                     | 1                   | H4                 | 37                          | S3              | W2              | 17.5         | 16           | NAUI     | —   |
| NWA 2846 | Er-Mor                                      | 2004                          | 352                     | 1                   | L4                 | 52                          | S4–6            | W5              | 24.1         | 20.7         | NAUI     | Ringwoodite in melt veins   |
| NWA 2847 | Er-Mor                                      | 2004                          | 283                     | 1                   | H4                 | 51                          | S2              | W2              | 17.8         | 16.2         | NAUI     | —   |
| NWA 2848 | Er-Mor                                      | 2004                          | 538                     | 1                   | L4                 | 35                          | S4              | W2              | 23.8         | 20.5         | NAUI     | Maskelynite present   |
| NWA 2849 | Er-Mor                                      | 2004                          | 846                     | 1                   | L4                 | 37                          | S2              | W3              | 23.9         | 20.5         | NAUI     | —   |
| NWA 2850 | Er-Mor                                      | 2004                          | 561                     | 1                   | L5                 | 29                          | S3              | W3              | 24.7         | 20.8         | NAUI     | —   |
| NWA 2851 | Er-Mor                                      | 2004                          | 417                     | 1                   | L4                 | 22                          | S2              | W3              | 24.1         | 20.7         | NAUI     | —   |

Table 2. *Continued. Meteorites from Northwest Africa.*

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location | Comments   |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|----------|--|
| NWA 2852 | Unknown                                     | 2004                          | 398                     | 1                   | Pac                | 21.1                        | Low             | W3              | 22.2         | 18.7         | NAUI     | Olv = (FeO/MnO = 63.8)<br>Paired with NWA 3133   |
| NWA 2853 | Unknown                                     | 2004                          | 1000                    | 1                   | How                | 22                          | High            | –               | –            | –            | NAUI     | See separate entry   |
| NWA 2854 | Er-Mor                                      | 2004                          | 167                     | 1                   | CK3/4              | 20.3                        | S2              | W2              | 33.5         | –            | NAUI     | Olv (FeO/MnO = 101);<br>Mag (Cr <sub>2</sub> O <sub>3</sub> = 4.8,<br>Al <sub>2</sub> O <sub>3</sub> = 1.86)   |
| NWA 2856 | Er-Mor                                      | 2004                          | 1609                    | 1                   | H4                 | 41                          | S2              | W3              | 18.8         | 16.4         | NAUI     | –  |
| NWA 2857 | Er-Mor                                      | 2004                          | 5866                    | 1                   | L4                 | 40                          | S2              | W2              | 23.3         | 20.7         | NAUI     | –  |
| NWA 2858 | Er-Mor                                      | 2004                          | 2335                    | 1                   | L4                 | 40                          | S3              | W1              | 24.7         | 20.4         | NAUI     | –  |
| NWA 2859 | Er-Mor                                      | 2004                          | 920                     | 1                   | H4                 | 32                          | S2              | W2              | 18.1         | 16.1         | NAUI     | –  |
| NWA 2860 | Er-Mor                                      | 2004                          | 878                     | 1                   | H4                 | 23                          | S2              | W1              | 18.6         | 16.5         | NAUI     | –  |
| NWA 2861 | Er-Mor                                      | 2004                          | 2029                    | 1                   | L4                 | 63                          | S3              | W3              | 25.2         | 21.7         | NAUI     | –  |
| NWA 2862 | Er-Mor                                      | 2004                          | 486                     | 1                   | H4                 | 50                          | S2              | W3              | 18.5         | 16.3         | NAUI     | –  |
| NWA 2863 | Er-Mor                                      | 2004                          | 480                     | 1                   | L4                 | 41                          | S2              | W3              | 24.9         | 20.6         | NAUI     | –  |
| NWA 2864 | Er-Mor                                      | 2004                          | 90                      | 1                   | LL4                | 19                          | S3              | W2              | 28.9         | 23.8         | NAUI     | –  |
| NWA 2865 | Er-Mor                                      | 2004                          | 1503                    | 1                   | L4                 | 25                          | S2              | W2              | 24           | 20.9         | NAUI     | –  |
| NWA 2866 | Er-Mor                                      | 2005                          | 213                     | 1                   | Acp                | 21                          | Low             | W3              | 8.2 ± 0.2    | 9.2 ± 0.2    | NAUI     | Plag (An <sub>24.5</sub> Or <sub>2.8</sub> );<br>Cpx (Fs <sub>84.2</sub> Wo <sub>4.3</sub> );<br>Chromite (Cr# = 70);<br>Paired with NWA 2656<br>(FeO/MnO) in olivine =<br>133 |
| NWA 2867 | Er-Mor                                      | 2002                          | 60                      | 1                   | CK4                | 20                          | S1              | W0/1            | 30.5 ± 2.3   | –            | NAUI     | Magnetite (Cr <sub>2</sub> O <sub>3</sub> = 3.8,<br>Al <sub>2</sub> O <sub>3</sub> = 0.88 wt%);<br>Plag (An <sub>24.3</sub> );<br>Paired with NWA 2656<br>Chromite (Cr# = 71)  |
| NWA 2871 | Er-Mor                                      | 2005                          | 3467                    | 1                   | Acp                | 24.1                        | Low             | W3              | 8.4          | 9.1          | NAUI     | See separate entry   |
| NWA 2890 | Unknown                                     | 2004                          | 132                     | 1                   | How                | 22.1                        | –               | –               | –            | –            | –        | See separate entry   |
| NWA 2899 | Unknown                                     | 2004                          | 11                      | 1                   | Oc                 | 2.5                         | –               | –               | –            | –            | –        | See separate entry   |
| NWA 2900 | Unknown                                     | 2004                          | 1375                    | 1                   | CV3                | 20.5                        | –               | –               | –            | –            | –        | See separate entry   |
| NWA 2901 | Unknown                                     | 2004                          | 308                     | 1                   | CV3                | 21.1                        | –               | –               | –            | –            | –        | See separate entry   |
| NWA 2902 | Unknown                                     | 2003                          | 1000                    | n                   | Oc-Imc             | 22.7                        | –               | –               | 12.9–23.9    | 11.2–19.7    | –        | See separate entry   |
| NWA 2904 | Unknown                                     | 2003                          | 29.04                   | 4                   | Euc                | 9.95                        | –               | –               | –            | –            | –        | See separate entry   |
| NWA 2905 | Er-Mor                                      | 2004                          | 205                     | 1                   | L4                 | 21                          | S2              | W2              | 24.3         | 21.1         | NAUI     | –  |
| NWA 2906 | Er-Mor                                      | 2004                          | 215                     | 1                   | L4                 | 26.8                        | S2              | W2              | 23.7         | 20.6         | NAUI     | –  |
| NWA 2908 | Er-Mor                                      | 2003                          | 148                     | 1                   | LL4                | 26.2                        | S2              | W1              | 28.7         | 24           | NAUI     | –  |
| NWA 2910 | Er-Mor                                      | 2005                          | 41                      | 1                   | Euc                | 8.2                         | M-H             | W2              | –            | –            | NAUI     | Py (Fs <sub>58.5</sub> Wo <sub>3.9</sub> );<br>Lamellae (Fs <sub>82.1</sub> Wo <sub>26.6</sub> );<br>Plag (An <sub>89</sub> )  |
| NWA 2911 | Er-Mor                                      | 2005                          | 113                     | 1                   | L3.5               | 20                          | S2              | W2              | 23.1 ± 0.08  | –            | NAUI     | Olv (Cr <sub>2</sub> O <sub>3</sub> = 0.13 ± 0.08)   |
| NWA 2912 | Unknown                                     | 2005                          | 203                     | 1                   | How                | 22.1                        | Low             | Min             | –            | –            | NAUI     | Dio (Fs <sub>59.2</sub> Wo <sub>4.2</sub> );<br>Euc (Fs <sub>89</sub> Wo <sub>2.7</sub> )  |
| NWA 2913 | Unknown                                     | 2005                          | 100                     | 3                   | Euc                | 23.1                        | –               | Min             | –            | –            | –        | See separate entry   |
| NWA 2914 | Unknown                                     | 2005                          | 399                     | 1                   | Euc                | 21.1                        | –               | Min             | –            | –            | –        | See separate entry   |
| NWA 2917 | Morocco                                     | 2005                          | 256                     | 1                   | LL4                | 20.8                        | S3              | W2              | 29.3         | 23.8         | NAUI     | –  |
| NWA 2918 | Morocco                                     | 2005                          | 237                     | 1                   | CO3.0              | 20                          | S1              | W2              | 1.2–64.5     | –            | NAUI     | Olv (Cr <sub>2</sub> O <sub>3</sub> = 0.38 ± 0.24)   |
| NWA 2919 | Morocco                                     | 2005                          | 1214                    | 3                   | H4                 | 20.6                        | S2              | W4              | 17.8         | 15.7         | NAUI     | –  |
| NWA 2920 | Morocco                                     | 2005                          | 734                     | 1                   | LL3.5              | 20.4                        | S2              | W4              | 29.7 ± 5.1   | 23.2 ± 3.2   | NAUI     | –  |
| NWA 2921 | Morocco                                     | 2005                          | 195                     | 1                   | CK3.8              | 20.5                        | S1              | W3              | 39.5 ± 1.3   | –            | NAUI     | Olv (FeO/MnO = 72–76)  |

Table 2. *Continued.* Meteorites from Northwest Africa.

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location | Comments  |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|----------|---|
| NWA 2922 | Morocco                                     | 2005                          | 186                     | 1                   | L5                 | 20.3                        | S3              | W3              | 24.4         | 20.2         | NAUI     | –   |
| NWA 2923 | Unknown                                     | 2005                          | 16,000                  | n                   | Dio                | 50                          | –               | –               | –            | –            | –        | See separate entry  |
| NWA 2924 | Unknown                                     | 2005                          | 606                     | 1                   | Mes                | 85                          | –               | –               | –            | –            | –        | See separate entry  |
| NWA 2927 | Er-Mor                                      | 2005                          | 171                     | 1                   | H4                 | 20.1                        | S2              | W2              | 18.8         | 16.7         | NAUI     | –   |
| NWA 2932 | Er-Mor                                      | 2005                          | 206                     | 1                   | Mes                | 23.2                        | S2              | W2              | 29.4         | 31.3         | NAUI     | Large metal nuggets;<br>Paired with NWA 2923  |
| NWA 2933 | Er-Mor                                      | 2005                          | 511                     | 1                   | L3.3               | 21.3                        | S2              | W2              | 25.2 ± 15    | –            | NAUI     | Olv (Cr <sub>2</sub> O <sub>3</sub> = 0.16 ± 0.07)<br>Paired with NWA 2685  |
| NWA 2935 | Er-Mor                                      | 2005                          | 460                     | 1                   | LL6/7              | 29                          | S3              | W2              | 31.7         | 25.7         | NAUI     | –   |
| NWA 2936 | Er-Mor                                      | 2005                          | 290                     | 1                   | H4                 | 24                          | S2              | W2              | 16.8         | 15.2         | NAUI     | –   |
| NWA 2937 | Er-Mor                                      | 2005                          | 351                     | 1                   | H5                 | 25                          | S2              | W3              | 18.8         | 17           | NAUI     | –   |
| NWA 2938 | Er-Mor                                      | 2003                          | 7000                    | 1                   | H4                 | 24.7                        | S1              | W4              | 17.7         | 15.9         | NAUI     | –   |
| NWA 2939 | Er-Mor                                      | 2003                          | 107                     | 1                   | R4                 | 20.2                        | S2              | W4              | 38.9 ± 0.05  | –            | NAUI     | Olv (FeO/MnO = 81–91);<br>Plag (An <sub>13</sub> )  |
| NWA 2940 | Er-Mor                                      | 2003                          | 138                     | 1                   | R3.9               | 20.4                        | S2              | W3              | 38.7 ± 2.1   | –            | NAUI     | Olv (FeO/MnO = 74–81);<br>Plag (An <sub>11.7</sub> )  |
| NWA 2941 | Er-Mor                                      | 2003                          | 115                     | 1                   | R3.8               | 20.4                        | S2              | W2              | 39.1 ± 3.4   | –            | NAUI     | Olv (FeO/MnO = 75–86);<br>Plag (An <sub>8.4–18.6</sub> )  |
| NWA 2942 | Unknown                                     | 2003                          | 345                     | 1                   | Euc                | 20.6                        | Mod             | W2              | –            | –            | NAUI     | Pyx (Fs <sub>82.1</sub> Wo <sub>2.3</sub> ,<br>Fs <sub>52</sub> Wo <sub>11.4</sub> );<br>Plag (An <sub>90.7</sub> );<br>Chromite (Cr# = 83) |
| NWA 2943 | Er-Mor                                      | 2005                          | 300                     | 1                   | R3–6               | 22.8                        | S1              | W3              | 28.2 ± 13    | 22.1 ± 7.2   | NAUI     | –   |
| NWA 2975 | Algeria                                     | 2005                          | 70.1                    | 1                   | Martian            | 20.2                        | –               | –               | –            | –            | –        | See separate entry  |
| NWA 2977 | Tag-Mor                                     | 11/2005                       | 233                     | 1                   | Lunar              | 20.1                        | –               | Min             | –            | –            | –        | See separate entry<br>Paired with NWA 773 and<br>NWA 2700   |
| NWA 2995 | Algeria                                     | 2005                          | 538                     | 1                   | Lunar              | 21.2                        | –               | –               | –            | –            | –        | See separate entry  |
| NWA 2999 | Tag-Mor                                     | 08/2004                       | 392                     | 12                  | Ang                | 22.0                        | –               | –               | –            | –            | –        | See separate entry  |
| NWA 3151 | Tag-Mor                                     | 04/2005                       | 1500                    | 1                   | Bra                | 20.0                        | –               | –               | –            | –            | –        | See separate entry  |
| NWA 3160 | Er-Mor                                      | 07/2005                       | 34                      | 3                   | Lunar              | 4.8                         | –               | –               | –            | –            | –        | See separate entry<br>Paired with NWA 2727  |
| NWA 3163 | Ouarzazate                                  | 08/2005                       | 1634                    | 1                   | Lunar              | 20.1                        | –               | –               | –            | –            | –        | See separate entry  |
| NWA 3164 | Rab-Mor                                     | 08/2004                       | 928                     | n                   | Ang                | 21.0                        | –               | –               | –            | –            | –        | See separate entry<br>Paired with NWA 2999  |
| NWA 3169 | Er-Mor                                      | 2003                          | 4500                    | 1                   | LL5                | 62.4                        | S2-5            | W3              | 27.6         | 23           | NAUI     | –   |
| NWA 3172 | Er-Mor                                      | 2003                          | 187                     | 1                   | H4                 | 24.7                        | S4              | W1              | 19.4         | 17.2         | NAUI     | –   |
| NWA 3342 | Morocco                                     | 1998                          | 1250                    | 1                   | H/L4               | 31.4                        | S2              | W1              | 17–28        | 11.2–19.9    | Shim1    | –   |
| NWA 3343 | Morocco                                     | 1998                          | 760                     | 1                   | L4                 | 23.2                        | S2              | W1              | 22–27        | 21           | Shim2    | –   |
| NWA 3347 | Dortmund                                    | 10/2000                       | 4480                    | 1                   | L3–5               | 26.0                        | S4              | W2-3            | 24 ± 1.2     | 20.5 ± 1     | Ifp2     | –   |
| NWA 3348 | Dortmund                                    | 10/2000                       | 654                     | 1                   | L5                 | 23.0                        | S3              | W2              | 22           | 20           | Ifp2     | –   |
| NWA 3349 | Hamburg                                     | 11/2000                       | 297                     | 1                   | L6                 | 22.0                        | S3              | W1-2            | 24.5         | 20.5         | Ifp2     | –   |
| NWA 3350 | Hamburg                                     | 11/2000                       | 110                     | 1                   | Ure                | 20.0                        | S2              | –               | 21           | 18.5         | Ifp2     | –   |
| NWA 3351 | Hamburg                                     | 11/2000                       | 132                     | 1                   | LL4–6              | 22.0                        | S2              | W3              | 28           | 22.5         | Ifp2     | –   |
| NWA 3352 | Hamburg                                     | 11/2000                       | 7430                    | 1                   | H6                 | 22.0                        | S4              | W3              | 17.5         | 15.5         | Ifp2     | –   |
| NWA 3353 | Stuttgart                                   | 02/2001                       | 3220                    | 1                   | L6                 | 20.0                        | S4              | W3              | 24.5         | 21           | Ifp2     | –   |
| NWA 3354 | Stuttgart                                   | 02/2001                       | 723                     | 1                   | H6                 | 21.0                        | S3              | W1-2            | 18           | 16.5         | Ifp2     | Ringwoodite   |
| NWA 3355 | Stuttgart                                   | 02/2001                       | 617                     | 1                   | L6                 | 19.0                        | S3              | W2              | 24           | 20           | Ifp2     | –   |
| NWA 3356 | Stuttgart                                   | 02/2001                       | 2075                    | 3                   | L6                 | 21.0                        | S4              | W2              | 24           | 20           | Ifp2     | –   |

Table 2. *Continued. Meteorites from Northwest Africa.*

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location | Comments           |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|----------|--------------------|
| NWA 3357 | Sainte Marie                                | 05/2000                       | 1030                    | 1                   | L4-6               | 30.0                        | S2              | W1              | 22.5         | 19.5         | Ifp2     | —                  |
| NWA 3358 | Sainte Marie                                | 05/2001                       | 1162                    | 1                   | H(L)3              | 23.0                        | —               | W2              | 13.5 ± 7     | 11 ± 10      | Ifp2     | —                  |
| NWA 3359 | Sainte Marie                                | 06/2005                       | 509                     | 1                   | Euc                | 20.0                        | S2              | W1              | —            | 55.5 ± 2     | Ifp2     | —                  |
| NWA 3360 | Dortmund                                    | 10/2000                       | 112                     | 1                   | H3-5               | 20.0                        | S3              | W2              | 19 ± 2.4     | 16 ± 2.1     | Ifp2     | —                  |
| NWA 3361 | Dortmund                                    | 10/2000                       | 29                      | 1                   | L3-6               | 8.5                         | S4              | W1-2            | 24 ± 2.7     | 19 ± 1.8     | Ifp2     | —                  |
| NWA 3362 | Dortmund                                    | 10/2000                       | 741                     | 1                   | L4-6               | 30.0                        | S4              | W1              | 23.5         | 19           | Ifp2     | —                  |
| NWA 3363 | Dortmund                                    | 10/2000                       | 801                     | 1                   | L4-6               | 27.0                        | S4              | W1              | 23           | 19           | Ifp2     | —                  |
| NWA 3364 | Sainte Marie                                | 2004                          | 538                     | 1                   | R3-5               | 41.0                        | S2              | W3              | 39.5 ± 2.5   | —            | Ifp2     | —                  |
| NWA 3365 | Sainte Marie                                | 2004                          | 84                      | 1                   | LL4-6              | 13.0++                      | S2              | W1              | 29           | 23.5         | Ifp2     | —                  |
| NWA 3366 | Unknown                                     | 2004                          | 245                     | 1                   | H6                 | 22.0                        | S2              | W2              | 18           | 14           | Ifp2     | —                  |
| NWA 3367 | Münster                                     | 2005                          | 214                     | 1                   | L3-6               | 20.0                        | S2              | W1              | 24.2 ± 1.4   | 17.6 ± 1.8   | Ifp2     | —                  |
| NWA 3368 | Unknown                                     | 2005                          | 1600                    | 1                   | Euc                | 23.0                        | —               | —               | —            | —            | —        | See separate entry |
| NWA 4000 | Morocco                                     | 23/10/2002                    | 1465                    | 1                   | L4                 | 89.5                        | S3              | W2              | 24-26        | 22-24        | NUM1     | —                  |
| NWA 4001 | Morocco                                     | 23/10/2002                    | 1113                    | 1                   | L4                 | 24.8                        | S3              | W3              | 23-27        | 21-26        | NUM1     | —                  |
| NWA 4002 | Morocco                                     | 2004                          | 93.8                    | 1                   | L3-6               | 92.7                        | S3-S6           | W1              | 24.3-24.7    | 12.5-21.2    | CML1-1   | —                  |
| NWA 4003 | Morocco                                     | 02/2002                       | 357                     | 1                   | H5                 | 353.5                       | S3              | W1              | 18.8 ± 0.5   | 17.2 ± 1.2   | CML1-1   | —                  |
| NWA 4004 | Morocco                                     | 2005                          | 76.1                    | 1                   | H5                 | 17.3                        | S2              | W2              | 17.2         | 15.3         | MNB1-1   | —                  |
| NWA 4005 | Morocco                                     | 2005                          | 470.9                   | 1                   | H5/6               | 25.2                        | S2              | W3              | 18.3         | 16.2         | MNB1-1   | —                  |
| NWA 4006 | Morocco                                     | 2004                          | 758.1                   | 3                   | L4 Imc             | 23.6                        | —               | W0/1            | 22.6         | 9.8-20.3     | MNB1-1   | —                  |
| NWA 4007 | Morocco                                     | 2005                          | 511                     | 1                   | H6                 | 61.6                        | S3              | W2              | 17.6         | 15.6         | MNB1-1   | —                  |
| NWA 4008 | Morocco                                     | 2005                          | 195                     | 1                   | L6                 | 21.1                        | S4              | W2              | 24.1         | 19.9         | MNB1-1   | —                  |
| NWA 4009 | Morocco                                     | 2005                          | 856                     | 1                   | H5/6               | 21.1                        | S2              | W2              | 18.2         | 16.1         | MNB1-1   | —                  |
| NWA 4010 | Morocco                                     | 2005                          | 1676                    | 1                   | L4                 | 29.6                        | S3              | W2              | 23.9         | 5.2-21.1     | MNB1-1   | —                  |
| NWA 4011 | Morocco                                     | 2004                          | 77.7                    | 1                   | H3                 | 17.3                        | S3              | W1              | 0.7-25.3     | 8.1-20.8     | MNB1-1   | —                  |
| NWA 4012 | Morocco                                     | 2005                          | 149.4                   | 1                   | L4                 | 24.4                        | S2              | W2              | 26.6         | 10.9-24.7    | MNB1-1   | —                  |
| NWA 4013 | Morocco                                     | 2004                          | 20.7                    | 1                   | H6                 | 4.5                         | S2              | W3              | 18.4         | 16.5         | MNB1-1   | —                  |
| NWA 4014 | Morocco                                     | 2004                          | 35.7                    | 1                   | L3                 | 7.8                         | S2              | W1              | 0.6-28.7     | 1.7-28.6     | MNB1-1   | —                  |
| NWA 4015 | Morocco                                     | 2004                          | 3.5                     | 1                   | H3                 | 0.7                         | S2              | W2              | 0.9-43.8     | 2.7-17.4     | MNB1-1   | —                  |
| NWA 4016 | Morocco                                     | 2005                          | 81.6                    | 1                   | L3                 | 18.0                        | S3/4            | W2              | 0.3-22.7     | 4.6-37.3     | MNB1-1   | —                  |
| NWA 4017 | Morocco                                     | 2005                          | 216.9                   | 1                   | Mes                | 23.4                        | Low             | M-M             | —            | —            | —        | See separate entry |
| NWA 4018 | Morocco                                     | 2005                          | 158.6                   | 1                   | Euc                | 23.1                        | Mod             | Mod             | —            | —            | —        | See separate entry |
| NWA 4019 | Morocco                                     | 2005                          | 504.7                   | 1                   | Euc                | 22.0                        | Low             | Min             | —            | —            | —        | See separate entry |
| NWA 4020 | Morocco                                     | 2005                          | 209.4                   | 10                  | L/LL3              | 21.9                        | S3/4            | W3              | 1.4-27.3     | 4.0-23.1     | MNB1-1   | —                  |
| NWA 4021 | Morocco                                     | 2005                          | 1212.5                  | 5                   | LL6                | 22.8                        | S2/3            | W0/1            | 27.7         | 22.9         | MNB1-1   | —                  |
| NWA 4022 | Morocco                                     | 2005                          | 886.9                   | 1                   | L3                 | 22.6                        | S3/4            | W2              | 1.2-41.7     | 5.3-30.3     | MNB1-1   | —                  |
| NWA 4023 | Morocco                                     | 2005                          | 17.2                    | 1                   | Euc                | 3.5                         | High            | Mod             | —            | —            | —        | See separate entry |
| NWA 4027 | Morocco                                     | 2005                          | 1024                    | 1                   | H6                 | 25.0                        | S2              | W2/3            | 18.6         | 16.6         | MNB1-1   | —                  |
| NWA 4028 | Morocco                                     | 2004                          | 810                     | 1                   | L6                 | 25.9                        | S4              | W2/3            | 24.5         | 21           | MNB1-1   | —                  |
| NWA 4029 | Morocco                                     | 2004                          | 2072                    | 1                   | L6                 | 24.6                        | S4              | W2/3            | 24.3         | 20.6         | MNB1-1   | —                  |
| NWA 4030 | Morocco                                     | 2004                          | 87                      | 1                   | L6                 | 20.5                        | S4              | W1              | 24.7         | 20.2         | MNB1-1   | —                  |
| NWA 4031 | Morocco                                     | 2004                          | 390                     | 1                   | LL6                | 21.1                        | S4              | W0/1            | 28.4         | 22.7         | MNB1-1   | —                  |
| NWA 4032 | Morocco                                     | 2004                          | 10.5                    | 1                   | Euc                | 2.2                         | High            | Min             | —            | —            | —        | See separate entry |
| NWA 4033 | Morocco                                     | 2004                          | 1042                    | 1                   | L/LL5              | 25.7                        | S3              | W1              | 26.6         | 21.3         | MNB1-1   | —                  |
| NWA 4034 | Morocco                                     | 2005                          | 1513                    | 1                   | Dio                | 24.5                        | Mod             | High            | —            | —            | —        | See separate entry |
| NWA 4035 | Morocco                                     | 2005                          | 355                     | 1                   | H6                 | 22.3                        | S2              | W1              | 17.4         | 15.6         | MNB1-1   | —                  |
| NWA 4036 | Morocco                                     | 2005                          | 22,000                  | 1                   | L6 Imc             | 36.2                        | W1              | W1              | 23.4         | 20.3         | MNB1-1   | —                  |
| NWA 4037 | Morocco                                     | 2005                          | >300,000                | n                   | L4-6               | 22.0                        | S3/4            | W1              | 23.3         | 2.1-20.9     | MNB1-1   | —                  |

Table 2. *Continued.* Meteorites from Northwest Africa.

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location | Comments             |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|----------|----------------------|
| NWA 4038 | Morocco                                     | 2005                          | 23                      | 1                   | CO3                | 4.6                         | S2              | W2/3            | 0.4–47.8     | 1–2.2        | MNB1-1   | –                    |
| NWA 4039 | Morocco                                     | 2005                          | 950                     | n                   | Euc                | 20.1                        | High            | Mod             | –            | –            | –        | See separate entry   |
| NWA 4040 | Morocco                                     | 2004                          | 226.3                   | 1                   | L3                 | 24.5                        | S2              | W2              | 0.4–29.5     | 2.8–23.9     | –        | See separate entry   |
| NWA 4041 | Morocco                                     | 2004                          | 42                      | 1                   | LL6                | 8.4                         | S2              | W1              | 28.7         | 23.2         | MNB1-1   | –                    |
| NWA 4042 | Morocco                                     | 2004                          | 56.2                    | 1                   | Aung               | 11.6                        | S2              | W2              | –            | –            | –        | See separate entry   |
| NWA 4043 | Morocco                                     | 2004                          | 40.05                   | 1                   | EL6                | 8.3                         | S2              | W0              | –            | 0.1–1.5      | MNB1-2   | –                    |
| NWA 4044 | Morocco                                     | 2005                          | 1365                    | 1                   | LL6                | 23.2                        | S3              | W1              | 29.9         | 24           | MNB1-1   | –                    |
| NWA 4045 | Morocco                                     | 2005                          | 1000                    | 1                   | L6                 | 33.0                        | S3/4            | W1              | 26.9         | 22.6         | MNB1-1   | –                    |
| NWA 4046 | Morocco                                     | 2005                          | 140                     | 1                   | LL6                | 20.1                        | S3              | W2              | 28.5         | 23           | MNB1-1   | –                    |
| NWA 4047 | Morocco                                     | 2005                          | 200                     | 1                   | H4–5               | 20.1                        | S2              | W0/1            | 16.9         | 4.8–16.3     | MNB1-1   | –                    |
| NWA 4048 | Morocco                                     | 2004                          | 60                      | 1                   | L3                 | 7.4++                       | S2              | W2              | 8.1–28.5     | 1.7–22.9     | MNB1-1   | –                    |
| NWA 4050 | Morocco                                     | 02/2002                       | 58                      | 1                   | H3–4               | 54.0                        | S2              | W2              | 19.3 ± 6.5   | 13.1 ± 4.5   | CSM1-1   | Kamacite (Co = 0.42) |
| NWA 4052 | Er-Mor                                      | Unknown                       | 66                      | 1                   | L3–6               | 14.2                        | S4              | W1              | –            | 1.9–20.8     | MNB1-1   | –                    |
| NWA 4053 | Er-Mor                                      | Unknown                       | 610                     | 1                   | H5                 | 26.8                        | S2              | W2/3            | –            | 15.6         | MNB1-1   | –                    |
| NWA 4054 | Er-Mor                                      | Unknown                       | 454                     | 1                   | H5                 | 23.2                        | S2              | W2              | –            | 15.5         | MNB1-1   | –                    |
| NWA 4055 | Er-Mor                                      | Unknown                       | 196                     | 1                   | L6                 | 22.2                        | S2/3            | W1              | –            | 19.8         | MNB1-1   | –                    |
| NWA 4057 | Er-Mor                                      | Unknown                       | 103.4                   | 1                   | L3                 | 23.8                        | S4              | W1              | –            | 8.6–21.1     | MNB1-1   | –                    |
| NWA 4058 | Er-Mor                                      | Unknown                       | 58                      | 1                   | LL6                | 12.8                        | S4              | W0              | –            | 24.3         | MNB1-1   | –                    |
| NWA 4060 | Er-Mor                                      | Unknown                       | 292                     | 1                   | L6                 | 22.2                        | S4              | W1              | –            | 19.9         | MNB1-1   | –                    |
| NWA 4061 | Er-Mor                                      | Unknown                       | 41.2                    | 1                   | H4/5               | 9.4                         | S2              | W3              | 18           | 16.1         | MNB1-1   | –                    |
| NWA 4063 | Er-Mor                                      | Unknown                       | 132.4                   | 1                   | L6                 | 24.2                        | S4              | W1              | 23.3         | 19.9         | MNB1-1   | –                    |
| NWA 4064 | Er-Mor                                      | Unknown                       | 130.6                   | 1                   | H6                 | 21.0                        | S2              | W3              | 17.4         | 15.5         | MNB1-1   | –                    |
| NWA 4065 | Er-Mor                                      | Unknown                       | 146.4                   | 1                   | H5                 | 23.8                        | S3              | W3              | 18.2         | 16.4         | MNB1-1   | –                    |
| NWA 4066 | Er-Mor                                      | Unknown                       | 254                     | 1                   | H5                 | 22.6                        | S2              | W2              | 16.5         | 14.8         | MNB1-1   | –                    |
| NWA 4067 | Er-Mor                                      | Unknown                       | 335.4                   | 2                   | L5                 | 23.6                        | S3              | W2              | 23.5         | 19.8         | MNB1-1   | –                    |
| NWA 4068 | Er-Mor                                      | Unknown                       | 460                     | 1                   | H4/5               | 22.4                        | S2              | W3              | 17.5         | 15.8         | MNB1-1   | –                    |
| NWA 4069 | Er-Mor                                      | Unknown                       | 506                     | 1                   | L6                 | 21.6                        | S3              | W2              | 23.9         | 20.6         | MNB1-1   | –                    |
| NWA 4072 | Er-Mor                                      | Unknown                       | 334.7                   | 2                   | L5                 | 23.0                        | S4              | W2              | 24.5         | 20.9         | MNB1-1   | –                    |
| NWA 4073 | Er-Mor                                      | Unknown                       | 242                     | 1                   | H6                 | 27.0                        | S3              | W2/3            | 18.9         | 16.5         | MNB1-1   | –                    |
| NWA 4074 | Er-Mor                                      | Unknown                       | 216                     | 1                   | H5/6               | 24.4                        | S4              | W1/2            | 16.7         | 15.5         | MNB1-1   | –                    |
| NWA 4076 | Er-Mor                                      | Unknown                       | 143.1                   | 1                   | L6                 | 23.2                        | S4              | W1              | 23.6         | 20           | MNB1-1   | –                    |
| NWA 4077 | Er-Mor                                      | Unknown                       | 89.9                    | 1                   | H6                 | 19.8                        | S2              | W1              | 17.9         | 15.7         | MNB1-1   | –                    |
| NWA 4078 | Er-Mor                                      | Unknown                       | 53.1                    | 1                   | L5                 | 12.2                        | S3              | W1              | 23.3         | 20.1         | MNB1-1   | –                    |
| NWA 4079 | Er-Mor                                      | Unknown                       | 192                     | 1                   | L5                 | 22.8                        | S4              | W2              | 24.3         | 20.5         | MNB1-1   | –                    |
| NWA 4080 | Er-Mor                                      | Unknown                       | 99.1                    | 1                   | L6                 | 22.0                        | S3              | W1              | 24.5         | 20.4         | MNB1-1   | –                    |
| NWA 4083 | Er-Mor                                      | Unknown                       | 77.5                    | 1                   | L4                 | 18.0                        | S2              | W1              | 27.2         | 14.3–22.2    | MNB1-1   | –                    |
| NWA 4084 | Er-Mor                                      | Unknown                       | 70                      | 1                   | H3                 | 16.2                        | S2              | W1              | 1.1–20.2     | 1.1–16.2     | MNB1-1   | –                    |
| NWA 4085 | Er-Mor                                      | Unknown                       | 68.7                    | 1                   | L6                 | 14.0                        | S4              | W3              | 24.1         | 20.4         | MNB1-1   | –                    |
| NWA 4087 | Er-Mor                                      | Unknown                       | 57.8                    | 1                   | L6                 | 12.8                        | S2              | W0/1            | 23.4         | 20.2         | MNB1-1   | –                    |
| NWA 4089 | Er-Mor                                      | Unknown                       | 96.8                    | 1                   | H/L4/5             | 20.2                        | S2              | W3              | 21.2         | 18.1         | MNB1-1   | –                    |
| NWA 4090 | Er-Mor                                      | Unknown                       | 776.3                   | 6                   | L6                 | 26.8                        | S4              | W1              | 24.4         | 20.6         | MNB1-1   | –                    |
| NWA 4092 | Er-Mor                                      | Unknown                       | 147.6                   | 2                   | L6                 | 23.8                        | S3/4            | W2              | 23.9         | 20           | MNB1-1   | –                    |
| NWA 4094 | Er-Mor                                      | Unknown                       | 143.5                   | 1                   | H4                 | 21.2                        | S3              | W1              | 17.9         | 14.0–17.4    | MNB1-1   | –                    |
| NWA 4097 | Er-Mor                                      | Unknown                       | 128.5                   | 1                   | H/L3               | 27.6                        | S2              | W2              | 7.9–24       | 4.7–18.8     | MNB1-1   | –                    |
| NWA 4098 | Er-Mor                                      | Unknown                       | 1470                    | 1                   | L4/5               | 21.4                        | S3              | W1              | 23.4         | 19.6         | MNB1-1   | –                    |
| NWA 4101 | Er-Mor                                      | Unknown                       | 261                     | 2                   | H4                 | 22.8                        | S2              | W2/3            | 17.6         | 14.0–17.0    | MNB1-1   | –                    |
| NWA 4102 | Er-Mor                                      | Unknown                       | 169                     | 1                   | L6                 | 23.8                        | S4              | W3              | 23.8         | 19.8         | MNB1-1   | –                    |

Table 2. *Continued.* Meteorites from Northwest Africa.

| Name     | Recovery area<br>(or purchase) <sup>a</sup> | Recovery date<br>(dd/mm/yyyy) | TKM <sup>b</sup><br>(g) | Pieces <sup>c</sup> | Class <sup>d</sup> | Ty mass <sup>e</sup><br>(g) | SS <sup>f</sup> | WG <sup>g</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Location  | Comments           |
|----------|---|-------------------------------|-------------------------|---------------------|--------------------|-----------------------------|-----------------|-----------------|--------------|--------------|-----------|--------------------|
| NWA 4103 | Er-Mor                                      | Unknown                       | 112.8                   | 1                   | L3                 | 23.8                        | S4              | W2              | 9.6–24.8     | 2.8–21.8     | MNB1-1    | –                  |
| NWA 4104 | Er-Mor                                      | Unknown                       | 60                      | 1                   | H4                 | 13.0                        | S2              | W2              | 17           | 15.6         | MNB1-1    | –                  |
| NWA 4107 | Er-Mor                                      | Unknown                       | 53.1                    | 1                   | L6                 | 12.0                        | S3              | W2              | 23.9         | 20.6         | MNB1-1    | –                  |
| NWA 4109 | Er-Mor                                      | Unknown                       | 233                     | 1                   | L6                 | 20.8                        | S4              | W3              | 24.3         | 20           | MNB1-1    | –                  |
| NWA 4110 | Er-Mor                                      | Unknown                       | 309                     | 1                   | L6                 | 21.4                        | S3              | W3              | 23.3         | 19.4         | MNB1-1    | –                  |
| NWA 4112 | Er-Mor                                      | Unknown                       | 45                      | 1                   | L6                 | 9.4                         | S3              | W1              | 24.4         | 20.3         | MNB1-1    | –                  |
| NWA 4113 | Er-Mor                                      | Unknown                       | 78.4                    | 2                   | L4                 | 17.8                        | S4              | W1              | 23           | 6.4–19.5     | MNB1-1    | –                  |
| NWA 4116 | Er-Mor                                      | Unknown                       | 90.6                    | 1                   | H5/6               | 20.0                        | S3              | W1              | 17.8         | 16           | MNB1-1    | –                  |
| NWA 4117 | Er-Mor                                      | Unknown                       | 70                      | 1                   | H3/4               | 14.4                        | S2              | W3              | 13.0–14.8    | 10.9–13.0    | MNB1-1    | –                  |
| NWA 4120 | Er-Mor                                      | Unknown                       | 15,200                  | 1                   | L6                 | 28.8                        | S4              | W2/3            | 24.5         | 21           | MNB1-1    | –                  |
| NWA 4121 | Er-Mor                                      | Unknown                       | 750                     | 1                   | L6                 | 22.8                        | S4              | W1              | 24.1         | 20.5         | MNB1-1    | –                  |
| NWA 4123 | Er-Mor                                      | 2004                          | 46.9                    | 1                   | Euc                | 10.0                        | Mod             | Min             | –            | –            | –         | See separate entry |
| NWA 4124 | Er-Mor                                      | Unknown                       | 81.8                    | 1                   | H6                 | 17.4                        | S2              | W3              | 18.1         | 15.7         | MNB1-1    | –                  |
| NWA 4125 | Er-Mor                                      | Unknown                       | 209                     | 1                   | L5/6               | 25.6                        | S4              | W2              | 24.1         | 21.3         | MNB1-1    | –                  |
| NWA 4127 | Er-Mor                                      | Unknown                       | 54.4                    | 1                   | L6                 | 11.8                        | S3              | W3              | 26.7         | 22.6         | MNB1-1    | –                  |
| NWA 4129 | Er-Mor                                      | Unknown                       | 42.9                    | 1                   | H3                 | 10.4                        | S2              | W1              | 0.9–18.7     | 6.3–20.5     | MNB1-1    | –                  |
| NWA 4135 | Er-Mor                                      | Unknown                       | 91.1                    | 1                   | L6                 | 27.2                        | S4              | W2              | 24.6         | 20.5         | MNB1-1    | –                  |
| NWA 4136 | Er-Mor                                      | Unknown                       | 78.2                    | 1                   | L6                 | 17.4                        | S2              | W2              | 24.8         | 20.8         | MNB1-1    | –                  |
| NWA 4137 | Er-Mor                                      | Unknown                       | 121.1                   | 1                   | L6                 | 22.0                        | S3              | W2              | 24.3         | 20.3         | MNB1-1    | –                  |
| NWA 4139 | Er-Mor                                      | Unknown                       | 341                     | 1                   | H6                 | 24.8                        | S2              | W3              | 16.7         | 15.0         | MNB1-1    | –                  |
| NWA 4140 | Er-Mor                                      | Unknown                       | 139.1                   | 1                   | L6                 | 23.6                        | S4              | W1              | 24.2         | 20.1         | MNB1-1    | –                  |
| NWA 4141 | Er-Mor                                      | Unknown                       | 70.3                    | 1                   | L6                 | 16.6                        | S3              | W3              | 22.9         | 20.2         | MNB1-1    | –                  |
| NWA 4142 | Er-Mor                                      | Unknown                       | 63                      | 1                   | H4                 | 13.4                        | S2              | W1              | 16.9         | 14.9         | MNB1-1    | –                  |
| NWA 4143 | Er-Mor                                      | Unknown                       | 74.4                    | 1                   | L4                 | 18.4                        | S3              | W2/3            | 23.8         | 15.3–20.0    | MNB1-1    | –                  |
| NWA 4145 | Er-Mor                                      | Unknown                       | 122.3                   | 1                   | H4                 | 21.6                        | S2              | W1              | 17.3         | 10.5–16.7    | MNB1-1    | –                  |
| NWA 4146 | Er-Mor                                      | Unknown                       | 305.6                   | 2                   | L6                 | 29.6                        | S3              | W3              | 24.1         | 20.6         | MNB1-1    | –                  |
| NWA 4150 | Er-Mor                                      | Unknown                       | 436.4                   | 2                   | H/L6 Imc           | 25.0                        | –               | W0/1            | 24.1         | 19.9         | MNB1-1    | –                  |
| NWA 4152 | Er-Mor                                      | Unknown                       | 91.7                    | 1                   | H/L6               | 24.0                        | S3              | W2              | 20.4         | 20.4         | MNB1-1    | –                  |
| NWA 4153 | Er-Mor                                      | Unknown                       | 75.4                    | 1                   | H/L6               | 19.8B                       | S4              | W3              | 20.1         | 20.1         | MNB1-1    | –                  |
| NWA 4154 | Er-Mor                                      | Unknown                       | 132.5                   | 1                   | H/L6               | 24.8                        | S4              | W4              | 20.6         | 20.6         | MNB1-1    | –                  |
| NWA 4155 | Er-Mor                                      | Unknown                       | 247                     | 1                   | H/L6               | 30.8                        | S4              | W3              | 21.2         | 21.2         | MNB1-1    | –                  |
| NWA 4156 | Er-Mor                                      | Unknown                       | 165                     | 1                   | H/L6               | 29.2                        | S4              | W3              | 19.9         | 19.9         | MNB1-1    | –                  |
| NWA 4157 | Er-Mor                                      | Unknown                       | 135                     | 1                   | L3                 | 33.3                        | S2              | W1              | 18.3         | 14.8         | Vernad1-1 | –                  |
| NWA 4158 | Hamburg                                     | Unknown                       | 77.7                    | 1                   | H5                 | 27.9                        | S4              | W0              | 18.8         | 16           | Vernad1-1 | –                  |
| NWA 4215 | Mha-Mor                                     | 04/2002                       | 46.4                    | 1                   | Dio                | 10.0                        | –               | –               | –            | –            | ENSL      | –                  |

<sup>a</sup>Recovery area (or purchase) = where the meteorites were recovered (or found) or where they were purchased. Ag-Mor = Agadir, Morocco; Er-Mor = Erfoud, Morocco; Mha-Mor = Mhamid, Morocco; Rab-Mor = Rabat, Morocco; Tag-Mor = Tagounite, Morocco; Hamburg = purchased in Hamburg from a dealer from Morocco.

<sup>b</sup>TKM = total known mass.

<sup>c</sup>Pieces = total number of pieces; n = numerous pieces.

<sup>d</sup>Class = classification. Ang = angrite; Aug = achondrite, ungrouped; Bra = brachinite; Dio = diogenite; Acp = acapulcoite; Euc = eucrite; How = howardite; Imc = impact melt rock; Mes = mesosiderite; Pac = primitive achondrite; Ure = ureilite.

<sup>e</sup>Ty mass = type specimen mass. Type masses with ++ next to them all have at least one thin section on deposit and additional materials are expected to be deposited in the future.

<sup>f</sup>SS = shock stage. Min = minimal; Mod = moderate; L-M = low to moderate; M-H = moderate to high.

<sup>g</sup>WG = weathering grade. Mod = moderate; M-M = minimal to moderate.

<sup>h</sup>Reclassification from an winonaite (MB 88) to an acapulcoite. All samples listed above were purchased. All reported analyses are given in wt%, unless otherwise stated.



Table 3. Meteorites from Grove Mountains, Antarctica.

| Name       | Latitude<br>(S) | Longitude<br>(E) | Mass<br>(g) | Date of find | Class <sup>a</sup> | SS <sup>b</sup> | WG <sup>c</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Comments           | Info   |
|------------|-----------------|------------------|-------------|--------------|--------------------|-----------------|-----------------|--------------|--------------|--------------------|--------|
| GRV 024516 | 73°00'          | 75°12'           | 24.7        | 01/2003      | Ure                | –               | –               | –            | –            | See separate entry | IGGCAS |
| GRV 024517 | 73°00'          | 75°12'           | 40.5        | 01/2003      | H5                 | S1              | W2              | 16.0 ± 0.4   | 13.6 ± 0.4   | –                  | IGGCAS |

<sup>a</sup>Class = classification.

<sup>b</sup>SS = shock stage.

<sup>c</sup>WG = weathering grade.

Table 4. Meteorites from North America.

| Name                   | Find site    | Find location      | Latitude<br>(N) | Longitude<br>(W) | Mass<br>(g) | Recovery<br>date<br>(dd/mm/<br>yyyy) | Pieces | Class <sup>a</sup> | SS <sup>b</sup> | WG <sup>c</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Wo<br>(mol%) | Ty<br>mass <sup>d</sup><br>(g) | Location | Comments <sup>e</sup> |
|------------------------|--------------|--------------------|-----------------|------------------|-------------|--------------------------------------|--------|--------------------|-----------------|-----------------|--------------|--------------|--------------|--------------------------------|----------|-----------------------|
| <b>Black Rock</b>      |              |                    |                 |                  |             |                                      |        |                    |                 |                 |              |              |              |                                |          |                       |
| BR 001                 | Dry lake     | Pershing, NV       | 40°52.030'      | 119°11.011'      | 152.0       | 29/08/2003                           | 1      | L6                 | S4              | W2              | 25.3 ± 0.2   | 21.6 ± 0.5   | 1.7 ± 0.2    | 21.1                           | UNMI     | –                     |
| <b>Buck Mountains</b>  |              |                    |                 |                  |             |                                      |        |                    |                 |                 |              |              |              |                                |          |                       |
| BM 001                 | Pediment     | Mojave, AZ         | 34°43.947'      | 114°13.354'      | 50.0        | 29/02/2004                           | 1      | H6                 | S2              | W3              | 18.3 ± 0.3   | 16.8 ± 0.6   | 1.2 ± 0.2    | 10.0                           | CML1-2   | –                     |
| BM 002                 | Pediment     | Mojave, AZ         | 34°43.16'       | 114°12.47'       | 18.4        | 10/04/2004                           | 1      | L6                 | S2              | W2-3            | 25.1 ± 0.7   | 21.1 ± 0.9   | 1.5 ± 0.2    | 9.1                            | CML1-2   | –                     |
| <b>Lucerne Valley</b>  |              |                    |                 |                  |             |                                      |        |                    |                 |                 |              |              |              |                                |          |                       |
| LV 028                 | Dry lake     | San Bernardino, CA | 34°29.25'       | 116°57.34'       | 3.0         | 10/11/2003                           | 1      | CK4                | S2              | –               | 33.2 ± 0.4   | –            | –            | 1.2                            | UCLA1    | See separate entry    |
| LV 029                 | Dry lake     | San Bernardino, CA | 34°29.25'       | 116°57.34'       | 10.1        | 10/11/2003                           | 1      | CK4                | S2              | –               | 33.2 ± 0.4   | –            | –            | 2.2                            | UCLA1    | Paired                |
| LV 030                 | Dry lake     | San Bernardino, CA | 34°29.25'       | 116°57.478'      | 4.1         | 03/07/2004                           | 1      | CK4                | S2              | –               | –            | –            | –            | –                              | UCLA1    | Paired                |
| LV 031                 | Dry lake     | San Bernardino, CA | 34°29.226'      | 116°57.468'      | 3.2         | 03/07/2004                           | 1      | CK4                | S2              | –               | –            | –            | –            | –                              | UCLA1    | Paired                |
| LV 032                 | Dry lake     | San Bernardino, CA | 34°29.409'      | 116°56.991'      | 11.9        | 03/07/2004                           | 1      | CK4                | S2              | –               | –            | –            | –            | 2.1                            | UCLA1    | Paired                |
| LV 035                 | Dry lake     | San Bernardino, CA | 34°29.208'      | 116°57.502'      | 0.9         | 13/03/2004                           | 1      | CK4                | S2              | –               | –            | –            | –            | 0.9                            | UCLA1    | Paired                |
| LV 036                 | Dry lake     | San Bernardino, CA | 34°29.276'      | 116°57.383'      | 2.5         | 13/03/2004                           | 1      | CK4                | S2              | –               | –            | –            | –            | –                              | UCLA1    | Paired                |
| LV 037                 | Dry lake     | San Bernardino, CA | 34°29.332'      | 116°57.134'      | 0.9         | 13/03/2004                           | 1      | CK4                | S2              | –               | –            | –            | –            | 0.9                            | UCLA1    | Paired                |
| LV 050                 | Dry lake     | San Bernardino, CA | 34°29.237'      | 116°57.500'      | 6.85        | 05/02/2005                           | 1      | H4                 | S1              | W2              | 17.8 ± 0.1   | –            | –            | 1.8                            | UCLA1    | –                     |
| Mohawk                 | Desert       | Yuma, AZ           | 32°43.8'        | 113°42.6'        | 586         | 01/10/2000                           | 1      | Iron               | –               | –               | –            | –            | –            | 37.4                           | –        | See separate entry    |
| Orlando                | Residential  | Orange, FL         | 28°32'51"       | 81°21'44"        | 180         | 11/08/2004                           | 1      | Euc                | –               | –               | –            | –            | –            | 20.0                           | –        | See separate entry    |
| Pedernales             | Ranch        | Gillespie, TX      | 30°20'          | 98°57'           | 691         | 12/01/1980                           | 1      | Iron               | –               | –               | –            | –            | –            | 23 and 86                      | –        | See separate entry    |
| Purmela                | Farm         | Coryell, TX        | 29°30'          | 98°3'            | 4500        | 10/01/1977                           | 1      | Iron               | –               | –               | –            | –            | –            | 406.3                          | –        | See separate entry    |
| <b>Sacramento Wash</b> |              |                    |                 |                  |             |                                      |        |                    |                 |                 |              |              |              |                                |          |                       |
| SaW 001                | Flat desert  | Mojave, AZ         | 34°45.382'      | 114°14.061'      | 98.9        | 11/08/2003                           | 1      | H4                 | S2              | W3              | 18.0 ± 3.5   | 15.8 ± 0.3   | 1.1 ± 0.2    | 28.2                           | ASU1     | Not paired            |
| SaW 002                | Desert ridge | Mojave, AZ         | 34°45.381'      | 114°14.058'      | 892.8       | 13/03/2004                           | 2      | H4                 | S1              | W1              | 18.6 ± 0.3   | 9.0 ± 1.0    | 0.4 ± 0.2    | 103.0                          | ASU1     | Not paired            |
| SaW 003                | Desert hill  | Mojave, AZ         | 34°45.524'      | 114°14.005'      | 89.2        | 22/03/2004                           | 1      | H4                 | S2              | W3              | 17.2 ± 5.1   | 15.7 ± 5.0   | 1.1 ± 0.5    | 21.6                           | ASU1-1   | Not paired            |
| SaW 004                | Desert ridge | Mojave, AZ         | 34°45.047'      | 114°13.630'      | 28.6        | 17/04/2004                           | 1      | H5                 | S1              | W3              | 18.3 ± 0.3   | 16.2 ± 0.3   | 1.4 ± 0.3    | 6.3                            | ASU1     | Not paired            |
| <b>Superior Valley</b> |              |                    |                 |                  |             |                                      |        |                    |                 |                 |              |              |              |                                |          |                       |
| SuV 013                | Dry lake     | San Bernardino, CA | 35°14.0'        | 117°02.367'      | 2.7         | 24/08/2002                           | 1      | L5                 | S2              | W4              | 24.7 ± 0.2   | –            | –            | 0.6                            | UCLA1-1  | –                     |

Table 4. *Continued.* Meteorites from North America.

| Name                    | Find site    | Find location       | Latitude (N) | Longitude (W) | Mass (g) | Recovery date (dd/mm/yyyy) | Pieces | Class <sup>a</sup> | SS <sup>b</sup> | WG <sup>c</sup> | Fa (mol%)  | Fs (mol%)  | Wo (mol%) | Ty mass <sup>d</sup> (g) | Location | Comments <sup>e</sup> |
|-------------------------|--------------|---------------------|--------------|---------------|----------|----------------------------|--------|--------------------|-----------------|-----------------|------------|------------|-----------|--------------------------|----------|-----------------------|
| SuV 014                 | Dry lake     | San Bernardino, CA  | 35°14.160'   | 117°02.527'   | 1.8      | 25/08/2002                 | 1      | Acp                | –               | –               | 4.6        | 6.9 ± 0.1  | 2.9 ± 1.6 | 0.5                      | –        | See separate entry    |
| SuV 018                 | Dry lake     | San Bernardino, CA  | 35°14.227'   | 117°00.910'   | 11.3     | 26/08/2002                 | 1      | H5                 | S2              | W5              | 19.1 ± 1.8 | 15.6 ± 2.5 | 1.1 ± 0.4 | 2.5                      | UCLA1    | –                     |
| Trilby Wash             | Basin        | Maricopa County, AZ | 33°55'       | 112°33'       | 846.0    | 13/01/2005                 | 22     | L4                 | S1              | W3              | 27.7 ± 0.2 | 20.7 ± 0.4 | 1.6 ± 0.2 | 87.7                     | ASU1-2   | –                     |
| Warm Springs Wilderness | Desert ridge | Mojave County, AZ   | 34°47.219'   | 114°15.031'   | 156.9    | 22/12/2003                 | 2      | H4–6               | S2              | W1              | 17.7 ± 2.6 | 15.6 ± 1.1 | 1.4 ± 0.2 | 35.2                     | ASU1-1   | Not paired            |

<sup>a</sup>Class= classification.<sup>b</sup>SS = shock stage.<sup>c</sup>WG = weathering grade.<sup>d</sup>Ty mass = type specimen mass.<sup>e</sup>Paired = paired with LV 028. See LV 028 for type specimen mass and depository; Not paired = sample is not paired with Franconia (H5).

Table 5. List of meteorites from the Middle East, the country of Oman.

| Name          | Recovery date (dd/mm/yyyy) | Latitude (N) | Longitude (E) | Mass (g) | Pieces <sup>a</sup> | Class <sup>b</sup> | Shock | WG <sup>c</sup> | Fa (mol%) | Fs (mol%) | Wo (mol%) | Ty mass (g) <sup>d</sup> | Location  | Comments           |
|---------------|----------------------------|--------------|---------------|----------|---------------------|--------------------|-------|-----------------|-----------|-----------|-----------|--------------------------|-----------|--------------------|
| <b>Dhofar</b> |                            |              |               |          |                     |                    |       |                 |           |           |           |                          |           |                    |
| Dho 1278      | 2005                       | 19°35'55.1"  | 55°00'10.6"   | 54.7     | 1                   | H6                 | S2    | W2/3            | 23.6      | 19.6      | –         | 13.5 MNB                 | MNB1-3    | –                  |
| Dho 1279      | 2005                       | 18°15'04.2"  | 54°11'53.6"   | 1588     | 1                   | L6                 | S4    | W3              | 23.8      | 19.3      | –         | 25 MNB                   | MNB1-3    | –                  |
| Dho 1280      | 2005                       | 18°16'37.8"  | 54°15'19.7"   | 4473     | n                   | H6                 | S4    | W2/3            | 16.8      | 15.5      | –         | 28.9 MNB                 | MNB1-3    | –                  |
| Dho 1281      | 2005                       | 18°13'10.1"  | 54°18'18.5"   | 140      | 1                   | L6                 | S2    | W3/4            | 23.2      | 19.3      | –         | 25.2 MNB                 | MNB1-3    | –                  |
| Dho 1282      | 2005                       | 18°33'07.2"  | 54°00'07.8"   | 582      | 1                   | L5                 | S4    | W3              | 23.2      | 20.4      | –         | 20.5 MNB                 | MNB1-3    | –                  |
| Dho 1283      | 2005                       | 18°33'14.1"  | 54°01'10.5"   | 502      | 4                   | L5                 | S4    | W3              | 22.8      | 19.8      | –         | 24.3 MNB                 | MNB1-3    | –                  |
| Dho 1284      | 2005                       | 18°31'55.2"  | 54°02'03.7"   | 558      | 1                   | L5/6               | S4    | W3              | 23.7      | 20.7      | –         | 22.6 MNB                 | MNB1-4    | –                  |
| Dho 1285      | 2002                       | 19°18.15'    | 54°33.3'      | 406      | 11                  | Ure                | –     | –               | –         | 92.48     | –         | –                        | –         | See separate entry |
| Dho 1286      | 12/2005                    | 18°25.579'   | 54°25.719'    | 898      | 2                   | –                  | –     | –               | –         | 78        | –         | 20.6                     | –         | See separate entry |
| Dho 1288      | 12/09/2004                 | 18°16.7'     | 54°17.1'      | 43,600   | n                   | H5                 | S3    | W2/3            | 17.80     | 17.40     | 1.10      | 1683                     | Vernad1-2 | –                  |
| Dho 1289      | 12/03/2004                 | 18°38.0'     | 54°25.3'      | 37,000   | n                   | L4                 | S2    | W3/4            | 24.70     | 21.30     | 1.50      | 1870                     | Vernad1-2 | –                  |
| Dho 1290      | 12/01/2004                 | 18°34.2'     | 54°25.4'      | 1211.3   | 20                  | LL4                | S3    | W0/1            | 28.0      | 21.30     | 1.60      | 270.8                    | Vernad1-2 | Breccia            |
| Dho 1291      | 11/04/2001                 | 18°19.0'     | 54°08.8'      | 66       | 1                   | H3.9               | S2    | W3              | 18.1–26.3 | 15.8–22.4 | 1.50      | 14.1                     | Vernad1-2 | –                  |
| Dho 1292      | 04/07/2001                 | 18°51.1'     | 54°39.7'      | 26       | 1                   | H3.9               | S2    | W3              | 8.2–18.3  | 16.6–21.7 | 1.50      | 10.6                     | Vernad1-2 | –                  |
| Dho 1293      | 14/07/2001                 | 18°50.8'     | 54°39.9'      | 69       | 2                   | H3.9               | S3    | W3              | 18.0–22.4 | 16.60     | 1.20      | 17.5                     | Vernad1-2 | –                  |
| Dho 1294      | 12/02/2003                 | 19°23.9'     | 54°31.8'      | 86       | 1                   | L/LL5              | S1    | W3              | 26.40     | 22.20     | 1.60      | 20.10/13.10              | Vernad1-2 | –                  |
| Dho 1295      | 11/02/2003                 | 19°07.5'     | 54°43.1'      | 30       | 1                   | L/LL6              | S4    | W2              | 25.80     | 22.20     | 1.40      | 7.5/6.8                  | Vernad1-2 | –                  |
| Dho 1296      | 18/01/2002                 | 18°10.0'     | 54°05.0'      | 48       | 1                   | L5                 | S3    | W3              | 25.0      | 22.40     | 1.40      | 15.1/9.8                 | Vernad1-2 | –                  |
| Dho 1297      | 12/12/2001                 | 19°08.4'     | 54°46.2'      | 130 + 54 | 2                   | H5                 | S3    | W3              | 19.40     | 16.70     | 1.70      | 40.7/38.8                | Vernad1-2 | –                  |
| Dho 1298      | 15/01/2002                 | 19°07.0'     | 54°47.1'      | 52       | 1                   | H5                 | S3    | W3/4            | 18.40     | 16.40     | 1.20      | 10.9/12.4                | Vernad1-2 | –                  |
| Dho 1299      | 15/01/2002                 | 19°08.2'     | 54°47.1'      | 20       | 1                   | H6                 | S2    | W3              | 19.90     | 17.30     | 1.40      | 6.5/4.9                  | Vernad1-2 | –                  |
| Dho 1300      | 16/01/2002                 | 19°08.3'     | 54°39.5'      | 42       | 4                   | H6                 | S3    | W2              | 17.20     | 15.70     | 1.20      | 14.6/9.9                 | Vernad1-2 | –                  |
| Dho 1301      | 12/07/2004                 | 18°25.579'   | 54°25.719'    | 898      | 2                   | Peuc               | –     | –               | –         | 32–59     | 1.8–38.9  | 206                      | Vernad2-1 | See separate entry |
| Dho 1302      | 03/12/2005                 | 18°55.168'   | 54°21.742'    | 23.90    | 1                   | How                | –     | –               | –         | 21–48     | 1.1–43.4  | 7.595                    | Vernad2-1 | See separate entry |
| Dho 1303      | 11/12/2002                 | 19°18.15'    | 54°33.3'      | 404      | 7                   | Ure                | S3    | W4              | –         | –         | –         | 92.48                    | Vernad2-1 | See separate entry |

## Recovery

| Recovery date     | Name              | Latitude    | Longitude   | Mass     | Pieces <sup>a</sup> | Class <sup>b</sup> | Shock | WG <sup>c</sup> | Fa     | Fs     | Wo     | Ty mass          | Location  | Comments   |   |
|-------------------|-------------------|-------------|-------------|----------|---------------------|--------------------|-------|-----------------|--------|--------|--------|------------------|-----------|--|---|
| (dd/mm/yyyy)      |                   | (N)         | (E)         | (g)      |                     |                    |       |                 | (mol%) | (mol%) | (mol%) | (g) <sup>d</sup> |           |  |   |
| Jiddat al Harasis | Dho 1304          | 18°40.032'  | 54°21.972'  | 85       | 1                   | H5                 | S1    | W3-4            | 18.7   | 16.6   | –      | 18.17            | Vernad2-1 | –  |   |
|                   | Dho 1305          | 18°47.705'  | 54°09.077'  | 34       | 1                   | H5                 | S2    | W3              | 18.5   | 16.3   | –      | 15.08            | Vernad2-1 | –  |   |
|                   | Dho 1306          | 18°20.697'  | 54°15.525'  | 104      | 1                   | L4                 | S1    | W4              | 22.6   | 19.9   | –      | 22.55            | Vernad2-1 | –  |   |
|                   | Dho 1307          | 19°19.719'  | 54°32.258'  | 90       | 1                   | H4                 | S1    | W4              | 17.6   | 17.9   | –      | 25.16            | Vernad2-1 | –  |   |
|                   | Dho 1308          | 18°43.795'  | 54°22.478'  | 192      | 12                  | H5                 | S1    | W4              | 18.9   | 16.8   | –      | 41.5             | Vernad2-1 | –  |   |
|                   | Dho 1309          | 19°17.463'  | 54°33.81'   | 204      | 1                   | H4                 | S2    | W4              | 17.4   | 15.8   | –      | 41.9             | Vernad2-1 | –  |   |
|                   | Dho 1310          | 17°01/2002  | 54°35.7'    | 76       | 1                   | L5                 | S4    | W3/4            | 24.90  | 21.40  | 1.60   | 17.3/15.9        | Vernad1-2 | –  |   |
|                   | Dho 1311          | 18°03/2005  | 54°31.8'    | 66       | 1                   | H5                 | S2    | W3              | 18.6   | 16.2   | 1.4    | 23.6             | Vernad3-1 | –  |   |
|                   | Dho 1312          | 19°6.3'     | 54°50.6'    | 86       | 1                   | L/L14              | S4    | W1              | 26.4   | 22.5   | 1.5    | 18               | Vernad1-2 | –  |   |
|                   | Dho 1313          | 18°02/2004  | 54°27.3'    | 154      | 1                   | H6                 | S5    | W2              | 17.7   | 15.3   | 1.5    | 44.5             | Vernad3-1 | –  |   |
|                   | Dho 1314          | 18°02/2004  | 54°34.1'    | 104      | 1                   | H5                 | S2    | W3              | 17.7   | 15.5   | 1.3    | 24               | Vernad3-1 | Ca-Pyx: Wo <sub>43.8</sub> Fs <sub>5.0</sub>                   |   |
|                   | Dho 1315          | 12/12/2001  | 54°46.5'    | 12       | 1                   | L/L15              | S4    | W3/4            | 26.4   | 21.8   | 1.7    | 4.1/4.3          | Vernad1-2 | –  |   |
|                   | Dho 1316          | 10/02/2003  | 54°23.6'    | 30       | 1                   | L/L16              | S3    | W4              | 26.4   | 22.5   | 1.5    | 11.2/8.4         | Vernad1-2 | –  |   |
|                   | Dho 1317          | 05/04/2001  | 18°25.2'    | 54°06.2' | 12                  | 1                  | L6    | S3              | W1     | 24.9   | 21     | 1.6              | 4.2       | Vernad1-2  | – |
|                   | Jiddat al Harasis | Jah 056     | 19°13.8'    | 55°10.9' | 20                  | 1                  | H5    | S2              | W3     | 19.20  | 16.60  | 1.40             | 4.5/4.6   | Vernad1-2  | – |
|                   |                   | Jah 057     | ~19.8°      | 54.6°    | 246                 | 1                  | L6    | S4              | W2     | 24.60  | 20.40  | –                | 56        | Vernad3-1  | – |
|                   |                   | Jah 123     | 19°44.0'    | 55°44.0' | 248                 | 1                  | L15   | S2              | W3     | 27.7   | 22.5   | 2                | 45.2      | Vernad3-1  | – |
| Jah 124           |                   | ~19.8°      | 54.8°       | 1414     | 12                  | L5                 | S3    | W4              | 24.30  | 20.60  | –      | 276              | Vernad3-1 | –  |   |
| Jiddat al Harasis | Jah 125           | 19°37.4'    | 55°0.6'     | 36       | 1                   | H6                 | S2    | W4              | 19.3   | 16.4   | 1.3    | 10.2             | Vernad3-1 | Ca-Px: Wo <sub>42.4</sub> Fs <sub>8.0</sub>                    |   |
|                   | Jah 130           | 19°35'55.1" | 55°00'10.6" | 54.7     | 1                   | L6                 | S2    | W2/3            | 23.6   | 19.6   | –      | 13.5 MNB         | MNB1-3    | –  |   |
|                   | Sayh al Uhaymir   | SaU 176     | 21°02.9'    | 57°15.5' | 19                  | 1                  | H6    | S2-3            | W4     | 19.00  | 16.8   | 1.6              | 4.3/4.4   | Vernad1-2  | – |
|                   |                   | SaU 177     | 20°59.7'    | 57°19.5' | 48                  | 1                  | L16   | S1              | W2     | 26.1   | 21.2   | 1.7              | 11.6/9.6  | Vernad1-2  | – |
| SaU 309           |                   | 20°55'39.6" | 57°16'56.4" | 714      | 49                  | L6                 | S4    | W3              | 23.7   | 20.7   | –      | 23.9 MNB         | MNB1-3    | –  |   |
| SaU 310           |                   | ~21.2°      | 57.1°       | 94       | 3                   | L6                 | S4    | W4              | 24.40  | 20.90  | –      | 30               | Vernad3-1 | –  |   |
| SaU 311           |                   | 15/02/2004  | 21°0.1'     | 214      | 1                   | H4                 | S2    | W3              | 16.8   | 14.7   | 1      | 40.7             | Vernad3-1 | Ca-Px: Wo <sub>43.4</sub> Fs <sub>4.8</sub>                    |   |
| SaU 312           |                   | 20/02/2004  | 20°36.7'    | 278      | 1                   | L6                 | S3    | W4              | 23.8   | 20     | 1.4    | 59               | Vernad3-1 | Ca-Px: Wo <sub>42.6</sub> Fs <sub>8.9</sub>                    |   |
| SaU 313           |                   | 20/02/2004  | 20°40.8'    | 412      | 1                   | H4                 | S2    | W3              | 17.2   | 15.7   | 1.2    | 82.7             | Vernad3-1 | Ca-Px: Wo <sub>40.1</sub> Fs <sub>5.5</sub>                    |   |
| SaU 314           |                   | 21/02/2004  | 21°0.7'     | 4        | 1                   | H5                 | S3    | W4              | 17.8   | 15.4   | 1      | 2                | Vernad3-1 | Ca-Px: Wo <sub>43.7</sub> Fs <sub>5.4</sub>                    |   |
| SaU 315           |                   | 21/02/2005  | 21°0.8'     | 6        | 1                   | H6                 | S2    | W4              | 17.4   | 15.5   | 1.4    | 4.7              | Vernad3-1 | –  |   |
| SaU 316           |                   | 21/02/2006  | 21°0.3'     | 10       | 1                   | L4                 | S4    | W4              | 22.8   | 19     | 1.2    | 3.6              | Vernad3-1 | –  |   |
| Sayh al Uhaymir   | SaU 317           | 15/02/2004  | 20°59.7'    | 8        | 1                   | H5                 | S2    | W3              | 17.5   | 15.3   | 1.2    | 2.2              | Vernad3-1 | Ca-Px: Wo <sub>43.4</sub> Fs <sub>8.0</sub>                    |   |
|                   | SaU 318           | 15/02/2005  | 20°59.9     | 70       | 3                   | H5                 | S2    | W4              | 17.5   | 15.3   | 1.5    | 32.8             | Vernad3-1 | –  |   |
|                   | SaU 319           | 15/02/2006  | 21°5.4'     | 550      | n                   | H5                 | S2    | W4              | 18     | 15.7   | 1.3    | 117.2            | Vernad3-1 | –  |   |
|                   | SaU 320           | 21°4.5'     | 57°15.8'    | 18       | 1                   | H6                 | S3    | W2              | 18.3   | 16.1   | 1.4    | 6.3              | Vernad3-1 | –  |   |
|                   | SaU 321           | 22/02/2004  | 21°5.9'     | 220      | 12                  | L5                 | S3    | W3              | 24.1   | 19.8   | 1.4    | 53.9             | Vernad3-1 | –  |   |
|                   | SaU 322           | 05/03/2005  | 21°3.6'     | 468      | 1                   | H5                 | S6    | W2              | 17.4   | 15     | 1      | 102.3            | Vernad3-1 | Ca-Px: Wo <sub>43.7</sub> Fs <sub>5.1</sub><br>Partial melting |   |
|                   | SaU 323           | 10/12/2001  | 20°59.8'    | 112      | 1                   | H6                 | S3    | W2              | 19.7   | 16.7   | 1.3    | 24.3/30.3        | Vernad1-2 | –  |   |
|                   | SaU 324           | 11/01/2002  | 20°59.8'    | 42       | 2                   | H/L6               | S3    | W3/4            | 20.3   | 17.6   | 1.5    | 17/13.5          | Vernad1-2 | –  |   |
|                   | SaU 325           | 22/02/2004  | 21°0.3'     | 76       | 1                   | H5                 | S4    | W2              | 17.8   | 15.5   | 1.2    | 19.6             | Vernad3-1 | Ca-Px: Wo <sub>41.9</sub> Fs <sub>4.6</sub>                    |   |
|                   | SaU 326           | 22/02/2004  | 21°0.5'     | 1352     | 1                   | L6                 | S4    | W3              | 23.1   | 19.4   | 1.5    | 218              | Vernad3-1 | Ca-Px: Wo <sub>42.3</sub> Fs <sub>6.8</sub>                    |   |
| SaU 402           | 06/26/1905        | 21°4'37"    | 57°16'11"   | –        | –                   | –                  | –     | –               | –      | –      | –      | –                | –         |  |   |

Table 5. *Continued.* List of meteorites from the Middle East, the country of Oman.

| Recovery     |  | Name      | date<br>(dd/mm/yyyy) | Latitude<br>(N) | Longitude<br>(E) | Mass<br>(g) | Pieces <sup>a</sup> | Class <sup>b</sup> | Shock | WG <sup>c</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Wo<br>(mol%) | Ty mass<br>(g) <sup>d</sup> | Location | Comments |
|--------------|--|-----------|----------------------|-----------------|------------------|-------------|---------------------|--------------------|-------|-----------------|--------------|--------------|--------------|-----------------------------|----------|----------|
|              |  |           |                      |                 |                  |             |                     |                    |       |                 |              |              |              |                             |          |          |
| <b>Shiṣr</b> |  |           |                      |                 |                  |             |                     |                    |       |                 |              |              |              |                             |          |          |
|              |  | Shiṣr 045 | 2005                 | 18°09'06.1"     | 53°56'53.6"      | 2298        | 1                   | H6                 | S3    | W2/3            | 17           | 15.2         | –            | 27.7                        | MNB1-4   | –        |
|              |  | Shiṣr 046 | 2005                 | 18°32'55.1"     | 53°57'01.8"      | 182         | 21                  | L5                 | S3    | W2              | 23.1         | 19.5         | –            | 23.2                        | MNB1-4   | –        |
|              |  | Shiṣr 047 | 2005                 | 18°32'50.8"     | 53°59'39.4"      | 396         | 3                   | L6                 | S4    | W3              | 22.7         | 19.1         | –            | 23.7                        | MNB1-4   | –        |
|              |  | Shiṣr 048 | 2005                 | 18°32'50.9"     | 53°59'59.5"      | 375         | 1                   | L5/6               | S3    | W3              | 23           | 20.2         | –            | 21.7                        | MNB1-4   | –        |

<sup>a</sup>n = numerous samples.  
<sup>b</sup>Class = classification. Euc = eucrite; How = howardite; Peuc = polymict eucrite; Ure = ureilite.  
<sup>c</sup>WG = weathering grade.  
<sup>d</sup>Ty mass = type specimen mass. MNB after a type mass indicates that it is deposited at *MNB*. For those masses where two entries occur in the cell, the first is on deposit at *l'ernad*, the second at *MSU*.

Table 6. Meteorites from South America.

| Name                 | Find site  | Find location           | Latitude<br>(S) | Longitude<br>(W) | TKM <sup>a</sup><br>(g) | Found<br>(dd/mm/yyyy)        | Pieces | Class <sup>b</sup> | SS <sup>c</sup> | WG <sup>d</sup> | Fa<br>(mol%) | Fs<br>(mol%) | Wo<br>(mol%) | Ty mass<br>(g) <sup>e</sup> | Location | Comments                 |
|----------------------|------------|-------------------------|-----------------|------------------|-------------------------|------------------------------|--------|--------------------|-----------------|-----------------|--------------|--------------|--------------|-----------------------------|----------|--------------------------|
|                      |            |                         |                 |                  |                         |                              |        |                    |                 |                 |              |              |              |                             |          |                          |
| Mercedes             | Small hill | Argentina               | 34°40'          | 59°20'           | 3301.0                  | 22/12/1994 and<br>25/05/1995 | 26     | H5                 | –               | W3              | 18.9 ± 0.7   | 17.6 ± 2.6   | 1.2 ± 0.5    | 22.0                        | LPL1     | –                        |
| Minas Gerais b       | Collection | Minas Gerais,<br>Brazil |                 |                  | 42.6                    | 00/00/2001                   | 1      | H4                 | S3              | W1–2            | 19.2 ± 2.0   | 15.1 ± 5.6   | 1.4 ± 1.1    | 11.3                        | MIN-KB1  | –                        |
| San Pedro<br>Jacuaro | Mountains  | Ocampo,<br>Mexico       | 19°46'          | 100°39'          | 460                     | 12/01/1968                   | 1      | LL6                | S2              | W0              | 27.7         | 23.2         | 2            | 20.0                        | –        | See<br>separate<br>entry |

<sup>a</sup>TKM = total known mass.  
<sup>b</sup>Class = classification.  
<sup>c</sup>SS = shock stage.  
<sup>d</sup>WG = weathering grade.  
<sup>e</sup>Ty mass = type specimen mass.