

LATE PLEISTOCENE-RECENT ATMOSPHERIC $\delta^{13}\text{C}$ RECORD IN C_4 GRASSES¹

LAURENCE J. TOOLIN

NSF Accelerator Facility for Radioisotope Analysis, The University of Arizona, Tucson, Arizona 85721 USA

and

CHRISTOPHER J. EASTOE

Department of Geosciences, The University of Arizona, Tucson, Arizona 85721 USA

ABSTRACT. Samples of *Setaria* species from packrat middens, herbarium specimens and modern plants preserve a record of $\delta^{13}\text{C}$ of atmospheric CO_2 from 12,600 BP to the present. No secular trend is detected between 12,600 and 1800 BP, when the mean value of $\delta^{13}\text{C}$ during that period was $-6.5 \pm 0.1\text{‰}$ (the error is the standard deviation of the mean). Our value agrees with $\delta^{13}\text{C}$ averages of pre-industrial CO_2 from polar ice cores, and differs significantly from modern regional ($-8.2 \pm 0.1\text{‰}$) and global (-7.7‰) values, which are higher because of fossil fuel burning.

INTRODUCTION

Recently, Marino and McElroy (1991) showed that carbon isotope ratios in a C_4 grass, *Zea mays* (cultivated corn) tracked changes in the carbon isotope ratios of atmospheric CO_2 from the years 1948–1987. Here, we present $\delta^{13}\text{C}$ and ^{14}C age data for species of the C_4 grass genus, *Setaria*. C_4 plants have advantages over C_3 plants as proxies for atmospheric carbon isotope studies. Briefly, in C_3 plants, the fractionation of C isotopes that occurs during photosynthesis is affected by environmental factors, such as temperature and atmospheric CO_2 concentrations. Fractionation in C_4 plants is little influenced by such factors and is more uniform (O’Leary 1988). Tiezen and Boutton (1989) confirm reduced variation in C_4 grasses. Henderson, von Caemmerer and Farquhar (1992) explore details of photosynthetic fractionation in C_4 dicots and monocots.

Two *Setaria* species, *S. macrostachya* and *S. leucopila*, are common today across the southwest USA and into central Mexico (Rominger 1962). Both respond to summer rains, growing and flowering from July to October. These species are commonly preserved in middens constructed by packrats (*Neotoma*). Packrat middens are well-known reservoirs of paleoecological information and have been the subject of intensive study (Betancourt, Van Devender & Martin 1990). Packrats forage up to 50 m from their dens for plants for food and den-building, sampling the local vegetation. In dry shelters in arid climates, middens of plant matter and fecal pellets at the den may become indurated with crystallized urine, and persist for tens of thousands of years.

One of us (L. J. T.) has studied grasses preserved in dozens of packrat middens from sites across the American Southwest (e.g., Betancourt 1984; Van Devender & Toolin 1983; Van Devender, Toolin & Burgess 1990). Floral parts of *Setaria macrostachya* have been found in many midden assemblages from Arizona to Mexico (Fig. 1). The fragments studied for identification include the distinctive, indurated floral bracts (lemma and palea) that enclose the reproductive organs.

METHODS AND RESULTS

All samples were given standard acid-base-acid treatment to remove carbonate and humic contaminants, and combusted to CO_2 . Values of $\delta^{13}\text{C}$ of the CO_2 were measured with a 1σ analy-

¹This paper was presented at the 14th International ^{14}C Conference, 20–24 May 1991, Tucson, Arizona.




Fig. 1. Sample location map. Locations of Holocene packrat middens are indicated by letters (e.g., BB, Big Boy). Locations of modern (1990) specimens are within the cross-hatched area near Tucson.

tical precision of 0.13‰, based on repeated measurements of a laboratory standard calibrated to PDB. The CO_2 was then reduced to graphite (Slota *et al.* 1987) for ^{14}C dating by accelerator mass spectrometry (AMS) (Linick *et al.* 1986; Donahue, Jull & Toolin 1990). Although it is preferable to measure $\delta^{13}\text{C}$ on plant cellulose rather than on a mix of tissues, the small mass of *Setaria* fragments found in most samples (<1mg–5mg) precludes further loss of datable material that would occur on cellulose extraction. Direct dating of the *Setaria* was imperative because all plant species in a given midden may not be contemporaneous (Van Devender *et al.* 1985). In fact, in the course of this study, we found that the ^{14}C age of the *Setaria* differed greatly from previous measurements on other plants from the same middens. For example, Waterman Mountains #2 was originally dated on juniper twigs at 21,510 BP, whereas the *Setaria* from this midden dated to 9895 BP. The lack of contemporaneity of different plant species in a single midden calls into question the value of results in which the age of one plant, or of bulk materials, is assumed to be the same as for other plants. This situation appears to have been overlooked by Marino *et al.* (1992).

Five midden samples were large enough to compare cellulose (method modified after Green (1963)) and mixed-tissue values for $\delta^{13}\text{C}$ (Table 1). These results can be compared to those from modern (AD 1990) *Setaria* samples. Modern samples were collected in eastern Pima and adjacent Santa Cruz counties, Arizona, and from The University of Arizona (ARIZ) herbarium. Cellulose *versus* mixed-tissue values for midden fragments differed less in the older specimens than in modern material (Table 1). We believe this is because the modern material contained more varied tissues (anthers, pistils, *etc.*) than the midden samples, where only the indurated, higher-cellulose-content lemmas and paleas remain. This effect is consistent with variations in different tissues of maize kernels (Tiezen & Fagre 1993).

TABLE 1. Mixed-Tissue *versus* Cellulose $\delta^{13}\text{C}$ (PDB)

Sample no.	Mixed-tissue (‰)	Cellulose (‰)	Difference (‰)
<i>Modern Samples</i>			
AA-6630	-11.8	-11.0	0.8
AA-6631	-11.9	-11.4	0.5
AA-6646a	-11.7	-10.9	0.8
AA-6646b	-11.7	-10.9	0.8
AA-6342a	-11.8	-11.1	0.7
AA-6342b	-11.6	-11.1	0.5
AA-6864	-12.4	-11.5	0.9
AA-6644	-11.6	-10.9	0.7
AA-6343	-11.4	-10.8	0.6
Mean = 0.7			
<i>Holocene Samples</i>			
AA-6634	-9.9	-9.6	0.3
AA-7056	-9.7	-9.5	0.2
AA-7058	-9.5	-9.1	0.4
AA-7059	-10.3	-10.1	0.2
AA-7061	-10.3	-10.0	0.3
Mean = 0.3			

TABLE 2. Herbarium and Modern Samples

Arizona county	Year	Herbarium no. (ARIZ)	AA-no.	$\delta^{13}\text{C}$ (‰, PDB)
Pima	1939	23344	7704	-10.4
Pima	1949	77865	7648	-9.8
Pima	1957	123359	7649L*	-10.4
Pima	1968	169494	7650L	-11.9
Pima	1978	211886	7651L	-11.8
Pima	1986	285825	7652M	-11.9
Pima	1990	290001	6644L	-11.6
Pima	1990	289153	6343L	-11.4
Pima	1990	---	6864M	-12.4
Pima	1990	---	6630L	-11.8
Pima	1990	290010	6631M	-11.9
Santa Cruz	1990	289593	6646aL	-11.7
Santa Cruz	1990	289593	6646bL	-11.7
Santa Cruz	1990	289594	6342aM	-11.8
Santa Cruz	1990	289594	6342bM	-11.6

*Sample no. suffixes: L = *S. leucopila*; M = *S. macrostachya*; a, b indicate different plants collected under the same field number.

TABLE 3. Specimens from Packrat Middens

Midden	Locality*	AA-no.	¹⁴ C age (BP)	δ ¹³ C
Waterman Mts 1A	Pima, AZ	7058	2990 ± 55	-9.5
Waterman Mts 1C	Pima, AZ	7313	1795 ± 70	-10.6
Waterman Mts 1E	Pima, AZ	7059	2080 ± 55	-9.6
Waterman Mts 2	Pima, AZ	6862	9895 ± 160	-9.4
Waterman Mts 9A2	Pima, AZ	7049	10335 ± 75	-9.6
Waterman Mts 9B	Pima, AZ	7050	10415 ± 70	-9.6
Waterman Mts 9C	Pima, AZ	7033	9485 ± 120	-9.4
Waterman Mts 9D	Pima, AZ	7051	5610 ± 70	-10.0
Waterman Mts 10	Pima, AZ	7034	8935 ± 105	-9.5
Waterman Mts 12A	Pima, AZ	7035	6015 ± 75	-9.0
Waterman Mts 12A	Pima, AZ	7060	6125 ± 70	-9.4
Waterman Mts 13A1	Pima, AZ	6863	9340 ± 90	-8.8
Ajo Mts AC1B	Pima, AZ	6951	11100 ± 175	-9.6
Ajo Mts MH1D	Pima, AZ	6953	11240 ± 115	-11.0
Big Boy 3	Otero, NM	6952	10365 ± 255	-9.7
Tank Trap 2	El Paso, TX	6860	9825 ± 175	-9.5
Navar Ranch 1C1	El Paso, TX	7056	6005 ± 65	-9.7
Navar Ranch 4B	El Paso, TX	7061	10065 ± 85	-10.0
Navar Ranch 11	El Paso, TX	7047	9785 ± 75	-9.9
Navar Ranch 12	El Paso, TX	7048	9560 ± 85	-9.7
Navar Ranch 14D	El Paso, TX	7037	9885 ± 90	-10.5
Navar Ranch 18C	El Paso, TX	7038	7950 ± 60	-8.9
Navar Ranch 19C	El Paso, TX	7036	9125 ± 60	-9.6
Maravillas Canyon 13	Brewster, TX	7046	2790 ± 50	-10.2
Maravillas Canyon 16	Brewster, TX	7055	6800 ± 60	-9.4
Puerto de Ventanillas	Coahuila, Mex.	7314	12605 ± 190	-9.4

*AZ = Arizona; NM = New Mexico; TX = Texas; Mex. = Mexico. No county given for Mexican sample.

Table 2 presents mixed-tissue values of δ¹³C for all recent samples. δ¹³C measurements for modern *S. macrostachya* and *S. leucopila* (indicated by M or L in the table) were essentially the same, ranging from -11.6 to -12.4‰ and -11.5 to -12.8‰, respectively. For nine modern (1990) plants, δ¹³C averages -11.90 ± 0.14‰.

Table 3 presents the results of our measurements on *Setaria* specimens from fossil packrat middens. The two species cannot be distinguished in this material. We show the values of δ¹³C of florets (δ_p¹³). The data for 26 midden samples of *Setaria* (12,600–1800 BP) average -9.67 ± 0.10‰, and the data from the herbarium samples (AD 1939–1986) range between the data for ancient and 1990 samples (Table 2).

Farquhar (1983) and Henderson, von Caemmerer and Farquhar (1992) have established a function for relating the δ¹³C of C₄ plants (δ_p¹³) to δ_a¹³, the δ¹³C of the atmospheric CO₂ fixed by the plant. For our *Setaria*, the relation between δ_p¹³ of florets and δ_a¹³ (Farquhar, personal communication) is

$$\delta_a^{13} = 2.9 + \delta_p^{13} \quad (1)$$

To account for the offset between mixed-tissue and cellulose δ^{13}_p values, we added the mean difference of the Holocene samples, 0.3‰ from Table 1, to Equation (1); thus the relation for midden florets becomes

$$\delta^{13}_a = 3.2 + \delta^{13}_p \quad . \quad (2)$$

For the AD 1939–1990 samples, we added the mean differences for 1990 samples (0.7, from Table 3) to Equation (1) and applied the relation

$$\delta^{13}_a = 3.6 + \delta^{13}_p \quad . \quad (3)$$

We applied these corrections (Eq. 2, 3) to relate all of our δ^{13}_p measurements to a common datum, independent of variability in the character of the mixed tissues, following Marino and McElroy (1991). Figure 2 presents the resulting δ^{13}_a data for all samples, derived with the above equations.




Fig. 2. $\delta^{13}\text{C}$ values of Holocene atmospheric CO_2 . • = *Setaria* data; — = mean $\pm 1\sigma$ of pre-1939 samples. --- = scatter of individual measurements about the mean; ● = mean of data for samples collected in 1990; the error bar = 1σ . Direct measurements: Δ , \square , \circ = the mean of Arctic and Antarctic ice core pre-industrial $\delta^{13}\text{C}$ measurements.

The mean δ^{13}_a from nine 1990 *Setaria* samples is $-8.2 \pm 0.14\text{‰}$, lower than the 1986 global mean of -7.7‰ (Keeling *et al.* 1989), but in line with differences between $\delta^{13}\text{C}$ values from clean-air and inland sites (Keeling 1961). Our value is empirically supported by comparison with an N_2O -corrected, seasonally averaged, mean atmospheric $\delta^{13}\text{C}$ value of -8.2‰ , measured in 1983–1984 at Kitt Peak, southern Arizona (Leavitt & Long 1989).

From Equation (2), the mean value of δ^{13}_a for all 26 midden samples is $-6.5 \pm 0.1\text{‰}$, which differs significantly from the modern and global values. Our Holocene mean agrees with $\delta^{13}\text{C}$ data for CO_2 in pre-industrial ice from Antarctica (Siple, $-6.5 \pm 0.07\text{‰}$; South Pole $-6.70 \pm 0.13\text{‰}$; Byrd, $-6.49 \pm 0.05\text{‰}$, and Dye 3, Greenland, $-6.41 \pm 0.09\text{‰}$) (Leuenberger, Siegenthaler & Langway 1992). The scatter of each of our Holocene measurements (0.5‰) is close to that of the modern samples

samples (0.42‰). This suggests that long-term changes in δ_{a}^{13} from 12,600–1800 BP could not have exceeded a few tenths of 1‰. Although our data indicate no secular trend in δ_{a}^{13} during the Holocene, events of short duration or small amplitude might remain undetected.

The nearly step-wise drop of 1.8‰ between 1957 and 1968 may be real, the few data notwithstanding. This interval of time corresponds approximately with a surge in the burning of fossil fuels in the southwestern USA, as a result of urban growth and ore smelting, and to a global increase in fossil fuel CO₂ production (Marland 1990). We plan to make more measurements on 20th century specimens to determine this segment of the δ_{a}^{13} vs. time curve with better precision.

Recently, Martinelli *et al.* (1991) showed that large outputs of biogenically fractionated CO₂ from the Amazon River produced a gradient in δ_{a}^{13} that could mimic secular changes produced, for example, by changes in atmospheric CO₂ concentration. In our region, there is no such large biogenic effect today, and none appears to have operated in the past. If our Holocene samples are grouped by region, there is no distinction between the western group (Arizona) with mean $\delta_{\text{p}}^{13} = -9.6 \pm 0.15$ ‰ and the eastern group (Texas and New Mexico) with mean $\delta_{\text{p}}^{13} = -9.7 \pm 0.13$ ‰.

CONCLUSION

Our results provide the first measurements of C-isotope values of Holocene atmospheric CO₂, constrained by AMS ¹⁴C dating of the carbon used for the stable-isotope measurements. Reconstructions of δ_{a}^{13} such as ours may help clarify the interpretation of δ_{p}^{13} changes in C₃ plant tissues (e.g., tree rings) in which C-isotope fractionation is strongly influenced by environmental factors (Leavitt & Danzer 1991; Krishnamurthy & Epstein 1990).

ACKNOWLEDGMENTS

We are greatly indebted to our colleagues at the University of Arizona. D. J. Donahue and A. J. T. Jull reviewed the manuscript and made important suggestions. T. Lange provided support for the laboratory work. C. T. Mason, Jr., gave access to Herbarium specimens. T. R. Van Devender of the Arizona-Sonora Desert Museum, Tucson, provided all midden *Setaria*. We also thank B. Marino, U. Siegenthaler and G. Farquhar for helpful reviews of an earlier manuscript. S. Leavitt provided further review and much valuable discussion.

REFERENCES

- Betancourt, J. L. 1984 Late Quaternary plant zonation and climate in southeastern Utah. *Great Basin Naturalist* 44: 1–35.
- Betancourt, J. L., Van Devender, T. R. and Martin, P. S., eds., 1990 *Packrat Middens – The Last 40,000 Years of Biotic Change*. Tucson, The University of Arizona Press: 467 p.
- Donahue, D. J., Jull, A. J. T. and Toolin, L. J. 1990 Radiocarbon measurements at the University of Arizona AMS Facility. *Nuclear Instrumentation and Methods in Physics Research B52*: 224–228.
- Farquhar, G. D. 1983 On the nature of carbon isotope discrimination in C₄ species. *Australian Journal of Plant Physiology* 10: 205–226.
- Green, J. W. 1963 Wood cellulose. In Whistler, R. L., ed., *Methods of Carbohydrate Chemistry III*. New York, Academic Press: 9–21.
- Henderson, F. A., Von Caemmerer, G. D. and Farquhar, G. D. 1992 Short-term measurements of carbon isotope discrimination in several C₄ species. *Australian Journal of Plant Physiology* 19: 263–285.
- Keeling, C. D. 1961 The concentration and isotopic abundances of carbon dioxide in rural and marine air. *Geochimica et Cosmochimica Acta* 24: 227–298.
- Keeling, C. D., Bacastow, R. B., Carter, A. F., Piper, S. C., Whorf, T. F., Heimann, M., Mook, W.G. and Roeloffzer, H. 1989 A three dimensional model of atmospheric CO₂ transport based on observed winds: 1. Analysis of observational data. *American Geophysical Monograph* 55: 165–236.
- Krishnamurthy, R. V. and Epstein, S. 1990 Glacial-interglacial excursion in the concentration of atmospheric CO₂: Effect in the ¹³C/¹²C ratio in wood cellulose. *Tellus* 42B: 423–434.
- Leavitt, S. W. and Danzer, S. R. 1992 $\delta^{13}\text{C}$ variations in C₃ plants over the past 50,000 years. In Long, A.

- and Kra, R. S., eds., Proceedings of the 14th International ^{14}C Conference. *Radiocarbon* 34(3): 783–791.
- Leavitt, S. W. and Long, A. 1989 Variation of concentration, ^{14}C activity, and $^{13}\text{C}/^{12}\text{C}$ ratios of CO_2 in air samples from Kitt Peak, Arizona. In Long, A. and Kra, R. S., eds., Proceedings of the 13th International ^{14}C Conference. *Radiocarbon* 31(3): 464–468.
- Leuenberger, M., Siegenthaler, U. and Langway, C. 1992 Carbon isotope composition of atmospheric CO_2 during the last ice age from an Antarctic ice core. *Nature* 357: 488–490.
- Linick, T. W., Jull, A. J. T., Toolin, L. J. and Donahue, D. J. 1986 Operation of the NSF-Arizona accelerator facility for radioisotope analysis, and results from selected collaborative research projects. In Stuiver, M. and Kra, R. S., eds., Proceedings of the 12th International ^{14}C Conference. *Radiocarbon* 28(2A): 522–533.
- Marino, B. D. and McElroy, M. B. 1991 Isotope composition of atmospheric CO_2 inferred from carbon in C_4 plant cellulose. *Nature* 349: 127–131.
- Marino, B. D., McElroy, M. B., Salawitch, R. J. and Spaulding, W. G. 1992 Glacial-to-interglacial variations in the carbon isotopic composition of atmospheric CO_2 . *Nature* 357: 461–466.
- Marland, G. 1990 Global CO_2 emissions. In Boden, T. A., Danciruk, P. and Farrell, M. P., eds., *Trends '90: A Compendium of Data on Global Change*. Oak Ridge National Laboratory/CDIAC-36: 92 p.
- Martinelli, L. A., Devol, A. H., Victoria, R. L. and Rickey, J. E. 1991 Stable carbon isotope variation in C_3 and C_4 plants along the Amazon River. *Nature* 353: 57–59.
- O'Leary, M. H. 1988 Carbon isotopes in photosynthesis. *Bioscience* 38: 328–336.
- Rominger, J. 1962 Taxonomy of *Setaria* (Gramineae) in North America. *Illinois Biological Monograph* 29. Urbana, University of Illinois Press: 1–132.
- Slota, P. J., Jull, A. J. T., Linick, T. and Toolin, L. J. 1987 Preparation of small samples for ^{14}C accelerator targets by catalytic reduction of CO . *Radiocarbon* 29(2): 303–306.
- Tiezen, L. L. and Boutton, T. W. 1989 Stable carbon isotopes in terrestrial ecosystem research. In Rundel, P. W., Ehleringer, J. R. and Nagy, K. A., eds., *Stable Carbon Isotopes in Ecological Research, Ecological Studies Series*. New York, Springer Verlag: 167–195.
- Van Devender, T. R., Martin, P. S., Thompson, R. S., Jull, A. J. T., Long, A., Toolin, L. J. and Donahue, D. J. 1985 Fossil packrat middens and the tandem accelerator mass spectrometer. *Nature* 317: 610–613.
- Van Devender, T. R. and Toolin, L. J. 1983 Late Quaternary vegetation of the San Andres mountains, Sierra County, New Mexico. In Eidenbach, P. L., ed., *Prehistory of Rhodes Canyon, New Mexico*. Tularosa, New Mexico, Human Systems Research: 33–54.
- Van Devender, T. R., Toolin, L. J. and Burgess, T. L. 1990 The ecology and paleoecology of grasses in selected Sonoran Desert plant communities. In Betancourt, J. L., Van Devender, T. R. and Martin, P. S., eds., *Packrat Middens – The Last 40,000 Years of Biotic Change*. Tucson, The University of Arizona Press: 326–349.