

ISOTOPIC ANALYSIS OF GROUNDWATER AND CARBONATE SYSTEM IN THE SURDULICA GEOTHERMAL AQUIFER¹

MUNEVERA HADŽIŠEHOVIĆ, NADA MILJEVIĆ, VOJISLAVA ŠIPKA
DUŠAN GOLOBOČANIN and RADULE POPOVIĆ²

Boris Kidrič Institute of Nuclear Sciences, Vinča, P. O. Box 522, 11001 Belgrade, Yugoslavia

ABSTRACT. We present here results of our investigation of the isotopic chemical composition of groundwater and carbonates in the Surdulica geothermal aquifer, Serbia. We considered the effects of carbonate dissolution and measured ¹³C, ¹⁴C, D, ¹⁸O, ³H, field pH, temperature, Na⁺, Ca²⁺, Mg²⁺, HCO₃⁻ and other aqueous species from 30 springs and boreholes. Geothermal waters are supersaturated with calcite. Carbon isotope compositions vary with carbonate mineral dissolution. The δD and δ¹⁸O of groundwater samples fit the meteoric water line, and indicate that groundwater is recharged mainly from higher altitudes and the cold season. Different groundwater residence times point out two mechanisms for their formation: fissure flow for young waters and standard diffusion processes for old ones.

INTRODUCTION

The Surdulica geothermal system is a major segment of the Serbia-Macedonian massif. The ~250 km² area is a potential source of geothermal energy and drinking water (~250 liter s⁻¹, 45–126°C). The largest hot springs are concentrated near the Bujanovac and Vranjska Spas (ca. 30 springs); this group of springs, which seems to discharge at locations of the main aquifer system reaching ca. 3000 m deep, has not yet been described. Also, relations between groundwater flow patterns and the physical structure of the aquifer have been poorly understood. Deep drilling has been limited to a few exploratory boreholes (up to 2000 m). Faults, particularly along the periphery of the South Morava valley, appear to be significant factors controlling the discharge from many of the parts of the Surdulica-Vranje-Bujanovac area. Several researchers have investigated geophysical and hydrological conditions (Žujović 1893; Dimitrijević *et al.* 1965; Babović *et al.* 1970; Anderson 1985; Popović 1990). Since 1986, we (Milovanović *et al.* 1989) have investigated the isotopic chemistry of the aquifer carbonate and groundwater circulation systems, range of residence times and the origin of the geothermal water. This study surveys carbon isotopic compositions of the system and groundwater ¹⁴C ages and variations, and attempts to understand the processes upon which the isotopic values and their spatial distributions are based.

BASIC CHARACTERISTICS OF THE SURDULICA AQUIFER

The Vranje-Surdulica geothermal system is a narrow catchment area of the South Morava valley between Grdelica and Bujanovac (Fig. 1) in the Besna Kobila, Kukavica and Kozjak Mountains. The system is situated mostly in the Morava massif, consisting of Paleozoic gneiss, mica schists, amphibolites and greenstone (metagabbros, metadiabases, calc-schists, quartz and other chlorite schists) complexes. The Tertiary Surdulica system contains granitoids and dacite-andesite volcanics.

A large amount of CO₂ was introduced to the system during Tertiary magmatism, which could explain the strong carbonization of older rocks. This is apparent in metamorphic rocks from Boreholes 10 and 10.1 (drilled to 1603 m). Abundant sulfates in thermal waters (2–5 mmol liter⁻¹) and little sulfide suggest an anaerobic environment; thus, anaerobic decay with sulfate reduction

¹This paper was presented at the 14th International ¹⁴C Conference, 20–24 May 1991, Tucson, Arizona.

²Geoinstitute, Rovinjska 12, 11000 Belgrade, Yugoslavia




Fig. 1. Geological map of the study area with sampling points: 1. Tertiary and Quaternary sediments; 2. Tertiary volcanic rocks; 3. Surdulica granitoid; 4. Bujanovac granitoid; 5. Kukavica granitoid; 6. green complex; 7. gneiss and mica schist (Morava massif); 8. gneiss (Pelagonia-Rhodope massif); 9. contact of these two massifs; 10. contact of gneiss and the green complex in the Morava massif

as a source of HCO_3^- can be ignored. However, the possibility that some modern CO_2 in thermo-mineral waters of this system has a mantle origin should not be excluded.

The study region lies between moderate continental and Mediterranean climates, with mean annual air temperature around 10.5°C . Runoff from Besna Kobila (1923 m asl) drains into the Aegean and Black Seas. The average annual precipitation ranges from 915 mm in the high areas to 655 mm

in the lower elevations, where most of the precipitation occurs from October to March. Stream flow is also highest during these months.

The system discharges to wells and springs about 150 liter s⁻¹ of cold and nearly 100 liter s⁻¹ of thermal waters (45–126°C). Relatively impermeable layers in the divided aquifer formation direct the subsurface flow to large springs (up to 80 liter s⁻¹), which discharge along many faults. The configuration and geological structure of the aquifer and D, ¹⁸O, ¹³C, ¹⁴C isotopic data indicate that the Besna Kobilja Mountain is a major recharge area. Natural thermal springs appear along the foothills of a schist massif at about 400 m asl, mostly near the Vranjska and the Bujanovac Spas.

The regional geothermal gradient is 1°C/25 m (Anderson 1985) with a local high gradient of up to 1°C/8 m. A plot of temperature vs. depth (Fig. 2) shows that even higher temperatures are found at some loci (here 2, 3, 4, 10). From this gradient, the maximal depth of circulation is ca. 2500 m. We also estimated the depth from which hot water discharges, based on silica and cation-ratio geothermometers (Henley *et al.* 1985), and concluded that two temperature regimes exist: for the Bujanovac Spa (sampling point 30, Table 1), with <700 m and temperature of 100°C, and for the Vranjska Spa, with up to 4000 m depth and temperature up to 200°C.




Fig. 2. Relation of water temperature to depth of discharge (all points represent boreholes, see Table 1)

SAMPLING AND ANALYSES

From 1986 to 1990, we measured ¹³C, ¹⁴C, ³H, D and ¹⁸O of groundwater, CO₂ gas and rocks at 30 sites. To define modern isotope input parameters, we collected samples from soil CO₂ and vegetation, and determined pH values, alkalinity and major ion chemistry in water samples.

We analyzed the ¹³C and ¹⁸O content in the CO₂ gas prepared from samples according to IAEA (1980), using a Micromass 602 mass spectrometer. We obtained deuterium data by reducing water with zinc at 460°C (Coleman *et al.* 1982). We report stable isotope data in δ values (‰) related to the V-SMOW standard for D, ¹⁸O and for δ¹³C to PDB with 1 σ error of 2, 0.2 and 0.4‰, respectively. The δ¹³C value refers to the total dissolved carbon (TDC).

TABLE 1. Chemical Analyses of Waters from the Surdulica Aquifer (concentrations in mmol liter⁻¹ unless otherwise indicated)

Sample	Sampling point	Collection date	Alt* (m)	Temp. °C	pH	Na ⁺	K ⁺	Ca ²⁺	Mg ²⁺	Si ²⁺	Mn ²⁺	Zn ²⁺	HCO ₃ ⁻	CO ₃ ²⁻	SO ₄ ²⁻	F ⁻	Cl ⁻	SiO ₂ (mg/liter)
<i>Freshwater</i>																		
Spring	14	10/20/86	1475	8.0	7.7	0.25	0.02	0.44	0.2	0.02	0.02	0.01	0.8		0.05	0.01	0.10	3
Capt. spring**	21	10/10/86	1300	7.5	7.5	0.15	0.01	0.45	0.3	0.02	0.01	0.01	1.0		0.05		0.05	5
Fountain	22	07/14/88	912	13.1	7.7	0.23	0.02	0.60	0.3	0.02	0.02	0.01	1.3		0.10		0.10	
Capt. spring	5	07/12/88	700	17.6	6.8	0.24	0.05	0.60	0.4	0.01			1.2		0.10		0.05	3
River	6	12/28/88	687	4.4	7.3	0.13	0.02	0.60	0.2	0.01	0.01		1.0		0.10	0.01	0.10	9
River	28	10/01/87	430	6.3	7.6	0.46	0.05	0.74	0.4	0.02	0.01		1.4		0.20		0.20	5
<i>Transition and low-temperature thermal water</i>																		
Mine water, d [†] = 600	16	06/18/86	1310	11.2	7.3	0.20	0.01	0.30	0.2		0.01	0.01	0.5		0.10	0.02	0.10	7
B [†] , d = 112	25	07/12/88	1354	10.6	7.3	0.19	0.01	0.40	0.1		0.01	0.01	1.2		0.10	0.05	0.10	10
Tunnel, d = 300	27	09/14/88	693	20.5	7.6	0.90	0.10	1.20	0.7	0.02	0.01		1.8		0.80	0.20	0.10	20
Capt. thermal spring	29	06/05/86	393	23.0	8.4	12.10	0.10	0.10	0.1			0.005	10.6	0.58	0.60	0.07	0.50	22
Capt. thermal spring	29	05/25/89	393	22.8	8.4	11.80	0.10	0.10	0.1				10.2	0.39	0.60	0.10	0.70	15
Capt. thermal spring	9	06/18/86	430	17.1	7.0	5.60	0.20	1.20	0.1				5.3		0.90	0.10	1.10	14
Capt. thermal spring	8	05/25/89	415	16.8	7.5	7.70	0.50	0.80	0.5				5.3		3.10	0.10	1.50	25
<i>Hot thermal water</i>																		
Capt. thermal spring	1	05/05/86	425	78.2	7.6	12.50	0.30	0.40	0.2				7.2		4.00	0.60	1.40	42
B, d = 120	3	05/06/86	294	80.5	7.5	11.50	0.20	0.70	0.1				6.4		4.20	0.60	1.50	90
B, d = 520	10	01/25/88	-105	79.4	8.2	13.50	0.30	0.50	0.4				9.8		3.20	0.20	1.50	39
B, d = 1063	10	04/14/88	-648	111 [†]	8.3	10.70	0.60	0.40	0.2				8.0		3.00	0.20	1.30	30
B, d = 450	10.1	05/25/89	-33	67	7.5	12.80	0.40	0.30	0.2				8.1		3.10	0.50	1.70	88
B, d = 535	30	07/12/88	-125	43	7.1	58.90	1.60	1.20	0.5	0.02			58.7		2.00		1.60	41
B, d = 535	30	05/26/89	-125	42.2	7.1	57.30	1.50	1.02	0.6	0.02			56.5	0.38	1.70	0.13	1.70	51

*Alt = altitude of water horizon

**Capt. = captured

[†]d = depth (m)

[†]B = borehole

[†] = measured temperature at 1063 m depth = 126°C, flow = 27 liter s⁻¹

We measured ^{14}C activities using a Packard 3380 liquid scintillation spectrometer or gas proportional counting system (Srdoč, Breyer & Sliepčević 1971). ^{14}C values are expressed in percent modern carbon (pMC), with a measurement error of $\pm 3\text{--}25\%$ (for hot waters); we measured tritium activity after electrolytic enrichment (Hut 1986).

RESULTS AND DISCUSSION

Water Chemistry and Isotopic Composition of Groundwater

Surdulica groundwaters are of $\text{Na}(\text{Ca})\text{--HCO}_3(\text{SO}_4)$ type, with pH values ranging from 7.0–8.4 (Table 1). The partial pressure of CO_2 is within $10^{-0.3}$ to $10^{-2.9}$ atm (Table 2) and increases downgradient from cold to geothermal water (GTW). The cold springs and most low-temperature thermal waters are unsaturated with calcite. Ionic concentrations (especially those of Na^+ , HCO_3^- and SO_4^{2-}) and groundwater temperature increase with depth. This is associated with considerable carbonization and pyritization of older rocks.

A plot of δD vs. $\delta^{18}\text{O}$ of groundwater samples fits a local meteoric water line of $\delta\text{D} = (7.3 \pm 0.2) \delta^{18}\text{O} + (7 \pm 2)$. Similarity between isotopic content of precipitation and GTW demonstrate their

TABLE 2. Isotopic Analyses and Chemical Data of Studied Waters

Sample	Sampling point	Log P_{CO_2} * (atm)	SI_c **	$\delta^{13}\text{C}_{\text{PDB}}$ (‰)	^{14}C (pMC)	^3H (TU)	^3H , TU interval†
<i>Freshwater</i>							
Spring	14	-2.9	-1.0	-13.0	72	59	55–65
Capt. spring	21	-2.8	-1.2	-12.9	90	58	32–69
Fountain	22	-2.9	-0.7	-13.5	67	37	28–46
Capt. spring	5	-1.9	-1.0	-13.4	71.1	19	17–37
Capt. spring	5	-2.3	-1.4	-16.3	94.1	23	17–37
Capt. spring	11	-2.8	-0.2	-10.2	61.3	18	15–35
<i>Transition and low-temperature thermal water</i>							
Mine water	16	-2.9	-1.8	-12.7	39.7	14	11–45
B [‡] , d [§] = 212	25	-2.5	-1.3	-10.7	38.7	38	38
Tunnel water	27	-2.7	0.1	-13.8		7	7
Thermal spring	29	-2.7	0.6	-7.8	26.9	12	5–18
Thermal spring	29	-2.5	0.5	-8.2	21.1	5	5–18
Thermal spring	9	-1.9	-0.1	-12.2	77.5	13	4.5–16
Thermal spring	8	-2.0	-0.1	-3.1	16.4	<2	
<i>Hot thermal water</i>							
Thermal spring	1	-1.7	0.9	-6.6	4.5	<2	
B, d = 120	3	-1.7	1.0	-3.7	6.0	<1	
B, CO_2 gas	3			-4.5			
B, d = 520	10	-2.2	1.8	-4.8	5.3	<0.9	
B, d = 867	10	-2.5	2.4	-1.8	5.4	<1.4	
B, d = 450	10.1	-1.6	0.9	-2.4	4.8	<3	
B, d = 535	30	-0.3	1.3	-0.9	3.3	<1	
B, d = 535	30	-0.4	1.2	-0.4	6.9	<2	

*Log P_{CO_2} = $\log \frac{[\text{H}_2\text{CO}_3]}{K_{\text{CO}_2}}$; ** SI_c = $\log \frac{\text{ion activity product}}{\text{equilibrium constant}}$

†Study period – 1986–1990; ‡B = borehole; §d = depth (m)

meteoric origin. Distribution and seasonal variations of D and ^{18}O isotopes in the water indicate that GTW have been recharged predominantly from higher elevations (>500 m asl) and during colder seasons.

All of the water samples, except for a few old geothermal samples, contain tritium from 5 to 59 tritium units (TU; $1 \text{ TU} = 0.118 \text{ Bq l}^{-1}$) (Table 2); this large interval is a result of different drainage rates throughout the system due to variable rock porosities. In rivers and shallow groundwater (freshwater, Table 2), ^{14}C concentrations are between 61.3 and 94.1 pMC, whereas $\delta^{13}\text{C}$ values range from -16.3 to -10.2‰ (Table 2). Transition- and low-temperature thermal water from deeper portions of the aquifer have lower ^{14}C (16.4–77.5 pMC) and higher ^{13}C (-13.8 to -3.1‰). The samples from hot thermal springs and boreholes had the lowest ^{14}C (3.3–6.9 pMC) and the highest $\delta^{13}\text{C}$ values (-6.6 to -0.4‰). The $\delta^{13}\text{C}$ value of the CO_2 gas analyzed from Borehole 3 is -4.5‰ . The ^{13}C content of core samples from Boreholes 10, 10.1 and 26 (-8.4 to -1.1‰) are very similar to those in GTW from deep circulations. Samples taken from marble rocks ranged from $\delta^{13}\text{C} = +0.41$ to $+2.7\text{‰}$.

Isotopic composition of dissolved carbonates formed during water interaction with rock matrix is plotted vs. the calcite saturation (Fig. 3). P_{CO_2} and $\delta^{13}\text{C}$ generally increase with calcite saturation, while ^{14}C decreases. To determine conditions of calcite mineral dissolution, we calculated the relation between HCO_3^- and ^{13}C content of water for systems open and closed to a CO_2 gas reservoir (Fig. 4). Under open-system conditions (Fig. 4A), cold waters gather between calcite saturation curves corresponding to initial values of ^{13}C content of soil gas, $\delta^{13}\text{C}_g = -24$ to -20‰ at initial pH values, $\text{pH}_i = 5$ – 6 . The isotopic composition of GTW is out of this range. This could be explained by significant dissolution of rocks and isotopic exchange between carbonate species in the aquifer. In Figure 4, we give the saturation curves for the supposed systems, assuming their initial parameters, $\delta^{13}\text{C}_g = -13\text{‰}$ (mean value for freshwater), -8‰ (low thermal water, point 29) and $\text{pH}_i = 5$ – 7 . Similar analysis derived for closed-system conditions (Fig. 4B) shows that cold groundwater recharge should have $\text{pH}_i = 5$ – 6 and $\delta^{13}\text{C}_g = -27$ to -24‰ , and P_{CO_2} s may range from $10^{-1.5}$ to $10^{-2.5}$. The ^{13}C content of the dissolved carbonate from rocks, $\delta^{13}\text{C}_r$, could be between -3.4‰ (mean for Borehole 10, Table 3) and $+1.4$ (mean for marble). Taking into account $\delta^{13}\text{C}_g = -13$ and -8‰ , and $\text{pH}_i = 5$ – 6 , as initial parameters, we show that most measured ^{13}C values for warm waters are still out of the range of the saturation curves. These values are greater than for open-system conditions.

Age of Groundwater

As shown in Table 2, freshwaters contain considerable tritium, and to a certain degree, reduced concentration of ^{14}C . Thus, for an estimate of mean residence time (MRT), we used the finite-state, mixing-cells model (Yurtsever 1983). For most samples, this model gives 2 or 3 solutions for MRT and an adjustment factor Q (a parameter for non-decay ^{14}C loss induced by chemical interaction with carbonate rock). There may be two reasons for this: 1) the sampling was performed at the end of tritium-response curves of the groundwater system (1986–1990); 2) water takes fast and slow paths through the system. The analyzed water is a mixture of these two components. This assumption is validated by the fact that most of the studied waters cluster around $Q = 0.7$ – 0.75 , which corresponds to a MRT of 100–700 yr (Table 4). On the other hand, tritium was detected in the same samples, which leads to 10–80-yr-old water, *i.e.*, young water. If waters with two very different MRTs exist simultaneously, we must assume two different mechanisms for their formation. In the case of fast water, it passes through fissures that conform to the geological structure of the aquifer, whereas a standard diffusion process is responsible for old water. We estimate the MRT at less than 80 yr for the fissured process and to 200–500 yr for the older waters. Isotopic




Fig. 3. Dependence of water ^{14}C , ^{13}C and P_{CO_2} on the saturation index of calcite




Fig. 4. Relation between $\delta^{13}\text{C}$ and HCO_3^- for open and closed systems for various initial conditions (10, 30°C).

A. Open-system solution paths:

(1) $\delta^{13}\text{C}_s = -24$; (2) $\delta^{13}\text{C}_s = -20$ ‰; (3) $\delta^{13}\text{C}_s = -13$ ‰; (4) $\delta^{13}\text{C}_s = -8$ ‰

B. Closed-system solution paths:

(1) $\delta^{13}\text{C}_s = -27$ ‰, $\delta^{13}\text{C}_r = -3.4$ ‰, $\text{pH}_i = 5$; (2) $\delta^{13}\text{C}_s = -27$ ‰, $\delta^{13}\text{C}_r = +1.4$ ‰, $\text{pH}_i = 5$; (3) $\delta^{13}\text{C}_s = -24$ ‰, $\delta^{13}\text{C}_r = -3.4$ ‰, $\text{pH}_i = 5$; (4) $\delta^{13}\text{C}_s = -24$ ‰, $\delta^{13}\text{C}_r = +1.4$ ‰, $\text{pH}_i = 6$; (5) $\delta^{13}\text{C}_s = -13$ ‰, $\delta^{13}\text{C}_r = -3.4$ ‰, $\text{pH}_i = 5$; (6) $\delta^{13}\text{C}_s = -13$ ‰, $\delta^{13}\text{C}_r = +1.4$ ‰, $\text{pH}_i = 5$; (7) $\delta^{13}\text{C}_s = -13$ ‰, $\delta^{13}\text{C}_r = +1.4$ ‰, $\text{pH}_i = 6$; (8) $\delta^{13}\text{C}_s = -8$ ‰, $\delta^{13}\text{C}_r = -3.4$ ‰, $\text{pH}_i = 6$; (9) $\delta^{13}\text{C}_s = -8$ ‰, $\delta^{13}\text{C}_r = +1.4$ ‰, $\text{pH}_i = 5$; (10) $\delta^{13}\text{C}_s = -8$ ‰, $\delta^{13}\text{C}_r = +1.4$ ‰, $\text{pH}_i = 6$. Subscripts, s and r, stand for soil and rock, respectively. - - - = calcite saturation curves; --- = calcite solution paths for various P_{CO_2} atm, Δ sample.

analyses indicate a complex carbon origin (biogenic CO_2 from the soil zone, carbonate species from carbonate mineral dissolution, CO_2 from the mantle); now, the question of age for these waters is open. Tritium-free waters suggest ages older than 100 yr. The low ^{14}C content (ca. 5 pMC) would infer waters ca. 20,000 yr old (hydrochemical age, Fontes 1979). Measured TDIC $\delta^{13}\text{C}$, > -6.6 ‰, indicates dilution by carbon of unknown origin. Other models (Fontes 1979) give unrealistic, even negative ages, and are not applicable to the Surdulica system. We believe that these waters are considerably younger than 20,000 yr because of high bicarbonate concentrations (up to 60 mmol

liter⁻¹) in the basin, which cannot be explained only by calcite and dolomite dissolution ($m_{Ca} + m_{Mg} = 0.5\text{--}1.5 \text{ mmol liter}^{-1}$).

CONCLUSION

Geothermal waters are of meteoric origin, formed in recharge areas at higher elevations and during colder seasons. The results of tritium and ¹⁴C measurements indicate fast (fissure) and slow (intergranular) flow paths through the system. Water composition is characterized by bicarbonate sodium (calcium) sulfates. The concentration of the main cations, anions and silica increases by a factor of almost ten, descending from the recharge area to the deep waters. Calcite saturation increases with depth, and GTWs are almost supersaturated.

It appears that cold water recharges under closed-system conditions, with initial parameters, $\delta^{13}C_s = -27\pm 3\text{‰}$, $pH_i = 5\text{--}6$, reaching a value of -13‰ for $\delta^{13}C_s$ at the discharge. The concentrations of ¹³C in GTW are significantly higher (up to 0‰), which indicate processes of intensive interaction between water and rocks, isotopic exchange, and a possible interior CO₂ source. The MRT for mixed waters calculated by finite-state mixing models is less than 80 yr for the fast component, and 200–500 yr for the older one. The age of geothermal waters based on isotopic analyses ranged from 100 to 20,000 yr.

ACKNOWLEDGMENTS

Partial financial support of this study was provided by the IAEA (Research Contract No. 4852/R1/R2) and Scientific Funds of Serbia.

REFERENCES

- Anderson, E. 1985 *Review of Vranjska Banja Geothermal Prospect, Scientific and Engineering Aspects*. Wellington, New Zealand, GENZ-Geothermal Energy New Zealand, Ltd.: 12 p.
- Babović, M., Vukanović, M., Krajačić, V., Rakić, M. and Dimitrijević, M. 1970 *Basic Geological Map, Chart Vranje 1:1000 000*. Geozavod, Federal Geological Institute, Belgrade.
- Coleman, M. L., Shepherd, T. J., Durham, J. J., Rouse, J. B. and Moore, G. R. 1982 Reduction of water with zinc for hydrogen isotope analysis. *Analytical Chemistry* 54: 993–995.
- Dimitrijević, M., Petrović, B., Cikin, M., Mozina, A., Vikanović, M. and Karajičić, L. J. 1965 *Interpreter of the Basic Geological Map of the SFRJ, Chart Leskovac 1:100000*. Federal Geological Institute, Belgrade.
- Fontes, J. Ch. and Garnier, J. M. 1979 Determination of the initial ¹⁴C activity of the total dissolved carbon: A review of the existing models and a new approach. *Water Resources Research* 15: 399–413.
- Henley, R. W., Truesdell, A. H., Barton, P. B., Whitney, J. A. and Robertson, J. M., eds., 1985 Fluid-mineral equilibrium in hydrothermal systems. *Review of Economic Geology* 1: 31–43.
- Hut, G. 1986 *Intercomparison of Low-Level Tritium Measurements in Water*. Vienna, IAEA.
- IAEA (ms.) 1980 Preparation of carbon dioxide for C-14 measurements. *Technical Procedure Note*: 25 p.
- Milovanović, B., Stanković, S., Komatina, M., Hadžišehović, M., Zupančić, M., Miljević, N., Stepić, R. and Obelić, B. 1989 Isotopic investigation of the Surdulica geothermal system. In Long, A. and Kra, R. S., eds., *Proceeding of the 13th International ¹⁴C Conference, Radiocarbon* 31(3): 893–901.
- Popović, R. 1990 The Serbian-Macedonian mass – yes or no? *Records of the Serbia Geological Society*, 1990, Belgrade.
- Srdoč, D., Breyer, B. and Sliepčević, A. 1971 Rudjer Bošković Institute radiocarbon measurements I. *Radiocarbon* 13(1): 35–140.
- Yurtsever, Y. 1983 Models for tracer data analysis. In IAEA/UNESCO Guidebook in Nuclear Techniques in Hydrology. IAEA *Technical Reports Series* 91: 381–400.
- Žujović, J. M. 1893 Thermal springs of Vranjska Spa. In *Geology of Serbia Vol. II-Granite*. Belgrade, Serbian Royal Academy: 54–73.