

## NANCY NATURAL RADIOCARBON MEASUREMENTS V

R COPPENS, B GUILLET, R JAEGY and P RICHARD

Laboratoire de Radiogéologie. Ecole Nationale Supérieure de Géologie  
BP 452, 54001 NANCY, France

The following list includes some measurements made during 1975-1978 since our last list (R, 1978, v 20, p 62-67) in the National Radiocarbon Laboratory of the Ecole Nationale Supérieure de Géologie Appliquée et de Prospection Minière (ENSG) de Nancy.

Laboratory procedures and techniques are as reported in Hassko *et al* (1974) where the sample is synthesized to benzene. The only difference is in the counting equipment. We now use a Packard Tricarb 3305 liquid scintillation spectrometer where samples are counted for 24 or 48 hours. Radiocarbon ages are calculated using  $^{14}\text{C}$  half-life of 5568 years and 95% activity of NBS oxalic acid is used as modern standard. Anthracite coal and Merck commercial benzene are used for the dead carbon run.

Counting errors are expressed at  $1\sigma$  confidence level. AD/BC\* dates are corrected using  $^{14}\text{C}$  half-life of 5730 years and according to the Masca correction curve (Ralph *et al*, 1973).

Description and comments are generally based on information supplied by those who submitted the samples.

### SAMPLE DESCRIPTIONS

#### I. ARCHAEOLOGIC AND HISTORIC SAMPLES

##### A. Peru

The following samples from different prehistoric villages in Peru are a continuation of the series previously described (R, 1978, v 20, p 62-67). Coll and subm by Frederic Engel, Dir Mission archaeol française au Pérou and Inst Antropol y Agric Precolombina, Univ Nac Agraria, Lima, Peru.

##### Chillon program series

*Sites with Neolithic IV or V pottery*

##### Ny-367. Chillon valley, V 3563

**2060 ± 70**  
**2080\***

Plant material from Village 11 b X-131, Chacra alta Layer 100, alt 15m (11° 56' 00" S, 77° 05' 00" W). *Comment* (FE): dates a post-Chavin, pre-early Lima settlement in the Chillon valley. Cf Willey's white on red coll 1973 by F Engel and B Ojeda.

##### Southern deserts series

*1. Preagricultural sites*

##### Pampa Colorado Fog oasis series

##### Ny-383. Fog oasis V 3653

**10,200 ± 140**

Charcoal from Encampment 17 c VIII-280A, Layer 400, alt 30 to 50m (16° 31' 30" S, 72° 51' 30" W). *Comment* (FE): could be one of oldest

Holocene settlements known in deep south of Peru. Located in fog oasis extending to bluff overhanging ocean. Estimated age: 9000 BP. Coll 1975 by F Engel.

**Ny-381. Fog oasis V 3651** **9310 ± 120**

Shells from Encampment 18 b IX-5, Las Higueras, alt 10m, (16° 14' 20" S, 71° 34' 57" W). *Comment* (FE): will date 1 of early Holocene encampments found on bluff overhanging ocean. Coll 1975 by F Engel.

**Ny-387. Fog oasis V 3657** **8490 ± 150**

Shells from Encampment 17 c VIII-2015, El Carrizal, alt 200m (16° 29' S, 72° 55' W, central point). *Comment* (FE): will date early settlement in Pampa Colorado fog oasis. Coll 1975 by F Engel.

**Ny-386. Fog oasis, V 3656** **6620 ± 100**

Shell from Encampment 19 a V-300, Punta Icu, Layer 200, alt 25m (17° 49' 00" S, 71° 08' 20" W). *Comment* (FE): will help date encampments found on bluff overhanging ocean. Coll 1975 by F Engel.

2. Pottery-yielding settlements, Neolithic

**Ny-380. Fog oasis, V 3650**

**3510 ± 80**  
**3990\***

Charcoal from very large, La Buitrera village, 19 a II-100, from foot of fog oasis in marshy area. Alt 150m (17° 45' S, 71° 10' W, central area). *Comment* (FE): La Buitrera was occupied at least twice, first during pre-forming days. Sample is from Layer 200, typologically showing early occupation by settlers using pottery. Coll 1975 by F Engel.

3. Pottery yielding settlements, alloyed metal period

**Ny-384. Fog oasis, V 3654**

**1100 ± 100**  
**AD 890\***

Bones from Village 19 b VII-105, Loma el Platanillo, Layer 100 (18° 02' 25" S, 76° 46' 00" W) alt 270m. *Comment* (FE): correct date for Tiahuaneco influenced village would read 1200 to 900 BP, AD 750 to 1050.

**Ny-358. Fog oasis**

**870 ± 80**

Village at mouth of Ica R; raft made of reeds was found hidden in sand dune. Some reeds were used for dating. Alt 20m (14° 52' 00" S, 75° 33' 34" W). *Comment* (FE): raft was decorated with fragments of shells (*Spondylus* sp), typical Tiahuanaco cultural trait in Peru (only Pre-colombian raft known in South America, now in Mus Nat Agrarian Univ, Lima). Coll 1974 by F Engel.

**Ny-385. Fog oasis, V 3655**

**650 ± 70**  
**AD 1300\***

Bones from Encampment 17 c VII-715, Palo Parado, on bluff. Alt 75 to 100m. Stone bldg (16° 29' 00" S, 73° 03' 00" W central point). *Comment* (FE): could be late Pre-Incaic. Coll 1975 by F Engel.

**Central coast program series***1. Preceramic horizons*

- 4885 ± 90**  
**5650\***
- Ny-414. Fog oasis, V 3704**  
Charcoal from Village 13 a II-90, Quilmana I, Layer 200. Alt 440m (12° 53' 58", 76° 26' 03"). *Comment* (FE): will help date early farming settlements on central coast. Coll 1976 by B Ojeda.
- 4250 ± 70**
- Ny-415. Fog oasis, V 3708**  
Charcoal from Village 12 b XI-500, Omaniso I, Layer 100, alt 440m (12° 57' 20" S, 76° 27' 54"). *Comment* (FE): same as Ny-414. Coll 1976 by B Ojeda.
- 3580 ± 80**  
**5530\***
- Ny-418. Fog oasis, V 3757**  
Charcoal from Village 13 a II-45, Carretilla, Layer 200, alt 150m (12° 58' 26" S, 76° 27' 56" W). *Comment* (FE): same as Ny-414. Coll 1976 by B Ojeda.
- 3550 ± 80**  
**5500\***
- Ny-416. Fog oasis, V 3709**  
Charcoal from Village 13 a II-70, Quilmana II, Layer 100, alt 350m (12° 55' 30" S, 76° 26' 45" W). *Comment* (FE): same as Ny-414. Coll 1976 by B Ojeda.
- 3430 ± 80**  
**5380\***
- Ny-417. Fog oasis, V 3750**  
Charcoal from Village 13 a II-10, Quilmana III, Layer 100, alt 310m (12° 56' 50" S, 76° 26' 42" W). *Comment* (FE): same as Ny-414. Coll 1976 by B Ojeda.
- 3240 ± 80**
- Ny-419. Fog oasis, V 3760**  
Charcoal from Village 13 a II-20, Cerro Grande, Layer 200, alt 330m (12° 56' 55" S, 76° 26' 52" W). *Comment* (FE): same as Ny-414. Coll 1976 by B Ojeda.
- 4740 ± 100**  
**5535\***
- Ny-392. Fog oasis, V 3700**  
Carbonized wood fragment from village 12 b IV-490 (12° 20' 30" S, 76° 43" W). Los Icasos, alt 280m. *Comment* (FE): estimated age: 5000 BP?. Could be a coastal Neolithic I village. Coll 1975 by F Engel.
- 3965 ± 100**  
**4520\***
- Ny-393. Fog oasis, V 3702**  
Structure seems to be related to late preceramic or Neolithic III without maize. It seems to be a common structure which is very rare at that time. Estimated age: 3500 BP. Coll 1975 by F Engel.

## 2. Alloyed metal periods

**Ny-477. V 6170 A** **680 ± 75**  
**AD 1275\***

Fragment of beam supporting roof of stone house, Village 12 b II-15, Orcocoto, Lurin Valley, alt 2070m (12° 06' 40" S, 76° 27' 00" W). *Comment* (FE): will help date "Cuculi period" villages of 13th-14th centuries AD. Coll 1977 by F Engel.

*B. Ivory Coast*

Samples were dated to study history of Kingdom of Kong, commercial area in savannah between forest and Sahel, Sahara and North Africa.

**Kong (Ferkessidougou Dept) series**

(9° 10' N, 4° 30' W)

**Ny-395. Kong, Site 5000** **350 ± 120**  
**AD 1440\***

Estimated age: AD 1400 to 1800. Coll 1975 by V Diabaté Tiégré.

**Ny-405. Kong, Site 6000** **195 ± 85**  
**AD 1650\***

Charcoal coll at 210cm below surface. Estimated age: AD 1400 to 1800. Coll 1976 by V Diabaté Tiégré.

**Ny-406. Kong, Site 7000** **185 ± 90**  
**AD 1650\***

Charcoal coll at 130cm below surface. Estimated age: AD 1400 to 1800. Coll 1976 by V Diabaté Tiégré.

**Ny-409. Kong, Mosquée de Sitafa** **410 ± 90**  
**AD 1440\***

Charcoal coll 150cm below surface. Estimated age: AD 1400 to 1800. Coll 1976 by V Diabaté Tiégré.

**Ny-410. Kong, Grande Mosquée** **335 ± 85**  
**AD 1490\***

Charcoal coll 60cm below surface. Estimated age: AD 1400 to 1800. Coll 1976 by V Diabaté Tiégré.

**Ny-404. Labiné, 2km SE Kong, Site 1000** **160 ± 65**  
**AD 1760\***

Charcoal coll 150cm below surface. Estimated age: AD 1300 to 1700. Coll 1976 by V Diabaté Tiégré.

**Ny-407. Ténéguéla, 9km Sud Kong** **340 ± 90**  
**AD 1470\***

Charcoal coll 180cm below surface. Estimated age: AD 1400 to 1800. Coll 1976 by V Diabaté Tiégré.

**Ny-399. Ténéguéla, 9km Sud Kong** **2290 ± 310**  
**130 BC\***

Humus 15 to 20cm. Coll 80cm below surface. Coll 1975 by J Polet.

**Sud Aby lagoon series**

Samples were dated to establish historic population maps of SE Ivory Coast.

- |  |                  |
|--|------------------|
|  | <b>125 ± 85</b>  |
| <b>Ny-401. Assoco—Mossobaha Island</b>   | <b>AD 1740*</b>  |
| Wood fragments from piles (3° 30' W, 5° 20' N). Coll 75cm below surface. Coll 1975 by J Polet. |                  |
|  | <b>2250 ± 95</b> |
| <b>Ny-408. Nyamva Island</b>   | <b>410 BC*</b>   |
| Charcoal under dense forest (10cm humus). Coll 1975 by J Polet 130cm below surface.            |                  |

*C. Afghanistan***Shortugai series**

Charcoal from hearths and ash layers. Ash with burned soils (marl) at different stratigraphic levels, coll from Protohistoric site of Shortugai (NE Afghanistan). Project was undertaken to study relations between Indus Valley and Central Asian civilizations in Bronze age period (Francfort & Pottier, 1978). Coll by H P Francfort and subm by J C Gardin.

- |   |                   |
|---|-------------------|
| <b>Ny-421. SHB 76, late Period III, Level 4</b>                                 | <b>3535 ± 165</b> |
| Estimated age: ca 3450 BP.  | <b>4000*</b>      |
| <b>Ny-424. SHB 76, Level 3, early Period III</b>                                | <b>3180 ± 335</b> |
| Estimated age: ca 3650 BP. Sample was very small. Benzene obtained was minimal. | <b>3480*</b>      |
| <b>Ny-427. SHB 76, Level 5, early Period III</b>                                | <b>3710 ± 100</b> |
| Estimated age: ca 3650 BP.  | <b>4110*</b>      |
| <b>Ny-428. SHB 76, Level 5, early Period III</b>                                | <b>4190 ± 125</b> |
| Estimated age: ca 3650 BP.  | <b>4875*</b>      |
| <b>Ny-422. SHB 76, Level 2, Period II</b>                                       | <b>3050 ± 250</b> |
| Estimated age: ca 3750 BP.  | <b>3330*</b>      |
| <b>Ny-429. SHB 76, Level 3, early Period II</b>                                 | <b>4375 ± 160</b> |
| Estimated age: ca 3850 BP.  | <b>5106*</b>      |
| <b>Ny-430. SHB 76, Level 1, late Period I</b>                                   | <b>4075 ± 95</b>  |
| Estimated age: ca 3950 BP.  | <b>4710*</b>      |
| <b>Ny-425. SHB 76, Level 1, early Period I</b>                                  | <b>4040 ± 100</b> |
| Estimated age: ca 4050 BP.  | <b>4605*</b>      |

*General Comment:* Masca corrected dates run generally several centuries older than tentative chronology based on ceramic typology. Discrepancy perhaps is due to contamination of samples by older mineralogic carbon.

*D. France*

**Gordes series, Vaucluse, France**

Wood fragments from cellar, estimated age: 15th or 16th century. Coll and subm by J L Morand.

**Ny-473.** **380 ± 50**  
**AD 1450\***  
Wood from an ancient oil mill.

**Ny-541.** **200 ± 90**  
**AD 1644\***  
Wood from beam in wall of cellar.

**Ny-542.** **255 ± 95**  
**AD 1580\***  
Well-preserved wood from wooden chimney funnel in wall of cellar.

*General Comment (RC):* beam and chimney funnel seem to be same age, ca AD 1600, late 16th or early 17th century, and suggest redistribution of cellar. Oil mill seems older, 15th century. Either is truly older or corresponds to re-use of old wood in new structure.

**Ny-485. Limoges, Haute Vienne, rue du Clos Adrien** **1980 ± 80**  
**30 BC\***

Charcoal from Gallo Roman well with pottery and tegulae, of 2nd century. Coll 1977 by J P Loustaud and subm by J M Desbordes. Estimated age: 2nd century. *Comment:* date appears older than expected based on historic correlations, but wood may be really older.

**Ny-498. Roman church of Meymac, Corrèze** **770 ± 75**  
**AD 1270\***

Charcoal from glass-founder furnace in chancel of church. Sample No. 2 c/III. Durst hole furnace No. 2. Coll and subm by J M Desbordes. Estimated age: 12th century.

**Ny-497. Roman church of Meymac, Corrèze** **265 ± 75**  
**AD 1570\***

Charcoal from site below choir of church in Roman surroundings. Coll 1978 and subm by J M Desbordes. Estimated age: 12th century.

**Ny-499. Tarnac, Corrèze** **2020 ± 70**  
**55 BC\***

Finely dispersed charcoal from base of funeral mound embedded in brown and tamped soil. Coll 1977 and subm by J M Desbordes. Estimated age: 12th century Iron age. *Comment:* date is younger than expected. Sample does not belong to Prehistoric occupation or may be contaminated by rootlets or humic acids.

**Ny-486. Jarnages, Creuse, lieu-dit Martinfort** **2440 ± 85**  
**490 BC\***  
Carbonized wood from early Gallo-Roman site with pottery and tiles. Coll 1977 by M Bord and subm by J M Desbordes. Estimated age: 1st century.

**Ny-489. Razes, Haute-Vienne, lieu-dit Augères** **660 ± 80**  
**AD 1305\***  
Carbonized wood from fill of underground gallery with post Gallo-Roman pottery. Coll 1977 by R Saumande and subm by J M Desbordes. Estimated age: post-Gallo-Roman period.

**Ny-488. Saint Pardoux, Haute Vienne, lieu-dit La Ribière** **600 ± 90**  
**AD 1380\***  
Carbonized wood from fill of underground gallery with post-Gallo-Roman pottery. Coll 1977 by R Saumande and subm by J M Desbordes. Estimated age: post-Gallo-Roman period.

**Ny-487. Cloister of Obazine, Corrèze** **960 ± 65**  
**AD 1020\***  
Wood fragment from stick (bishop's crook). Coll 1977 by B Barrière and subm by J M Desbordes. Estimated age: 13th century. Obazine abbey was founded AD 1130. *Comment:* date older than expected. Crook may have been transmitted by inheritance.

**Jabreilles-les-Bordes series, Haute-Vienne, lieu-dit Le Chatelard**

Study of ditch fill at foot of Historic fence (late Tene III or early Gallo-Roman period). Archaeal study made to determine absolute age and influence of human activities on environment.

**Ny-479. CH 77—1 and 1 bis** **820 ± 80**  
**AD 1165\***  
Charcoal from excavation at back of fields under cultivation. Sample may date early ploughing. Coll by B Valadas and subm by J M Desbordes. Estimated age: 12th century. *Comment:* <sup>14</sup>C date confirms interpretation above.

**Ny-482. CH 77—6** **940 ± 135**  
**AD 1030\***  
Charcoal from fill of ditch. Date may indicate surrender of ramparts and beginning of fill of ditch. Coll 1977 and subm by J M Desbordes. Estimated age: probably 10th century. *Comment:* <sup>14</sup>C date agrees with expected age.

**Ny-480. CH 77—3 and 3 bis** **670 ± 100**  
**AD 1275\***  
Charcoal from fill of ditch. Coll 1977 and subm by J M Desbordes. No age expected.

**Modern****Ny-481. CH 77—5****<180**

Charcoal from top of fill of ditch. Dates end of fill of ditch. Coll 1977 and subm by J M Desbordes. Estimated age: modern. <sup>14</sup>C date agrees with expected age.

*E. French West Indies***Ny-500. Marie Galante Island****1515 ± 85****AD 470\***

(61° 17' W, 15° 50' N). Charcoal from Taliseronde site. Coll and subm by D Emond.

**Ny-478. La Martinique, Fond Brulé site****1600 ± 250****AD 390\***

Charcoal from hearth, depth 130cm. Coll 1977 by M Mattoni. Site is between La Salle site (S), dated by Yale (Y-1116: 1770 ± 80 BP, R, 1963, v 5, p 336). Vive site (N), dated by FSU, 1730 ± 100 BP. Masca corrected dates are AD 190 ± 100\* and AD 235 ± 125\*. Our sample was of poor quality, very small (0.4173g C). Benzene obtained was minimal, yielding large statistical error, but result seems archeologically acceptable.

## II. GEOLOGIC SAMPLES

**Etang de Batéguier, Ile Ste Marguerite series, Alpes Maritimes France**

Shells from lacustrine sediments, Etang de Batéguier (5 gr 21, 9' W, 48 gr 35, 8' N). Cores taken in sediments of pool, depth 2.5m with level of numerous shells (*Bytinidae* sp, *Clausilidae* sp), undoubtedly killed by sudden disturbance of environment. Coll and subm 1975 by G Palausi.

**Ny-389. Ile Ste Marguerite****3920 ± 90****2530 BC\*****Ny-390. Ile Ste Marguerite****1410 ± 80****AD 580\*****Ny-391. Ile Ste Marguerite****1490 ± 80****AD 500\***

*General Comment* (GP): Ny-389, taken near shore, is probably not significant because of possible contamination by older shells. Two dates, Ny-390 and -391, agree statistically and help estimate rate of sedimentation for pool. Mean age: 1450 BP. Rate of sedimentation: 1.80 mm/yr.

## III. SOIL SAMPLES

*A. Andosols*

Andosols are soils developed on volcanic ashes. Most are mainly characterized by amorphous hydrous silica-alumina mineral such as allophanes, which can adsorb much organic matter.

Samples from Auvergne and Cantal coll 1974 by J M Hétiér and B Guillet; samples from Vivarais, by J Moinereau, E N S A Montpellier.

*Puy du Mercoeur, Auvergne*

(45° 42' 12" N, 2° 55' 12" E)

<b>Ny-558.</b>	<b>A<sub>12</sub> horizon, 8.8% C, 20 to 30cm</b>	<b>1680 ± 80</b>
<b>Ny-559.</b>	<b>B<sub>1</sub> horizon, 7.2% C, 40 to 50cm</b>	<b>3000 ± 80</b>
<b>Ny-560.</b>	<b>B<sub>2</sub> horizon, 3.4% C, 60 to 70cm</b>	<b>3700 ± 90</b>

*Comment:* Puy du Mercoeur is volcanic cone of basaltic cinders and scoria, S of Chaînes des Puys, 15 km SW Clermont-Ferrand, France. Volcanic eruption of Puy du Mercoeur is estimated at 4000 to 6000 yr, based on local stratigraphy (Hétier, 1975). Mean residence time of B<sub>2</sub> horizon organic matter probably expresses beginning of andosolization processes.

*Puy de Dôme, Auvergne*

(45° 45' 25" N, 2° 55' 14" E)

<b>Ny-561.</b>	<b>A<sub>1</sub> horizon, 14.2% C, 15 to 25cm</b>	<b>950 ± 70</b>
<b>Ny-562.</b>	<b>A<sub>1</sub> B<sub>1h</sub> horizon, 13.4% C, 30 to 40cm</b>	<b>2040 ± 80</b>
<b>Ny-563.</b>	<b>A<sub>1</sub> B<sub>2</sub> horizon, 10% C, 40 to 50cm</b>	<b>2590 ± 80</b>
<b>Ny-564.</b>	<b>II A<sub>1</sub> horizon, 1% C, 150 to 160cm</b>	<b>8210 ± 120</b>

*Comment:* mean residence time of organic matter of buried II A<sub>1</sub> horizon agrees well with dates of charcoals separated from similar buried soils (Brousse *et al*, 1969). Upper ando-podzolic soil is developed on domite material 150cm thick, forming of which began ca 8300 yr BP.

*Puy Mey, Auvergne*

(45° 42' 33" N, 2° 57' 57" E)

<b>Ny-565.</b>	<b>Buried III A<sub>1</sub> horizon, 6% C, 240 to 245cm</b>	<b>1280 ± 70</b>
<b>Ny-566.</b>	<b>Charcoal from buried III A<sub>1</sub> horizon</b>	<b>1510 ± 50</b>

*Comment:* Puy Mey is a small cone on Puy du Mercoeur and Puy de la Vache, covered by unweathered volcanic scoria. Two buried andosols are observed. Organic matter of deepest, Ny-565, had low residence time, suggesting that recovering by basaltic scoria is recent. Conclusion supported by age of charcoals, Ny-566, measured twice from 2 different samplings. If upper volcanic deposit was really caused by volcanic eruption, age of charcoals would date one of most recent volcanic events in Chaînes des Puys, ca AD 500.

*Montagne de Marlieux, Cantal*

(45° 13' N, 2° 35' 10" E)

<b>Ny-567.</b>	<b>A<sub>1</sub> horizon, 12.4% C, 15 to 25cm</b>	<b>650 ± 70</b>
<b>Ny-568.</b>	<b>B<sub>1</sub> horizon, 7.3% C, 40 to 50cm</b>	<b>3190 ± 90</b>

**Ny-569. B<sub>2</sub> horizon, 6% C, 60 to 75cm 3910 ± 90**

*Comment:* andosol developed on weathered material of early Pleistocene basalt, ankaramite, rejuvenated by glacial erosion. Mean residence time gradient of organic matter is very comparable to data obtained for Puy du Mercoeur andosol.

*Bois des Chabottes, Vivarais*

(44° 40' 23" N, 4° 21' 10" E)

**Ny-570. A<sub>1</sub> horizon, 36% C, 0 to 7cm undecomposed plant debris, size >200µm Modern****Ny-571. A<sub>1</sub> horizon, 17.4% C, 0 to 7cm humified material, <200µm 120 ± 50****Ny-572. B<sub>ir</sub> horizon, 8.4% C, 7 to 25cm 1340 ± 90****Ny-573. B<sub>h</sub> horizon, 10.2% C, 25 to 45cm 3500 ± 90**

*Comment:* this profile corresponds to evolution from andosol to podzolic soil by differentiation of ochrous B<sub>ir</sub> horizon (Moinereau, 1977).

*General Comment:* as proposed by Sharpenseel (1972), a regression curve may be established between mrt (Y value in yr) and depth of date samples (X value, in cm). Equation for andosols ( $Y = 60.9 X + 80$ ,  $r = 0.918$ ,  $n = 15$ ) suggests that gradient of mrt with depth is higher than in other soils, such as brownearths, vertisols, chernozems (Guillet, 1979). This is explained by great influence of allophanic and amorphous alumina material that biologically stabilizes adsorbed organic matter, as breakdown and turnover is delayed.

*B. Paleosols*

*Golbey, Vosges*

(48° 12' 47" N, 6° 26' 12" E). Surface of old fluvial terrace of Moselle was covered by loam, 1.40m thick, on which sol lessivé glossique (agric glossaqualf) is differentiated. Old surface corresponds to cryogenic humic soil with charcoals. Samples coll and subm 1976 by M Cailler, CPB Nancy.

**Ny-588. Humic paleosol, <500 µm, 1.8% C, 140 to 150cm 26,390 +830 -750****Ny-589. Humic acids from isolated charcoals 27,100 +2500 -1900****Ny-590. Charcoals >40,000**

*Comment:* humic paleosol was thought to have developed during Kesselt interstade of Würm glaciation (ca 27,000-30,000 BP). But treatments of isolated charcoals with NaOH 0.5 N and Cl H N solutions dated to >40,000 yr BP for residual charcoals. Paleosurface is older than expected and should date at Brorup interstade. Rejuvenation of paleosol humus

and of humic acid fraction of charcoals was probably caused by continuous percolation of organic matter and water through cracks of upper loamy soil.

*Le Bernardan, Haute Vienne*

(46° 19' 18" N, 1° 14' 41" E) Indurated iron pans in arenaceous granitic parent material coll and subm 1975 by J C Flageollet, Univ Nancy II.

**Ny-586. Le Bernardan I, 0.3% C, 30 to 40cm 2230 ± 80**

**Ny-587. Le Bernardan II, 0.1% C, 80 to 90cm 15,690 ± 430**

*Comment:* for deepest pan, older age was expected. Rejuvenation by root decomposition is possible.

*C. Soil organic matter fractionation*

Various fractions of A<sub>1</sub> and B<sub>1</sub> horizons of Puy du Mercoeur andosol were processed to determine different turnover rates of organic matter fractions. Soil organic matter extractions were performed with successive Na-pyrophosphate 0.1 M and NaOH N solutions. Fulvic acids are soluble when alkaline organic solutions are acidified (pH 1) causing humic acids to precipitate. Humine is non-extractable organic matter. Values in brackets report organic matter extracted as Carbon percent of total C of soil sample.

*Puy du Mercoeur, A<sub>1</sub> horizon*

**Ny-574. Fulvic acids 1, (15) 1330 ± 70**

Precipitated at pH 4.8: β humus.

**Ny-575. Fulvic acids 2, (13) 650 ± 70**

Fulvic acid fraction that does not precipitate at pH 4.8.

**Ny-576. Humic acids, (13) 1510 ± 70**

**Ny-577. Humine, (45) 1460 ± 70**

*Comment:* except for Fulvic acids 2, mean residence times of humus -C fractions are very similar.

*Puy du Mercoeur, B<sub>1</sub> horizon*

**Ny-578. Fulvic acids 1 (11) 2580 ± 80**

**Ny-579. Fulvic acids 2 (12.4) 2130 ± 80**

**Ny-580. Humic acids (17.5) 2760 ± 80**

**Ny-581. Humic acids: acid hydrolyzate (3.5) 1830 ± 80**

**Ny-582. Humic acids: hydrolyzate residue (14) 3270 ± 80**

**Ny-583. Humine fraction (55.5) 2870 ± 80**

**Ny-584. Humine: acid hydrolyzate (27) 2340 ± 80**

**Ny-585. Humine: hydrolyzate residue (29.5) 3180 ± 80**

*Comment:* as expected, hydrolyzate fractions of humic acids and humine are relatively more renewed than residual carbon fractions. This may be considered a constant characteristic of soil organic chemistry, resulting from higher turnover rate of aminopolysaccharides and other metabolic N-products, which may be liberated from humic polycondensates by 6 N HCl hydrolysis procedure for 16 hrs.

## REFERENCES

- Brousse, R, Delibrias, G, Labeyrie, J, and Rudel, A, 1969, *Eléments de chronologie des éruptions de la Chaîne des Puys: Soc Géol France Bull*, v 7, no. 11, p 770-793.
- Francfort, H P and Pottier, M H, 1978, *Sondage préliminaire sur l'établissement proto-historique harappéen et post-harappéen de Shortugai (Afghanistan du NE): Arts Asiatiques* 1978, v 34, p 29-79.
- Guillet, B, 1979, *Etude du renouvellement des matières organiques des sols par les radioisotopes (<sup>14</sup>C) en Pédologie, constituants et propriétés: Paris, Masson éd*, p 210-226.
- Hassko, B, Guillet B, Jaegy R, and Coppens R, 1974, *Nancy natural radiocarbon measurements III: Radiocarbon*, v 16, no. 1, p 118-130.
- Hétier, J M, 1975, *Formation et évolution des andosols en climat tempéré: Thesis, Univ Nancy*, 194 p.
- Moinereau, J, 1977, *Altération des roches, formation et évolution des sols sur basalte, sous climat tempéré humide (Velay-Vivaraï-Coïrons): Thesis, Univ Montpellier*, 139 p.
- Ralph, E K, Michael, H N, and Han, M C, 1973, *Radiocarbon dates and reality: Masca Newsletter*, v 9, no. 1.
- Sharpenseel H W, 1972, *Natural radiocarbon measurement of soil and organic matter fractions and on soil profiles of different pedogenesis: Internatl conf on radiocarbon dating, 8th Proc, Wellington, New Zealand, E*, p 3-11.