

UDINE RADIOCARBON LABORATORY DATE LIST III

FRANCO CALLIGARIS, STEFANO COLONNELLO
ADRIANO DEL FABBRO and SERENA PERRONE

Centro di Ricerca Applicata e Documentazione, Via Pradamano 2A, 33100 Udine, Italy

INTRODUCTION

This list includes some significant measurements carried out by CRAD Radiocarbon Laboratory from 1979 to 1985. The dates were obtained by liquid scintillation counting of benzene using the laboratory procedures described in previous lists (Barbina *et al* 1982: 214-216; Barbina, Calligaris & Ciuti 1979). Samples were pretreated according to generally applied methods, depending on type of material and particular archaeological and geologic features.

GEOLOGIC SAMPLES

Italy

The results reported here are part of a study of the paleogeography of the lagoon of Venice made by Istituto Studio della Dinamica delle Grandi Masse CNR, Venice, as previously described (Barbina *et al* 1982; Gatto 1980). The list contains dates of samples selected from the same core at different depths in the stratigraphic sequence, in order to verify the sedimentation rate. In all cases, samples were prepared from the most suitable materials, *ie*, peat or, if not available, mollusk shells.

Laguna di Venezia

Ca' Bianca series

Peat from lagoon of Venice Ca' Bianca (45°25'33"N, 12°21'18"E). Coll 1973 by P Da Roit, Lab Geol Appl CNR Padova, and subm by P Gatto, Ist Studio Dinamica Grandi Masse, CNR Venice.

UD-82. Ca' Bianca **21,700 ± 500**

Peat from Core 8 at depth 17.40m.

UD-87. Ca' Bianca **> 30,000**

Peat from Core 8 at depth 26.00m.

Forte di S Andrea series

Peat and shells from lagoon of Venice Forte di S Andrea (45°26'01"N, 12°22'53"E). Coll 1974 by P Da Roit and subm by P Gatto.

UD-79. Forte di S Andrea **19,500 ± 200**

Peat from Core 13 at depth 15.03m.

UD-80.	Forte di S Andrea	23,500 ± 1000
Peat from Core 9 at depth 27.83m.		
UD-81.	Forte di S Andrea	22,200 ± 1000
Peat from Core 13 at depth 18.00m.		
UD-86.	Forte di S Andrea	23,600 ± 1000
Peat from Core 12 at depth 16.50m.		
UD-89.	Forte di S Andrea	3900 ± 60
Shells from Core 9 at depth 7.15m.		

Cavallino Punta Sabbioni series

Peat from lagoon of Venice Cavallino Punta Sabbioni (45°26'00"N, 12°25'52"E). Coll 1974 by P Da Roit and subm by P Gatto.

UD-83.	Cavallino Punta Sabbioni	21,600 ± 650
Peat from Core 15 at depth 15.32m.		

Cavallino Ca' Pasquali series

Carbonate (mollusk shells) from lagoon of Venice Cavallino Ca' Pasquali (45°27'14"N, 12°29'01"E). Coll 1974 by P Da Roit and subm by P Gatto.

UD-88.	Cavallino Ca' Pasquali	7950 ± 50
Carbonate (mollusk shells) from Core 17 at depth 15.40m.		
UD-90.	Cavallino Ca' Pasquali	5340 ± 80
Carbonate (mollusk shells) from Core 17 at depth 13.05m.		

Alberoni series

Peat and carbonate (mollusk shells) from Alberoni Golf 1 (45°20'49"N, 12°19'20"E) and Golf 2 (45°20'46"N, 12°29'33"E). Coll 1976 by P Da Roit and subm by P Gatto.

Golf 1

UD-136.	Alberoni Golf 1	2940 ± 50
Carbonate at depth 2.40–2.48m.		
UD-137.	Alberoni Golf 1	3590 ± 50
Carbonate at depth 2.84–2.91m.		
UD-138.	Alberoni Golf 1	3980 ± 50
Carbonate at depth 4.35–4.47m.		

UD-139. Alberoni Golf 1 3350 ± 60

Carbonate at depth 5.76–5.92m.

UD-140. Alberoni Golf 1 2970 ± 50

Carbonate at depth 6.78–6.97m.

UD-141. Alberoni Golf 1 4490 ± 50

Carbonate at depth 7.70–7.80m.

UD-142. Alberoni Golf 1 2970 ± 50

Carbonate at depth 8.85–9.05m.

UD-143. Alberoni Golf 1 4950 ± 150

Peat at depth 9.10–9.60m.

UD-144. Alberoni Golf 1 4260 ± 60

Carbonate at depth 10.93–11.07m.

UD-145. Alberoni Golf 1 3250 ± 50

Peat at depth 12.25–12.70m.

UD-146. Alberoni Golf 1 3760 ± 50

Carbonate at depth 12.60–12.70m.

UD-147. Alberoni Golf 1 3090 ± 50

Peat at depth 13.25–13.70m.

Golf 2

UD-148. Alberoni Golf 2 3250 ± 50

Peat at depth 7.50–8.00m.

Fiume Brenta Series 1

Peat and clay from Rubano, Padova (46°26'30"N, 11°47'10"E) at different depths in Brenta area. Radiocarbon dating has been carried out in the frame of a general paleogeographic study of the alluvial plane between Vicenza and Padova (Castiglioni *et al* 1987).

UD-127. Rubano 1 16,000 ± 500

Peat from Rubano 1 at depth 5–7m. Coll and subm 1981 by GB Castiglioni, Dept Geography, Univ Padova.

UD-209. Rubano 4 10,190 ± 150

Silty clay from Rubano 4, Sample L, at depth 2.40–2.70m. Coll and subm 1985 by GB Castiglioni.

UD-210. Rubano 4 18,100 ± 700

Black clay from Rubano 4, Sample N, at depth 7.94–8.00m. Coll and subm 1985 by GB Castiglioni.

UD-211. Rubano 5 19,200 ± 250

Black clay from Rubano 5, Sample P, at depth 7.80–7.86m. Coll and subm 1985 by GB Castiglioni.

UD-212. Via Po 4650 ± 220

Peat from Via Po, Sample Q, at depth 2.63–2.72m. Coll and subm 1985 by GB Castiglioni.

Verona series

Radiocarbon dating of different stratigraphic samples has been carried out to solve some sedimentologic problems in the area of Verona (Sorbini *et al* 1984). Dates range from 470 ± 40 BP to 35,000 ± 2500 BP. In particular, Avesa and S Felice wells indicate that the great fan of the Adige River has to be located in the Würmian period, whereas sediments SE of Verona (Zevio) evidence most recent flood episodes. In this connection, a radiocarbon date (470 ± 40 BP, tree trunk, *Quercus farnia*) fits with archaeological findings.

UD-149. S Felice 30,000 ± 2500

Peat from S Felice, Verona (45°15'00"N, 11°46'20"E) at depth 35–39.5m. Coll and subm 1982 by L Sorbini, Mus Nat Hist, Verona.

UD-150. Basso Acquar 18,800 ± 2000

Peat from Basso Acquar, Verona (45°15'00"N, 11°46'20"E) at depth 16–21m. Coll and subm 1982 by L Sorbini.

UD-151. Avesa 25,000 ± 2000

Peat from Avesa Well no. 2, Verona (45°28'00"N, 11°48'12"E) at depth 20.1–20.2m. Coll and subm 1982 by L Sorbini.

UD-204. Basso Acquar > 31,000

Clayey peat from Basso Acquar Well no. 34, Verona (45°15'00"N, 11°46'20"E) at depth 70–72m. Coll and subm 1984 by L Sorbini.

UD-205. Basso Acquar 24,600 ± 800

Clayey peat from Basso Acquar Well no. 34, Verona (45°15'00"N, 11°46'20"E) at depth 16–21m. Coll and subm 1984 by L Sorbini.

Cordevole Valley series

These determinations are part of a study made by GB Pellegrini, Dept Geography, Univ Padova, to verify the Quaternary evolution of Cordevole Valley, Piave River basin, Belluno (46°09'00"N, 12°06'03"E). All samples coll and subm 1985 by GB Pellegrini, Dept Geography, Univ Padova (Pellegrini & Zambrano 1979).

UD-230. Vallone Bellunese 1 **1230 ± 80**

Fragment of tree trunk found at depth 1m in Mas, Sedico, Belluno (46°08'20"N, 12°06'56"E).

UD-231. Vallone Bellunese 2 **850 ± 80**

Same as UD-230, above.

UD-232. Valle del Cordevole 3 **> 43,000**

Fragment of wood found at depth 2m in Mas.

UD-233. Valle del Cordevole 4 **38,000 ± 2000**

Fragment of wood found at depth 1m in Mas.

UD-234. Valle del Cordevole 5 **42,300 ± 3100**

Fragment of wood found at depth 3m in Mas.

UD-235. Valle del Cordevole 6 **430 ± 80**

Peat found at depth 2m in Mas.

UD-238. Fadalto **640 ± 60**

Fragment of wood found at depth 1m in Fadalto, Belluno (46°04'34"N, 12°20'38"E).

Wood samples

UD-95. Val Belluna **11,180 ± 100**

Wood found during geomorphologic and geophysical research near Levego, Belluno (46°09'09"N, 12°15'30"E) at depth 3.5m to verify morphologic evolution of Val Belluna in Quaternary, with particular regard to Ponte nelle Alpi zone and nearby Valle di S Croce. Coll and subm 1979 by GB Pellegrini.

UD-119. Garda **10,570 ± 200**

Wood found during geomorphologic research near Garda, Verona (45°34'47"N, 10°42'57"E). Fragment from uprooted tree trunk at depth 2.2m. Coll and subm 1981 by U Sauro, Dept Geography, Univ Padova.

UD-164. Levico Lake **1730 ± 100**

Fragment from uprooted tree trunk (*Quercus petraea* Liebl) at depth 16m in Levico Lake, Trento (46°00'35"N, 11°10'25"E). Coll by Sub Club "Trento" and subm by G Tomasi, Tridentine Mus Nat Hist, Trento. Result is part of study of paleoclimate of Valsugana area.

*Somalia, Africa***Mogadiscio series**

Results reported here are part of study in Benadir on coast line between Mogadiscio and Danane to verify sequence of Quaternary sediments.

UD-185. **24,300 ± 500**

Carbonate, shells (*Goniastrea retiformis*) from Ceel Baqal (02°10'30"N, 45°17'20"E) at surface level. Coll and subm 1984 by E Robba, Dept Earth Sciences, Univ Milano, Italy.

UD-186. **3750 ± 100**

Carbonate, shells (*Achatina lactea*) from Ceel Baqal at surface level. Coll and subm 1984 by E Robba.

UD-187. **4250 ± 200**

Carbonate, shells (*Terebralia palustris*) from Ceel Baqal at surface level. Coll and subm 1984 by E Robba.

UD-188. **4990 ± 80**

Carbonate, shells (*Anadara antiquata*) from Ceel Baqal at surface level. Coll and subm 1984 by E Robba.

ARCHAEOLOGIC AND HISTORIC SAMPLES*Italy*

Survey made by G Bernardi to verify first appearance period of playing cards. Coll and subm 1979 by G Bernardi, Trieste.

UD-85. **925 ± 100**

Tissue paper used for producing playing cards found in printshop in Modena (45°12'13"N, 12°00'12"E).

Comment: radiocarbon date does not fit with age suggested by historic research (ca AD 1400), probably because tissue was produced long before it was used for paper production.

Torcello series

In northern part of Lagoon of Venice (Upper Adriatic Sea), sequence of Holocene sediments reveals evolution from subaerial sedimentary environment to marsh formation (Favero & Barbero 1983). Evidence of sea transgression is present in third stage. Archaeologic findings place this event between Roman imperial epoch and 5th century AD. Samples coll 1980 by P Da Roit and subm by V Favero, Ist Studio Dinamica Grandi Masse, CNR, Venice.

UD-100. Scanello 5 D **810 ± 70**

Fragment of tree trunk (*Carpinus betulus*) from the Scanello Channel (45°29'20"N, 12°00'50"E) at depth 1m.

Comment: sample is from brick kiln archaeologically dated to ca 1st century BC. Radiocarbon age does not agree with archaeologic findings, dated about 1st century BC.

UD-101. Scanello 5 **260 ± 60**

Fragment of tree trunk (*Salix viminalis*) from the Scanello Channel at depth 3m.

Comment: sample is from medieval building historically dated to ca end of 17th century.

UD-102. Canale di S Lorenzo **1900 ± 200**

Fragment from valve used for diverting water from San Lorenzo Channel at ground level.

Comment: channel is archaeologically dated to ca 1st–2nd century BC.

Venice series

Samples found in various locations of Lagoon of Venice during archaeologic survey to verify reliability of estimated ages and radiocarbon dating of origins of Venice (Dorigo 1983). All samples coll and subm 1981 by W Dorigo, Dept Letters, Univ Venice.

UD-109. 7 SOL 2, Sample 1 **2280 ± 60**

Fragment of lower part of wooden column from Barena del Vigno (45°31'00"N, 12°00'42"E).

UD-110. 7 SOL 2A **2350 ± 60**

Fragment of pole from Barena del Vigno.

UD-111. 7 SOL 2A **2810 ± 60**

Adhesive oysters (*Ostrea* sp) on pottery fragments from Barena del Vigno.

UD-112. 7 SOL 2A **2480 ± 60**

Wooden fragment with some nails, perhaps part of boat keel from Barena del Vigno.

UD-114. S M di Gaia **770 ± 80**

Fragment of pile at depth 2.70m in Motta di S Maria Mad di Gaia (45°29'50"N, 12°00'42"E).

UD-115. Salina Vetere **1360 ± 50**

Fragment of pole at depth 3.10m in S Erasmo (45°28'10"N, 12°02'00"E).

UD-116. Torcello **610 ± 80**

Fragment of lower part of pole from Torcello (45°29'50"N, 12°01'54"E).

UD-117. Scanello **1080 ± 80**

Fragment of plank with some nails, perhaps part of boat, from Scanello channel (45°29'20"N, 12°00'50"E).

UD-118. Arsenale **900 ± 100**

Fragment of lower part of pole from Arsenale (45°26'06"N, 12°06'20"E).

UD-165. Uzzo cave 6720 ± 80

Charcoal from Holocene levels in Uzzo cave on E slope of Cape S Vito, Trapani (38°06'36"N, 12°46'53"E) at depth 1.20m. Coll 1983 by M Piperno and subm by F Zevi, Mus Prehist Etn "L Pigorini," Rome.

Comment: research was carried out 1975–1983 in Uzzo cave to understand Neolithic process in W Sicily as well as use of cave as necropolis in Mesolithic period (Piperno 1985).

La Vela series

Results are part of study of La Vela cave to understand Neolithic sequence in different stages.

UD-175. La Vela, Gruppo Gaban 4850 ± 200

Charcoal from Neolithic layers of La Vela cave, Trento (46°04'55"N, 13°21'05"E) at depth 2.50m. Coll and subm by G Tomasi, Tridentine Mus Nat Hist, Trento.

UD-176. La Vela, Vasi a bocca quadrata 5370 ± 180

Same as UD-175, above.

S Daniele del Friuli series

Charcoal samples were found during archaeological surveys of Castello church in San Daniele del Friuli (46°13'18"N, 12°06'23"E). Coll and subm 1985 by F Piuze, Dept Antiquities, Udine. Charcoal was assoc with two furnaces, probably used to fuse metal and glass.

UD-213. Carboni I forno 1170 ± 120

Charcoal coll at depth 1.43m.

UD-214. Carboni II forno 790 ± 50

Charcoal coll at depth 0.95m.

REFERENCES

- Barbina, V, Calligaris, F and Ciuti, P 1979 An apparatus for beta counting, used in a radiocarbon dating laboratory. *Nuclear Instruments & Methods* 163: 545–548.
- Barbina, V, Calligaris, F, Del Fabbro, A, Turello, A and Ciuti, P 1982 Udine Radiocarbon Laboratory date list I. *Radiocarbon* 24(2): 214–216.
- Castiglioni, GB, Girardi, A and Rodolfi, G 1987 Le tracce degli antichi percorsi del Brenta per Monta' e Arcella nei pressi di Padova: studio geomorfologico. *Mem Geol Padova Ist Geol Univ* 39: 129–149.
- Dorigo, W 1983 *Venezia origini*. Milano: 1–247.
- Favero, V and Barbero, RS 1983 Oscillazioni del livello del mare ed evoluzione paleoambientale della laguna di Venezia nell'area compresa tra Torcello ed il margine lagunare. *Soc Ven Nat Sci* 8: 83–102.
- Gatto, P 1980 Il sottosuolo del litorale veneziano. *Venezia Ist Studio Dinamica Grandi Masse CNR Tech rept* 108: 1–18.
- Pellegrini, GB and Zambrano, R 1979 Il corso del Piave a Ponte nelle Alpi nel Quaternario. *Studi Tridentini di Scienze Naturali, Geologica* 56: 69–100.
- Piccoli, G and Robba, E 1983 The last marine submersion of the Bangkok area in Thailand. *Ist Geol Univ Mem Geol Sci Padova* 36: 343–352.
- Piperno, M 1985 Some C-14 dates for the palaeoeconomic evidence from the Holocene levels of Uzzo cave, Sicily. In Malone, C and Stoddart, S, eds, *The Cambridge Conf Part i: The human landscape. Papers in Italian Archaeology IV, Bar internatl ser* 244: 83–86.
- Sorbini, L, Accorsi, CA and Bandini Mazzanti, M 1984 Geologia e geomorfologia di una porzione della pianura a SE di Verona. *Verona Mus Nat Hist Mem* 2: 9–87.