UCLA RADIOCARBON DATES VIII*

RAINER BERGER and W. F. LIBBY

Institute of Geophysics, University of California, Los Angeles 90024

The measurements reported were made during the 2nd half of 1967 in the Isotope Lab. of the Inst. of Geophysics and Planetary Physics as a continuation of the UCLA date lists I through VII. Samples were analyzed as CO₂-gas at close to one atm in a 7.5 L proportional counter with 3 energy channels described earlier. Radiocarbon ages were calculated for uniformity on the basis of a 5568 yr half-life as was recommended by the Sixth International C¹⁴ and H³ Dating Conference, June 1965, in Pullman, Washington. The standard for the contemporary biosphere remains as 95% of the count rate of NBS oxalic acid for radiocarbon labs. Background determinations have been based on CO₂ obtained from marble. The error listed is always at least a 1σ statistical counting error. In critical cases C¹³/C¹² isotope ratio measurements were made to correct the dates for fractionation. All samples were subjected to accepted NaOH and/or HC1 pretreatments depending on the individual case as a minimum to exclude contamination.

A recent summary on the development of radiocarbon dating is that by Libby, 1965.

ACKNOWLEDGMENTS

We are indebted to the National Science Foundation for continued financial support (GA-628) and acknowledge the competent assistance of Jane Brown (Dept. of Mathematics), Mrs. Wanda Schmukler (Inst. of Geophysics), and Barbara Turring (Dept. of Psychology), the cooperation of I. R. Kaplan and Chari Petrowski (Inst. of Geophysics) in determining C¹³/C¹² isotope ratios, and the collaboration of T. Y. Ho (Esso Production Research, Houston, Texas), L. F. Marcus (Queens College, Flushing, New York) and R. E. Taylor (San Fernando Valley State College).

SAMPLE DESCRIPTIONS

A. United States

La Brea Tar Pit series

Bones from La Brea tar pits, Los Angeles, California (see UCLA V for details on locality and history). Part of extensive analysis of late Pleistocene fossil community now using bone of extinct species. Data with each bone represent pit, depth, and grid data on 3 foot square grid system (Howard, 1962).

Dense mid-shaft bone of femur of *Smilodon californicus* (sabre-tooth tiger) was used, except for UCLA-1292L which is tibia of same species.

Free amino acids were dated after liquid chromatography of collagen hydrolysate prepared according to procedures modified from Ho (1965)

* Publication number 650, Institute of Geophysics and Planetary Physics, University of California, Los Angeles.

by T. Y. Ho. This procedure permits complete removal of petroleum contamination. NaOH treatment for possible humic acid contamination was included in a few samples as indicated. Complete procedure will be published elsewhere (Ho, Marcus, and Berger, 1968). Subm. 1967 by L. F. Marcus.

LACMVP RLB A-675, A-511. Pit 3, coordinates E-5, 22 ft. Left femur. (UCLA-1292J is another sample from same bone treated with NaOH.)

UCLA-1292B. La Brea Tar Pits 12,650 \pm 160 \pm 10,700 B.C. LACMVP RLB A-642. Pit 3, coordinates C-4, 7 ft. Left femur.

14 700 . 16

 UCLA-1292C.
 La Brea Tar Pits
 $14,500 \pm 190$

 12,550 B.C.

LACMVP RLB A-398, A-719. Pit 3, coordinates E-2, 12 ft. Left femur. At same level as tree data (LJ-55, LJ-89, LJ-21; La Jolla I) (Y-354, Y-355; Yale IV), note UCLA-1292A is below this level, UCLA-1292B above, UCLA-1292E at approx. same level.

 UCLA-1292D.
 La Brea Tar Pits
 $28,000 \pm 1400$

 26,050 B.C.

LACMVP RLB A-411. Pit 4, coordinates D-2, 15.5 ft. Left femur.

 $14,400 \pm 2100$ UCLA-1292E. La Brea Tar Pits 12,450 B.C.

LACMVP RLB A-835. Pit 3, coordinates C-4, 11.5 ft. Right femur.

LACMVP RLB A-797. Pit 13, coordinates E-11, 11 ft. Left femur.

 UCLA-1292G.
 La Brea Tar Pits
 $26,700 \pm 900$

 26,700 ± 900 29,750 B.C.

LACMVP RLB A-846. Pit 4, coordinates D-2 and 4, 8 ft. Right femur.

LACMVP RLB A-4435. Pit. 60, coordinates C-10, 9 to 12 ft. Right femur.

UCLA-1292I.La Brea Tar Pits $15,300 \pm 200$ 13,350 B.C.

LACMVP RLB A-4436. Pit 13, coordinates F-10, 14.5 ft. Right femur.

 20.500 ± 900

18,550 в.с. UCLA-1292J. La Brea Tar Pits

LACMVP RLB A-675, A-511, Additional sample from same specimen as UCLA-1292A, treated with 1/10 N NaOH (Berger, Horney, and Libby, 1964).

 19.300 ± 395

UCLA-1292K. La Brea Tar Pits

17.350 в.с.

LACMVP RLB K-3407. Pit 3, coordinates E-4, 26 ft. Right femur. Treated with 1% NaOH.

 15.200 ± 800

UCLA-1292L. La Brea Tar Pits

13,250 в.с.

LACMVP RLB T-6101. Pit 4, coordinates C-2, 11.5 ft. This is suspected artifact; bone was carefully scraped from inside (included some cancellous bone). Sample treated with 1% NaOH.

General Comments (L.M.): the 5 specimen from Pit 3 indicate stratification in that deposit into 2 time periods. There is also preliminary indication that Pit 13 may be of same age as younger part of Pit 3. (R.B.): it had been thought that action of natural gas moving through levels of tar pits might exercize stirring action, greatly increasing difficulties of archaeologist and palaeontologist. Apparently, phenomenon was overestimated.

 260 ± 80

UCLA-1296. Abalone shell, Malibu, California A.D. 1690

Haliotis cracheroidi from cemetery of village site near junction of Virgines and Malibu Canyons, Malibu, California (34° 5′ 45" N Lat, 118° 45' W Long). Pit S-10, W-22, 30-39' below datum. Date corrected by -160 yr according to Berger, Taylor, and Libby (1966). Coll. and subm. by C. King, UCLA Archaeol. Survey.

B. Mexico

La Venta series

Continuation of series in UCLA VII concerning age of La Venta, Tabasco, site (18° 10' N Lat, 94° 5' W Long). Samples coll. July 1967 by R. F. Heizer, Univ. of California, Berkeley and P. Drucker, San Andreas Tuxtla, Mexico; subm. and commented on by R. F. Heizer and R. Berger.

 2300 ± 60

UCLA-1330. La Venta

350 в.с.

Charcoal from NW perimeter of Mound A-2. Stratigraphy not clear to which building phase of La Venta this belongs, but date indicates Phase IV.

 2660 ± 140

UCLA-1331. La Venta

710 в.с.

Charcoal stratigraphically below sample UCLA-1330 in same trench. Because no direct tie-in with phase-dated levels of 1955 excavation could be made, C¹⁴ age can be interpreted as belonging to Phase II or III.

UCLA-1332. La Venta

 2550 ± 80 600 B.C.

Charcoal from thin clay fill layer overlying Phase III old-rose floors inside and at W edge of Ceremonial Court area. Date agrees with recent revision of age of La Venta site (Berger, Graham, and Heizer, 1967).

 410 ± 80

A.D. 1540

UCLA-1288. Mexican idol

 $\delta C^{13} = -24.29\%$

15-cm long carved wooden idol from Guerrero. No date known for respective style. Coll. and subm. 1967 by P. T. Fürst, Univ. of California, Los Angeles.

 700 ± 80

UCLA-1298. Tecualillo, Navarit

A.D. 1250

Charcoal from site in Teacapan estuary (22° 27' N Lat, 105° 39' W Long). Test square A, 6 ft deep. Ceramic styles indicate late site for which confirmation was sought. Coll. by S. Scott, State Univ. of New York, Buffalo and subm. through C. W. Meighan, Univ. of California, Los Angeles.

C. South America

Northern Chile series

Undertaken under the Univ. of California/Univ. of Chile Cooperative Program to explore general archaeol. of N Chile which up till now is rather uncertain. Conclusions can only be drawn after more extensive investigation and dating. Samples coll. and subm. if not noted otherwise by D. L. True, Univ. of California, Davis, and C. W. Meighan.

 4690 ± 80

UCLA-1293.

2740 B.C.

Charcoal from small camp near Quebrada Tarapaca, Tarapaca (ca. 20° S Lat, 69° 30' W Long). Assoc. with preceramic complex in Tr-12-1. Coll. Jan. 1967.

 680 ± 80

UCLA-1294A.

A.D. 1270

Matting from Site A-Z-8, Burial N 2/5, San Miguel de Azapa, (18° 34' S Lat, 70° 12' W Long). Dates San Miguel culture. Coll. 1965 by O. Espoueys.

 280 ± 80

UCLA-1294B.

A.D. 1670

Textile from same site as UCLA-1294A. Dates Burial M-10, San Miguel Tardio phase. Coll. Nov. 1965 by J. C. Montane.

 580 ± 80

UCLA-1294C.

A.D. 1370

Matting from same site as UCLA-1294A. Dates Burial M27 of San Miguel Tardio phase. Coll. Nov. 1967, J. Montane.

 810 ± 80

UCLA-1294D.

а.р. 1040

Textile from same site as UCLA-1294A. Dates Burial M4/2, San Miguel culture. Coll. by O. Espoueys.

 610 ± 60

UCLA-1294E.

а.р. 1340

Matting. Another date for same feature as UCLA-1294C.

D. Pacific

New Hebrides series

Continuation of archaeol. exploration of New Hebrides, UCLA IV. Samples coll. and subm. by R. Shutler, Jr., San Diego State College, San Diego, California.

 645 ± 80

UCLA-1295A. Tanna

а.р. 1305

Charcoal from heavy occupation level in cave, TaRs 1, Pit 16, depth 36 to 42 in. (19° 32′ 30″ S Lat, 169° 15′ 30″ E Long). Assoc. with stone and shell artifacts, and shell and bone debris from human habitation. Sample dates Conus cup bead type (Conus miles). This type of bead also found on Futuan and Efate Islands. Coll. 30 Mar. 1964.

 1095 ± 80

UCLA-1295B. Tanna

A.D. 855

Charcoal from same location as preceding sample but from Pit 14, depth 24 to 30 in., just above heavy occupation level. Sample dates stone disc bead type. Coll. 30 Mar. 1964.

 $\textbf{455} \pm \textbf{80}$

UCLA-1295C. Efate

а.р. 1495

Charcoal from lowest level of rock shelter, EfRs 7, Tr. C, depth 26 to 32 in., on Efate (17° 45′ 0″ S Lat, 168° 17′ 30″ E Long). Assoc. with gorge fish hook and other debris. This hook type occurs also on Tanna, Futuan, and in Polynesia.

 170 ± 80

UCLA-1295F. Futuna

а.в. 1780

Charcoal from feature FuRs 12, Tr. 7, depth 18 to 24 in. (Level 5) on Futuna (19° 30′ 50″ S Lat, 170° 13′ 30″ E Long). Assoc. with shell and bone from human habitation. Might indicate introduction of taro into S New Hebrides as id. by pollen analysis.

 790 ± 80

UCLA-1295G. Futuna

A.D. 1160

Charcoal from Feature FuRs 12, Tr. 3, depth 18 to 24 in. near bottom of trench on Futuna (19° 30′ 30″ S Lat, 170° 13′ 30″ E Long). Assoc. with human burials, shell artifacts, and shell and bone debris from human occupation. Sample dates Conus cup bead type found also on Tanna (UCLA-1295A, above) and Efate.

Guam sherd series

Part of study to determine suitability of carbon-bearing potsherds for C14 dating by comparing sherds with assoc. charcoal samples (Taylor and Berger, 1968). Samples coll. by F. Reinman, Chicago Nat. Hist. Mus., 1965, to evaluate ceramic sequence on Guam (13° 30' N Lat, 144° 40' E Long). Samples obtained through courtesy of C. W. Meighan and subm. by R. E. Taylor and R. Berger.

 420 ± 100

UCLA-1232A. Guam

A.D. 1530

Charcoal from Pit 3, 6 to 12 in. depth.

 495 ± 80

UCLA-1232B. Guam

A.D. 1455

Sherds from same stratigraphic position as UCLA-1232A.

 965 ± 80

UCLA-1232C. Guam

A.D. 985

Charcoal from Pit 3, 24 to 30 in. depth.

 895 ± 120

UCLA-1232D. Guam

A.D. 1055

Sherds from same location as UCLA-1232C.

 320 ± 80

UCLA-1232G. Guam

A.D. 1630

Charcoal from Pit 4, 6 to 12 in. depth.

UCLA-1232H. Guam

 $275. \pm 80$ A.D. 1675

Sherds from same stratigraphic unit as UCLA-1232G.

 805 ± 80

UCLA-1232I. Guam

A.D. 1145

Charcoal from Pit 4, 18 to 24 in. depth.

 670 ± 100 A.D. 1280

UCLA-1232J. Guam

Sherds from same location as UCLA-1232I.

General Comment (R.B.): Reasonable agreement between sample pairs is indicated suggesting that C14 dating of suitable ceramics is feasible. C. W. Meighan thinks dates based on ceramics may be preferable to charcoal twin as chances of stratigraphically disturbing many sherds are less than for compact charcoal sample. Additional Guam samples are GaK-1356-1371 (ms. in preparation).

E. Europe

European medieval architecture series

Continuation of investigation in UCLA III-VII into potential and limitations of radiocarbon dating in Middle Ages. For maximum precision δC^{13} measurements are included, also sample location in timber as well as comparison with secular variations of C^{14} concentration in dendrochronologically dated wood (Suess, 1965). Samples coll. 1967, subm. and commented on by W. Horn, Univ. of California, Berkeley and R. Berger unless noted otherwise.

UCLA-1301. Canteloup

 $\mathbf{580} \pm \mathbf{80}$

 $\delta C^{13} = -26.88\%$

Oak sapwood from waney edge of Post V (carpenter's mark) S side of barn at Canteloup presently being dismantled and re-erected as church on grounds of Abbaye de Fontenelle, Saint Wandrille, Seine-Maritime, Ignace Dalle, Abbot. The building type of aisled and bay-divided medieval timber hall was and still is equally suitable as house, barn, festal hall or church as discussed in Horn and Born (1965). Coll. 19 July 1967.

 650 ± 60

UCLA-1302. Canteloup

 $\delta C^{13} = -24.16\% e$

Oak heartwood sample from same post used in UCLA-1301. Wood located ca. 30 yr from center of tree which was felled at age of ca. 100 yr. Comment (R.B.): when all correction factors are taken into account, Canteloup was erected most probably at very end of 13th or beginning of 14th century and will provide genuine medieval church at its new location (49° 32′ N Lat, 0° 45′ E Long).

 380 ± 80

UCLA-1303. Richelieu

 $\delta C^{13} = -24.54\%$

Oak bark from lowest longitudinal rail marked VII between Posts VI and VII on S side of central aisle of market hall in Richelieu built by cardinal as model village *in toto* between A.D. 1631 and ca. 1640 near Tours (47° 1′ N Lat, 0° 20′ E Long). Growth-increment of bark correction 30 yr. Coll. 30 July 1967. *Comment* (R.B.): calculated most probable historical age of erection: ca. A.D. 1630.

 930 ± 60

UCLA-1306. Sully

 $\delta C^{13} = -23.04\%$

Chestnut wood (Castania vesca) from curved long ashlar piece rising from floor to rafters in upper hall of castle of Sully-s-Loire (47° 46′ N Lat, 2° 22′ E Long). Castle built in A.D. 1363 according to L. Martin (1962). Comment: corrected date permits probable historical age from A.D. 1050 to 1220. Exact original position of timber in tree uncertain due to construction placement.

 $\textbf{515} \pm \textbf{60}$

UCLA-1307. Arpajon

 $\delta C^{13} = -23.83\% e$

Oak heartwood from market hall in Arpajon (48° 35′ N Lat, 2° 15′ E Long) built according to local tradition in A.D. 1450-1470 by Louis Mallet de Graville. Tree-ring correction 40 yr. Coll. 10 July 1967. *Comment*: probable historical date ca. A.D. 1450.

 400 ± 60

UCLA-1308. Troussures

 $\delta C^{13} = -24.42\%$

Chestnut sapwood from post replacing the original stone arcade pillar in 13th-century abbey barn of Troussures. Post bears date A.D. 1609 by carpenter. Coll. 17 July 1967. Comment: probable historical date ca. A.D. 1600 or 1450.

 860 ± 80

UCLA-1309. Maubuisson

 $\delta C^{13} = -22.58\%$

Waney edge of Truss 9, S brace with carpenter mark 9 of barn of Abbey of Maubuisson founded A.D. 1236. *Comment*: probable historical age ca. A.D. 1050-1220 due to plateau in curve of secular variations in C¹⁴ content of wood.

 330 ± 60

UCLA-1310. Questembert

 $\delta C^{13} = -26.02\%$

Bark from post of Truss IV (center aisle post), S side of building of market hall at Questembert (47° 40′ N Lat, 2° 36′ W Long). Barn built A.D. 1675 as marked on tie-beam. Bark growth increment correction —40 yr. *Comment*: probable historical age ca. A.D. 1660.

 420 ± 60

UCLA-1312. Milly

 $\delta C^{13} = -22.80\%_0$

Oak wood peg from original post of market hall sawed and trimmed into bench at St. Blais des Simples, burial chapel of Jean Cocteau (48° 24′ N Lat, 2° 28′ E Long). Market hall built A.D. 1479 by Louis Mallet de Graville. *Comment*: probable historical age ca. A.D. 1450.

 555 ± 60

UCLA-1313. Parcey-Meslay

 $\delta C^{13} = -24.23\%_0$

Oakwood from waney edge of Post 4 counted from W, S side of barn of Parcey-Meslay, near Tours (47° 28′ N Lat, O° 21′ W Long). Barn built in time of Abbot Hugue de Rochecourbon (A.D. 1211-1227). Coll. 31 July 1967. *Comment*: previous measurement UCLA-570 (UCLA IV) at A.D. 1215 is corroborated by this date which after modification yields time span of ca. A.D. 1400-1250.

 670 ± 60

UCLA-1314. Lenham

 $\delta C^{13} = -24.30\%e$

Charred oak sapwood from arcade plate in W part of minor barn at Lenham, Kent (51° 13′ N Lat, O° 38′ W Long). A 13th-century document mentions destruction of church and subsidiary buildings by fire in A.D. 1298. Tree-ring correction —15 yr. Coll. 19 August 1967. Comment: probable historical date based on C¹⁴ measurement after correction yields A.D. 1250.

 560 ± 60

UCLA-1315. Ter Doest

 $\delta C^{13} = -24.19\%$

Oakwood from waney edge from transverse brace reaching from suspended wall-piece to aisle-tie northernmost freestanding truss. Barn of Abbey of Ter Doest near Bruges (51° 16' N Lat, 3° 12' E Long).

Coll. 19 Aug. 1967. Comment: adjusted radiocarbon date yields probable historical span of erection from A.D. 1270-1400. Re-examination of previous dates UCLA-568A and B (UCLA IV) and UCLA-1036 and 1038 (UCLA VI) together with present date allow for time span of construction from late 13th through 14th centuries. Stylistically, barn is late 13th century when compared with elements of historically dated church a few mi. away.

 630 ± 60

UCLA-1316. Church Enstone

 $\delta C^{13} = -24.61\%$

Bark of oak from foot of N cruckblade of Truss 5, upper face, E edge, 12 in. above pad. Church dated by inscription A.D. 1382. Coll. 23 Aug. 1967. Growth increment correction 40 yr. *Comment*: probable historical date range: 14th century.

 120 ± 80

UCLA-1317. Beaumont-in-Gatinais

 $\delta C^{13} = -23.85\% e$

Oak bark from Post h' SW edge, directly under tie-beam of Beaumont Hall (48° 8' N Lat, 2° 29' E Long). Comment: probable historical age of post as old as mid-17th century.

 490 ± 60

UCLA-1318. Egreville

 $\delta C^{13} = -22.52\%$

Oak wood from central shake in Post c' under bracing strut, W face of market hall in Egreville (48° 10' N Lat, 2° 52' E Long). Coll. July 1967. Tree-ring allowance 30 to 50 yr. *Comment*: probable historical date first half of 15th century.

 890 ± 60

UCLA-1338. St. Pierre-sur-Dives

 $\delta C^{13} = -23.03\%$

Oak heartwood from central shake under aisle tie-beam of Post 1, NE corner of market hall (49° 1′ N Lat, 2′ W Long). One of apparently 4 old posts, rest all post-1945 replacements. Coll. 24 July 1967. *Comment*: probable historical age: 12th to mid-13th centuries.

 550 ± 60

UCLA-1340. Frindsbury

Oak sapwood from 2. post of W row counting from N, NW edge of barn at Frindsbury near Rochester (51° 24′ N Lat, 30′ E Long). Coll. 21 Aug. 1967. *Comment*: probable historical age: end of 13th through all of 14th centuries.

 450 ± 60

UCLA-1341. Brook

 $\delta C^{13} = -24.34\%$

Oakwood from waney edge from shoring beam, N aisle, easternmost truss, upper side, W edge of Court Lodge Barn at Brook near Wye (51° 11′ N Lat, 56′ E Long). Coll. 20 Aug. 1967. Comment: probable historical age: ca. A.D. 1450.

570 + 60

UCLA-1342. Leigh Court

 $\delta C^{13} = -25.25\%$

Oak sapwood from upper fare of N blade of Truss 2, 3'6" above springing of blade of Leigh Court Barn near Worcester (52° 11' N Lat, 2° 13' W Long). *Comment*: probable historical age: end of 13th through 14th centuries. Leigh Court appears to be most beautiful extant cruck barn discovered by F. W. D. Charles and is of great importance in history of cruck construction.

 390 ± 60

UCLA-1257. Harwell Church

 $\delta C^{13} = -26.37\%$

Oakwood from N transept tie-beam of Harwell Church (51° 37′ N Lat, 1° 18′ W Long). Some 60 tree-rings removed and older than UCLA-1250 (UCLA VII). Implications of this measurement and earlier ones with respect to secular variations in C¹⁴ content of wood are discussed in Fletcher, 1968. Coll. 1962 and subm. by J. M. Fletcher, Harwell, England.

Ukrainian Bronze age series

First radiocarbon dates on aspects of Ukrainian Bronze age in greater framework of Indo-European migrations. Samples obtained 1966 through Inst. of Archaeol., Soviet Acad. of Sci., Moscow by M. Gimbutas, Univ. of California, Los Angeles.

 $\textbf{4210} \pm \textbf{80}$

UCLA-1270. Tsatsa

2260 в.с.

Wood from Barrow 6, Grave 3 at Tsatsa (48° 13' N Lat, 44° 40' E Long). Early Yamna culture. Coll. 1962 by V. P. Shilov.

 $\textbf{4150} \pm \textbf{80}$

UCLA-1271. Ust'man

2200 в.с.

Wood from Barrow 1, Grave 13 at Ust'man. Early Yamna culture. Coll. 1962 by V. P. Shilov.

 3390 ± 80

UCLA-1272. Argadinskaja

1440 в.с.

Wood from Barrow 9, Grave 4 at Argadinskaja. Timber Grave culture. Coll. 1959 by V. P. Shilov.

 $\mathbf{3860} \pm \mathbf{80}$

UCLA-1273. Kudinov

1910 в.с.

Wood from Barrow 2, Grave 6 at Kudinov. Catacomb Grave culture. Coll. 1961 by A. P. Mantsevich.

 $\textbf{3525} \pm \textbf{80}$

UCLA-1274. Kudinov

1575 в.с.

Wood from Barrow 1, Grave 7 at Kudinov. Timber Grave culture. Coll. 1961 by A. N. Melent'ev. *Comments* (M.G.): dates fall in expected time ranges and confirm expectations. (R.B.): when tree-ring calibrated C¹⁴ dating is applied, age increases by 3 to 600 yr for some samples.

F. Africa

 7560 ± 1000 5610 B.C.

UCLA-1247E. Lothagam, Turkana, Kenya

Shells from basal level of fishing camp on 210 ft beach of Lake Rudolph (4° N Lat, 36° E Long) assoc. with burials showing robust physical type and mesolithic industry with bone harpoons and leister points plus micro- and macrolithic stone technology. Coll. 1966 by L. Robbins and subm. by J. D. Clark, Univ. of California, Berkeley. Comment (RB): date is uncorrected and may not represent best obtainable age until more data are forthcoming on radiocarbon content of Lake Rudolph. Margin of error has been increased to reflect uncertainty.

Fingira Rockshelter series

Located SW end of Nyika Plateau, Malawi (10° S Lat, 33° E Long) between 7 to 8000 ft. Occupied in Later Stone age times by short, robust group, of which 2 burials and other scattered human bones have been found. Material culture includes microlithic tools in quartz, interesting bone industry, shell beads, pendants, much ocher. Found also was much food waste. Schematic paintings occur on rock-shelter wall. Deposit was shallow, 33 in. where tested. Practically no evidence of Iron age occupation. Coll. Aug. 1966 by Beatrice Sandelowsky and K. R. Robinson; subm. by J. D. Clark.

 $\mathbf{3260} \pm \mathbf{80}$

UCLA-1258. Fingira

1310 в.с.

Charcoal from Trench E, 15 to 18 in. deep.

 3430 ± 80

UCLA-1259. Fingira

1480 в.с.

Charcoal from Test Pit E, 27 to 33 in. deep. *Comment* (J.D.C.). data indicates the Nyika was settled by hunters and gatherers earlier than expected. Archaeological assemblage very different from contemporary assemblage found in Malawi Rift.

 $\mathbf{885} \pm \mathbf{80}$

UCLA-1289. Mwavarambo

A.D. 1065

Charcoal from Test Pit B 2, 16 to 18 in. deep, Mwavarambo site already dated as UCLA-1242 (UCLA VII) at 655 ± 80 . Date confirms contemporaneity of Mwavarambo Ware at this site with two other early Iron age wares in Karonga district. Coll. July 1966 by K. R. Robinson; subm. by J. D. Clark.

 1100 ± 80

UCLA-1299. Vintukutu

a.d. 850

Charcoal with sherds at Vintukutu Iron age Site 54 8 to 10 in. below surface. This is earlier Iron age site 40 mi. S of those excavated in Karonga district (9° 54′ S Lat, 33° 55′ E Long) (UCLA VII) showing affinites with Mwamasapa pottery and derivative wares. Coll. 1966 by K. R. Robinson; subm. by J. D. Clark.


General Comment (R.B.): preceding African dates and those in UCLA VII are discussed in detail in individual papers in Proceedings of VI. Congrès Panafricain de Prehistoire et de L'Etude du Quaternaire, Dakar, Senegal, December 1968.

G. C¹⁴ in Atmospheric Carbon Dioxide

Atmospheric Radiocarbon Activity series, California

This series is continuation of data published in UCLA IV-VI. C¹⁴ content in ground level atmospheric CO₂ is monitored monthly at China Lake, California (35° 37′ N Lat, 117° 41′ W Long). Samples are coll. with cooperation of Gilbert Plain, Assoc. Head, Research Dept., Naval Ordnance Test Station, China Lake, California.

The following list contains exposure times of NaOH solutions to air and percent increase of C¹⁴ above reference level of 1890 or 0.95 NBS oxalic acid. Data are graphed in Fig 1. The statistical error is less than one-half percent.



Sample no.	Exposure time	C^{14} , %
UCLA-1154	26 Nov.—3 Dec. 1966	+66.9
UCLA-1155	28 Dec.—4 Jan. 1967	+66.8
UCLA-1132	28 Jan.—5 Feb. 1967	+67.5
UCLA-1168	25 Feb.—4 Mar. 1967	+67.3
UCLA-1174	10 Apr.—17 Apr. 1967	+66.9
UCLA-1175	29 Apr.—5 May 1967	+65.9
UCLA-1179	27 May—3 June 1967	+65.8
UCLA-1182	2 July–9 July 1967	+65.1
UCLA-1183	7 Aug.—13 Aug. 1967	+62.8
UCLA-1184	1 Sept.—8 Sept. 1967	+64.9
UCLA-1185	10 Oct.—17 Oct. 1967	+64.1
UCLA-1187	28 Oct.—4 Nov. 1967	+64.0
UCLA-1188	26 Nov.—3 Dec. 1967	+60.5

Atmospheric Radiocarbon Activity series, Brazil

 $\rm C^{14}$ content in ground level atmospheric $\rm CO_2$ was also monitored for several months at Escola de Engenharia, Sao Carlos, Brazil (22° S Lat, 47° 52′ W Long). Samples were coll. with cooperation of M. Tolentino, Dept. of Chem. and Geol. through J. H. Reynolds, Univ. of California, Berkeley and U. G. Cordani, Univ. of Sao Paulo.

Sample no.	Exposure time	Brazil	Calif.
		C^{14} , $\%$	C^{14} , $\%$
UCLA-1156	27 July–1 Aug. 1966	+64.8	+66.6
UCLA-1157	24 Aug.—31 Aug. 1966	+64.1	+72.3
UCLA-1158	24 Sept.—1 Oct. 1966	+65.4	+70.4
UCLA-1159	24 Oct.—31 Oct. 1966	+79.0	+67.2
UCLA-1180	23 Nov.—30 Nov. 1966	+62.5	+66.9
UCLA-1181	27 Dec.—30 Dec. 1966	+63.2	+66.8

For convenience, California data from UCLA VI for similar time periods have been added. Comparison shows almost complete gross equilibration of atmospheres of both hemispheres.

H. Geochemical Determination

UCLA-1186. Methane

>40,000

Gas sample from location near 4843 Longridge, Van Nuys, California. Question if natural gas or sewage gas leak which kills deeprooting trees in area. Subm. by F. J. Folmer and J. R. Mazelli, S California Gas Co., 1967. *Comment* (R.B.): natural gas origin is indicated, but it is still undecided if imported natural gas or seepage from geologic oil or gas bearing formations below Van Nuys.

UCLA-1228. Mt. Vesuvius, Italy

 $30,700 \pm 1100$ 28,750 B.C.

Presumably dates eruptive phase of Vesuvius. Carbonized wood enclosed in tuff considered from geomorphological evidence to be of Würm age. Subm. 1967 by E. Franco, Inst. di Mineralogia, Univ. of Naples.

I. Vegetation and Climate

Neotoma Midden series

Part of dating program begun with UCLA III-VI to use macrofossil plan rests of packrat middens to infer environmental conditions at time of deposition. Coll. 1966 and subm. by P. W. Wells, Univ. of Kansas, Lawrence and R. Berger.

 1860 ± 80

A.D. 90

UCLA-1098A. Laramie Basin, Wyoming

Midden from 7500 ft, Site 1, No. 25. Contains *Pinus ponderosa* and *Juniperus scopulorum* in abundance in the midden whereas no living *Pinus* at site today and only 2 *Juniperus* barely surviving. Indicates time of more favorable growth conditions for woodland as well as higher rainfall for area.

 4060 ± 80

UCLA-1098B. Laramie Basin, Wyoming

2110 в.с.

Midden from Site 2, No. 26. Contains Pinus ponderosa and Juniperus scopulorum.

 $\mathbf{2320} \pm \mathbf{80}$

UCLA-1098D. Laramie Basin, Wyoming

370 в.с.

Midden from 7500 ft, Site 4, No. 28. Contains Juniperus scopulorum.

 1735 ± 80

UCLA-1098E. Laramie Basin, Wyoming

A.D. 215

Wood from beneath sand dune (Juniperus scopulorum). No. 29. Corroborates midden dates.

 $16,400 \pm 250$

UCLA-1099. Pintwater Cave, Nevada

14,450 в.с.

Midden from Pintwater Cave, S Nevada, elev. 1280 m. containing xerophytic semidesert assemblage. Pintwater dates are discussed in Wells and Berger (1967) and provide evidence for pluvial expansion of the pinyon-juniper zone at the close of the Wisconsin glacial to 600 m below present lower limit of woodland. Coll. 18 Dec. 1966.

 $17,750 \pm 200$

UCLA-1218. North Muddy Mountains, Nevada 15,800 B.C.

Midden from 530 m. in N Muddy Mts., 50 km NE of Gipsum Cave. Discussed in same publ. Coll. 17 Dec. 1966.

 12.900 ± 180

UCLA-1219. North Muddy Mountains, Nevada 10,950 B.C.

Midden from 550 m. in same general location as UCLA-1218. Coll. Dec. 1966 and discussed as above.

REFERENCES

Date lists:	
La Jolla I	Hubbs, Bien, and Suess, 1960
UCĽA III	Fergusson and Libby, 1964
UCLA IV	Berger, Fergusson, and Libby, 1965
UCLA V	Berger and Libby, 1966
UCLA VI	Berger and Libby, 1967
UCLA VII	Berger and Libby, 1968
Yale IV	Deevey, Gralenski, and Hoffren, 1959

Berger, R., Fergusson, G. J., and Libby, W. F., 1965, UCLA radiocarbon dates IV: Radiocarbon, v. 7, p. 336-371.

Berger, R., Graham, J. A., and Heizer, R. F., 1967, A reconsideration of the age of the La Venta site: Contributions of the Archaeological Research Facility, No. 3, Univ. of California, Berkeley, p. 1-24.

Berger, R., Horney, A. G., and Libby, W. F., 1964, Radiocarbon dating of bone and shell from their organic components: Science, v. 144, p. 999-1001.

Berger, R. and Libby, W. F., 1966, UCLA radiocarbon dates V: Radiocarbon, v. 8, p. 467-497.

______ 1967, UCLA radiocarbon dates VI: Radiocarbon, v. 9, p. 477-504.

1968, UCLA radiocarbon dates VII: Radiocarbon, v. 10, no. 1, p. 149-160.

Berger, R., Taylor, R. E., and Libby, W. F., 1966, Radiocarbon content of marine shells from California and Mexican west coast: Science, v. 153, p. 864-866.

Deevey, E. S., Gralenski, L. J., and Hoffren, V., 1959, Yale natural radiocarbon measurements IV: Radiocarbon, v. 1, p. 144-172.

Fergusson, G. J. and Libby, W. F., 1962, UCLA radiocarbon dates III: Radiocarbon, v. 6, p. 318-339.

Fletcher, J. M., 1968, Radiocarbon dating of timber-framed cruck cottages: Applications of Science to Medieval Archaeology, in, Contributions of the UCLA Center for Medieval and Renaissance Studies, Univ. of California Press, Berkeley and Los Angeles, in press.

Ho, T., 1965, The amino acid composition of bone and tooth proteins in late Pleistocene mammals: [U.S.] Natl. Acad. Sci. Proc., v. 54, p. 26-31.

Ho, T., Marcus, L. M., and Berger, R., 1968, Radiocarbon dating of bones contaminated by petroleum tar. Manuscript on file.

Horn, W. and Born, E., 1965, The Barns of the Abbey of Beaulieu at its granges of Great Coxwell and Beaulieu St. Leonards: Univ. of California Press, Berkeley and Los Angeles.

Howard, H., 1962, A comparison of avian assemblages from individual pits at Rancho La Brea, California: Contrib. in Science, Los Angeles County Mus., No. 58.

Hubbs, C. L., Bien, G. S., and Suess, H. E., 1960, La Jolla natural radiocarbon measurements I: Radiocarbon, v. 2, p. 197-223.

Kigoshi, K. and Kobayashi, H., Gakushuin natural radiocarbon measurements, to be publ.

Libby, W. F., 1965, Radiocarbon dating: Univ. of Chicago Press, Chicago. First Phoenix Edition.

Martin, L., 1962, Sully-sur-Loire, Des Origines a nos jours: Edition A. Pornin, Sully-sur-Loire, p. 88.

Suess, H. E., 1965, Secular variations of the cosmic-ray-produced carbon 14 in the atmosphere and their interpretations: Jour. Geophys. Research, v. 70, p. 5937-5952.

Taylor, R. E. and Berger, R., 1968, Radiocarbon dating of the organic portion of ceramic and wattle-and-daub house construction materials of low carbon content: Amer. Antiquity, in press.

Wells, P. V. and Berger, R., 1967, Late Pleistocene history of Coniferous woodland in the Mohave desert: Science, v. 155, p. 1640-1647.