

**GEOLOGICAL SURVEY OF FINLAND
RADIOCARBON MEASUREMENTS IV**

E. HYYPPÄ, A. HEIKKINEN, and V. TOIVONEN

C¹⁴ Laboratory, Geological Survey of Finland, Otaniemi, Finland

The following results represent measurements carried out since our 3rd date list was prepared. The entire CO₂ purification technique (Radiocarbon, 1962, v. 4, p. 81-83) has been changed to "wet purification." No measurements of C¹²/C¹³ ratios have been made.

I. GEOLOGIC SAMPLES

Su-34. Kuusamo, E Finnish Lapland **7430 ± 190**
5480 B.C.

Bryales peat taken with piston sampler, depth 4.55 to 4.60 m, surface alt. 272 m, Takasuo bog (65° 56' N Lat, 29° 10' E Long). Coll. 1962 by E. Hyyppä. *Comment*: pollen analysis shows early Atlantic period.

Su-35. Kuusamo, E Finnish Lapland **9320 ± 170**
7370 B.C.

Same bog as Su-34, fine detritus gyttja taken with piston sampler, depth 4.80 to 4.85 m. Coll. 1962 by E. Hyyppä. *Comment*: pollen analysis shows later part of *Betula* maximum time. The C-14 date is possibly too old for this horizon in Lapland.

Su-36. Pello, W Finnish Lapland **8250 ± 150**
6300 B.C.

Fine detritus gyttja taken with piston sampler, depth 5.53 to 5.65 m, surface alt. 101.4 m, Kaakkuririipi bog (66° 47' N Lat, 24° 15' E Long). Coll. 1962 by E. Hyyppä. *Comment*: pollen and diatom analysis indicate early Littorina I stage at beginning of Atlantic period (Hyyppä, 1966).

Su-37. Virolahti, SE Finland **7835 ± 170**
5885 B.C.

Allochthonous peat and coarse detritus gyttja taken with piston sampler, depth 5.05 to 5.12 m, surface alt. 27.11 m, Ruokolampi bog (60° 32' N Lat, 27° 40' E Long). Coll. 1961 by M. Salmi. *Comment*: pollen and diatom analysis indicate beginning of Littorina I stage (Salmi, 1961).

Su-38. Sippola, SE Finland **9710 ± 230**
7760 B.C.

Sphagnum-Carex-Equisetum peat taken with piston sampler, depth 6.55 to 6.65 m, surface alt. 52.42 m, Hangassuo bog (60° 47' N Lat, 26° 55' E Long). Coll. 1948 by M. Salmi. *Comment*: pollen analysis shows Pre-Boreal *Betula* maximum (Salmi, 1948).

Su-39. Kuusamo, E Finnish Lapland **8080 ± 150**
6130 B.C.

Coarse detritus gyttja from drowned peat bog, taken with piston sampler, depth 3.03 to 3.08 m, surface alt. 251.06 m, Joukamojärvi (65°

52° N Lat, 30° E Long). Coll. 1963 by E. Hyyppä. *Comment*: pollen analysis shows *Betula* maximum in beginning of Atlantic period.

8030 ± 200

Su-40. Pello, W Finnish Lapland

6080 B.C.

Terrestrial peat taken with piston sampler, depth 6.0 to 6.1 m, surface alt. 228.76 m, Tornio R. Valley (66° 46' N Lat, 24° 04' E Long). Coll. 1961 by E. Hyyppä. Sample coll. from same profile as Su-24 (Radio-carbon, 1963, v. 5, p. 303) in which surface alt. 288 m was in error. *Comment*: sample lay directly above *Betula* wood dated 6200 B.C.

Su-42. Luumäki, SE Finland

Modern

Wood under sand layer, Käännetsalo Is. (60° 55' N Lat, 27° 30' E Long). Coll. 1965 by E. Hyyppä. *Comment*: wood belongs to very late transgression of Lake Saimaa.

8660 ± 240

Su-43. Virolahti, SE Finland

6710 B.C.

Fine fresh-water mud taken with piston sampler, depth 7.70 to 7.80 m, surface alt. 36.0 m, Huosiosuo bog (60° 33' N Lat, 27° 38' E Long). Coll. 1966 by E. Hyyppä. *Comment*: pollen and diatom analysis indicate Pre-Boreal regression maximum from Yoldia II to Yoldia IV; horizon represents isolation and is overlain by thin layer belonging to Yoldia IV transgression (Hyyppä, 1966).

2560 ± 100

Su-44. Enontekiö, NW Finnish Lapland

610 B.C.

Sorbus aucuparia wood from layer of fine sand and sand, taken with piston sampler, depth 2.5 to 3.0 m, Markkina (68° 30' N Lat, 22° 20' E Long). Coll. 1966 by H. Roivainen. *Comment*: horizon represents beginning of Sub-Atlantic flood period.

4285 ± 130

Su-45. Nurmo, W Finland

2335 B.C.

Sphagnum-Carex peat from hand-dug sec., depth 1.36 to 1.39 m, surface alt. 93.9 m, Kurjenneva bog (62° 46' N Lat, 23° 08' E Long). Coll. 1962 by M. Salmi. *Comment*: pollen analysis shows beginning of *Picea* curve (Salmi, 1963).

6420 ± 130

Su-46. Nurmijärvi, S Finland

2670 B.C.

Carex-Sphagnum peat taken with piston sampler, depth 0.4 to 0.5 m, surface alt. 69 m, Råbacka bog (60° 29' N Lat, 24° 57' E Long). Coll. 1966 by K. Virkkala. *Comment*: sample lay directly under horizon showing beginning of pollen-analytical *Picea* curve (Virkkala, 1966).

5950 ± 150

Su-47. Joensuu, E Finland

4000 B.C.

Betula wood from sand layer, taken with piston sampler, depth 3.15 m, surface alt. 84.68 m, Raatekangas (62° 40' N Lat, 29° 50' E Long).

Coll. 1967 by E. Hyypä. *Comment*: wood belongs to early transgression of Lake Saimaa.

Su-48. Hauho, S Finland

**1250 ± 70
700 A.D.**

Wood from layer of sand, depth 0.3 m (61° 15' N Lat, 24° 40' E Long). Coll. 1966 by E. Hyypä. *Comment*: wood possibly belongs to late transgression of Lake Vanajavesi.

Su-49. Imatra, SE Finland

**8250 ± 200
6300 B.C.**

Bryales peat containing remnants of deciduous trees from under silty mud, depth 11.0 m, surface alt. 75.65 m, Kaukopää (61° 10' N Lat, 28° 50' E Long). Coll. 1966 by E. Hyypä. *Comment*: pollen analysis shows latter half of Boreal period; horizon antedates transgression of Lake Saimaa.

Su-50. Joensuu, E Finland

**5810 ± 150
3860 B.C.**

Sandy peat from under fine sand, Surakka (62° 40' N Lat, 29° 40' E Long). Coll. 1960 by E. Hyypä. *Comment*: sample represents early stage of Lake Saimaa transgression.

Su-51. Asikkala, S Finland

**7550 ± 180
5600 B.C.**

Pinus wood from under stony gravel in Kalkkinen canal (61° 15' N Lat, 25° 30' E Long). Coll. 1963 by H. Peltola and E. Hyypä. *Comment*: wood indicates forest drowned by early Lake Päijänne transgression.

Su-52. Helsinki, S Finland

**4940 ± 140
2990 B.C.**

Alnus wood from hand-dug sec., depth 2.80 m, surface alt. ca. 30 m, Myllypuro (60° 14' N Lat, 25° 07' E Long). Coll. 1965 by E. Hyypä. *Comment*: horizon represents approx. time of Littorina IV, local shore alt. 23 m.

Su-54. Luumäki, SE Finland

**1130 ± 100
820 A.D.**

Carex peat under sand layer, alt. 75.6 m, Käännetsalo Is. (60° 55' N Lat, 27° 30' E Long). Coll. 1965 by E. Hyypä. *Comment*: peat belongs to late transgression of Lake Saimaa.

Porvoo series

Surface and threshold alt. 28.5 m of the Bastuberg peat bog (60° 21' N Lat, 25° 47' E Long). Su-41 and Su-53 coll. 1966, others 1967 by E. Hyypä.

Su-41.

**6870 ± 165
4920 B.C.**

Coarse detritus gyttja, depth 1.75 to 1.82 m. *Comment*: pollen and diatom analysis indicate beginning of Littorina II stage.

- Su-53.** **5560 ± 140**
3610 B.C.
Coarse detritus gyttja, depth 1.27 to 1.33 m. *Comment:* horizon represents time when basin had already isolated from Littorina II stage and Littorina III stage (local shore alt. 26 m) was established.
- Su-55.** **4110 ± 130**
2160 B.C.
Betula bark, depth 0.65 m. *Comment:* pollen analysis shows beginning of *Picea* curve.
- Su-56.** **3910 ± 180**
1960 B.C.
Peat containing remnants of deciduous trees, depth 0.60 m. *Comment:* same as Su-55.
- Su-57.** **5860 ± 200**
3910 B.C.
Coarse detritus gyttja, depth 1.30 to 1.35 m. *Comment:* same as Su-53.
- Su-58.** **5570 ± 200**
3620 B.C.
Phragmites leaves and cones, depth 1.30 m. *Comment:* same as Su-53 and Su-57.
- Su-59.** **6010 ± 105**
4060 B.C.
Phragmites leaves and cones, depth 1.60 m. *Comment:* pollen and diatom analysis indicate maximum of Littorina II stage.
- Su-60.** **7950 ± 190**
6000 B.C.
Wood, depth 2.60 m. *Comment:* horizon represents Littorina I stage directly after maximum stage.
- Su-61.** **8200 ± 270**
6250 B.C.
Pinus wood, depth 2.80 m. *Comment:* horizon represents beginning of maximum transgression of Littorina I stage.
- Su-62.** **6810 ± 165**
4860 B.C.
Wood, depth 1.80 m. *Comment:* horizon represents beginning of maximum transgression of Littorina II stage.
- Su-63.** **7100 ± 170**
5150 B.C.
Wood and *Phragmites* leaves, depth 1.70 to 1.75 m. *Comment:* horizon represents beginning of maximum transgression of Littorina II stage.

Su-64. Helsinki, S Finland

5110 ± 115

3160 B.C.

Same sec. as Su-52, coarse detritus gyttja, depth 3 m. Coll. 1965 by E. Hyypä. *Comment*: diatom analysis shows brackish-water facies; sample belongs to final phase of Littorina III stage.

Su-65. Virolahti, SE Finland

6310 ± 130

4360 B.C.

Fine detritus gyttja taken with piston sampler, depth 4.40 to 4.50 m, surface alt. 19.67 m, Reskovinsuo bog (60° 33' N Lat, 27° 40' E Long). Coll. 1966 by E. Hyypä. *Comment*: horizon represents time subsequent to Littorina II maximum.

Su-66. Ulvila, W Finland

Modern

Branch of *Betula* from sand pit, depth 15 m (61° 24' N Lat, 21° 58' E Long). Coll. 1967 by K. Virkkala. *Comment*: sample is from slumped portion in pit.

II. ARCHAEOLOGIC SAMPLE

Su-67. Sodankylä, Finnish Lapland

Modern

Pinus wood from under boat of ancient Lapp dwelling place, Sompiojärvi (68° 05' N Lat, 27° 27' E Long). Coll. 1964 by E. Hyypä.

REFERENCES

Date lists:

- Finland I Hyypä, Hoffrén, and Isola, 1962
- Finland II Hyypä, Isola, and Hoffrén, 1963
- Finland III Hyypä, Toivonen, and Isola, 1964
- Hyypä, E., 1966, The late-Quaternary land uplift in the Baltic sphere and the relation diagram of the raised and tilted shore levels: Acad. Sci. Fennicae. Annales, A, III, 90, p. 153-168.
- Hyypä, E., Hoffrén, V., and Isola, A., 1962, Geological Survey of Finland radiocarbon measurements I: Radiocarbon, v. 4, p. 81-83.
- Hyypä, E., Isola, A., and Hoffrén, V., 1963, Geological Survey of Finland radiocarbon measurements II: Radiocarbon, v. 5, p. 302-304.
- Hyypä, E., Toivonen, A. V. P., and Isola, A., 1964, Geological Survey of Finland radiocarbon measurements III: Radiocarbon, v. 6, p. 110-111.
- Salmi, Martti, 1948, Die Ancylostrostransgression in dem Moore Hangassuo im Süd-Finnland: Finland, Comm. Géol. B. no. 142, p. 1-20.
- 1961, Two Littorina transgression in Virolahti, southeastern Finland; Finland, Comm. Géol. B. no. 196, p. 417-436.
- 1963, Drei subfossile Sattelrobben aus Ostbotttnien; geologische Datierung der Funde und einige chronologische Beobachtungen: Arch. Soc. "Vanamo", v. 18:2, p. 82-95.
- Virkkala, K., 1966, Radiocarbon ages of the Råbacka bog, southern Finland: Finland, Comm. Géol. B. no. 222, p. 237-240.