

IFAN RADIOCARBON LABORATORY MEASUREMENTS I

Maurice Ndeye^{1,2} • Matar Sène³ • Alpha Oumar Diallo¹

ABSTRACT. Results obtained from a liquid scintillation counter using BGO ($\text{Bi}_4\text{Ge}_3\text{O}_{12}$) tubes have produced more precise radiocarbon dates in our laboratory. Duplicate analyses confirm the electronic stability of the counter with a background of 0.1 cpm. Our ^{14}C dates agree well with those from another laboratory (Paris 6-LOCEAN). Most of the ^{14}C dates in this study were obtained on samples taken from different archaeological sites. Calibration of the various dates with the appropriate software (CALIB 5.0 in our case) allows better interpretation of the results and their importance in this understudied region. In this paper, we investigate the performance of the counter by analyzing samples from archaeological and marine sites in Senegal and Mauritania, and report the results in our first laboratory date list.

INTRODUCTION

Since 2003, we have operated a new dating system in the Institut Francophone d'Afrique Noire (IFAN) Radiocarbon Laboratory in Dakar, Senegal. Counting was carried out using a Tri-Carb[®] 3170TR/SL liquid scintillation analyzer from Packard provided with BGO ($\text{Bi}_4\text{Ge}_3\text{O}_{12}$) tubes driven by the Quanta Smart software under a Windows[®] NT operating system. In “super low level” mode, the background noise is very low (Cook et al. 1990). The liquid scintillation counter is housed in an underground room, thus eliminating more background of atmospheric ^{14}C . Assessment of the performance of the dating system (background and factor of merit) is done using the known samples from Paris 6 and international standards from the International Atomic Energy Agency (IAEA). In the 13–85 kV energy range at an efficiency of 68% and a background of 0.2 cpm, the figure of merit is 23.400.

With the change in methods, the laboratory took on a new lab code, DK, replacing the previous one of DAK (Diop 1974) used in the 1970s by Prof Cheikh Anta Diop (former head of the laboratory). Physical and chemical pretreatments depend on the nature of the samples, but the classic acid-base-acid (ABA) procedure is usually adopted. The acid treatment is done with HCL 8%, and the base with NaOH 5%.

The liquid scintillation counter uses benzene as a scintillation solvent. For each sample, 2 g of benzene was placed in a standard (20 mL) pico glass vial. The scintillators employed were a bis MSB + Buthyl PBD mixture (6 mg + 6 mg) (Horrocks 1974). Age calculations are based on the Libby half-life of 5568 yr and dates are expressed in yr BP with AD 1950 as the reference year. Ages and standard deviations (2σ) of samples are adjusted for stable isotope fractionation to a normalized concentration ratio ($\delta^{13}\text{C} = 25\%$), according to Stuiver and Polach (1977). Calibration was done using the program CALIB 5.0 (Reimer and Reimer 2001; www.calib.org). In the list that follows, new dates are given along with their location coordinates and sample descriptions.

ARCHAEOLOGICAL SAMPLES

Khant Series

Shell samples (*Anadara sensilis*) were collected from the Khant site in northern Senegal (Figure 1). This site is a depression discovered by A Ravisé in 1969 (Ravisé 1970). The samples were submitted 13 January 2003 by H Bocoum of the IFAN laboratory, Université Cheikh Anta Diop de Dakar.

¹Laboratoire carbone 14, IFAN Cheikh Anta Diop BP 206, Dakar, Senegal.

²Corresponding author. Email: mndeye@ucad.sn.

³Institut de Technologie Nucléaire Appliquée (ITNA), Université Cheikh Anta Diop, Dakar, Senegal.




Figure 1 Location of the Khant site, 22 km N-NE of St. Louis (north of the Senegal River delta).

DK-1. Khant, northern Senegal	2448 ± 34 BP 409–598 BC
Bolé zone at 0.3 m depth.	
DK-4. Khant, northern Senegal	2633 ± 29 BP 777–837 BC
Xoor Nook zone at 0.2 m depth.	
DK-5. Khant, northern Senegal	2663 ± 49 BP 775–916 BC
Bolé zone at 2 m depth.	
DK-6. Khant, northern Senegal	2946 ± 86 BP 968–1393 BC
Bolé zone at 1 m depth.	
DK-7. Khant, northern Senegal	2912 ± 29 BP 1010–1212 BC
Bolé zone at 3.2 m depth.	

Comment: These dates (DK-4 and DK-6) could incite new investigations to confirm the initial level of the cluster above 4 m, which could date to the 2nd millennium. This period may be a time of transition that would mark the end of the Neolithic and the introduction of the Iron Age in the valley of the River Senegal (Bocoum et al. 1992). The sequence obtained with DK-1, DK-5, and DK-7 is coherent with regard to the respective calibrations. These results allow us to study the chronological sequence of a small entropic valley in this zone called Bolé.

Samples from IAEA, Vienna

Carbonate sample and cellulose samples were sent from the IAEA. We compare the pMC of the samples given by the IAEA to those calculated in our lab. The aim of this analysis is for calibration of the counter and intercomparison exercises (Rozanski et al. 1990).

DK-2, DK-3. IAEA, Vienna**IAEA = 129.46 pMC****LC14 DK = 129.36 pMC**

Sample of cellulose supplied by the IAEA. It is a series of cellulose (C3) that was produced in 1989 from a harvest during that season.

DK-8. IAEA, Vienna**IAEA = 0.02 pMC****LC14 DK = 0.04 pMC**

Sample of carbonate (C1) supplied by the IAEA.

Comment: The 2 values of pMC are almost the same (difference of 0.08%) and show good calibration with the scintillation counter. Sample C1 is used to measure the background noise of the machine by the low pMC (0.02), which is twice as low as the one calculated by our lab. Taking into account the uncertainty, one can say that the values belong in the same range (Ndeye et al. 2003).

Mauritania Series

Shell samples collected from sites in Mauritania (Figure 2). The samples were submitted in 2003 by Mr Robert Vernet, the head of the Khant Project (SCAC, Ambassade de France, Rabat, Morocco). The aim of the project is to date the consumption of shellfish in order to determine the Nouakchottan transgression.

DK-9. Dakhlet, Nouadibou, Mauritania**6150 ± 86 BP****4894–5305 BC**

Shell samples (*Anadara senilis*) collected in March 2003 from the Dakhlet site on the surface of the shell midden (20°51'N, 17°02'W).

DK-10. Dhraïna, Mauritania**5142 ± 49 BP****3966–4175 BC**

Shell samples (*A. senilis*) collected in May 2003 from the Dhraïna site in Mauritania at the shell midden (18°48'N, 15°29'W).

DK-11. Grandes dunes Nouakchott, Mauritania**6000 ± 115 BP****4651–5215 BC**

Shell samples (*Lopha stentina*) collected in May 2003 from site 32 in Mauritania.

DK-12. Targueuit, Keur massène, Rosso, Mauritania**5030 ± 80 BP****3660–3967 BC**

Shell samples (*A. senilis*) collected 29 January 2004 from the Bantar After Saheli site (Rosso in Mauritania).

DK-13. Akchar, Mauritania**6090 ± 85 BP****4791–5224 BC**

Shell samples (*L. stentina*) collected 30 January 2004 from site 28 of Bantar After Saheli (Akchar in Mauritania).




Figure 2 Senegal and Mauritania coastal sites

DK-14. Akchar, Mauritania

**4340 ± 40 BP
2890–3029 BC**

Shell samples (*Donax rugosus*) collected 31 May 1996 from the site of Bantar After Saheli (Akchar in Mauritania). Samples collected from the shell midden situated at 18°59'N, 16°15'W.

DK-15. Goula (Aftout es Saheli), Mauritania

**6485 ± 64 BP
5321–5557 BC**

Shell samples (*A. senilis*) collected 31 May 1996 from the site of Goula in Mauritania. Samples collected from the shell midden situated at 16°55'N, 16°14'W.

DK-16. Foum Arguin, Mauritania

**5147 ± 76 BP
3763–4079 BC**

Shell samples (*A. senilis*) collected 31 May 1996 from the site of Foum Arguin in Mauritania.

Comment: The sequence DK-9 to DK-16 are coherent with regards to their respective calibrations and allow the determination of the Nouakchottan transgression (Vernet 1993) climatic phase.

Ngayene Series

Samples of charcoal from wood collected and presented by A Holl from the Museum of Anthropology at the University of Michigan (USA) during archaeological excavations made at the Megalithic sites of the rural community of Ngayene in the Falmé region (northern Senegal) in June 2005.

DK-17. Ngayene, Nioro du Rip, central-west Senegal **305 ± 35 BP**
AD 1482–1654

Sample collected at 1.6–1.7 m depth.

DK-18. Ngayene, Nioro du Rip, central-west, Senegal **3075 ± 30 BP**
1268–1415 BC

Sample collected at 1.7 m depth.

DK-19. Ngayene, Nioro du Rip, central-west Senegal **820 ± 25 BP**
AD 1173–1264

Sample collected at 0.6 m depth.

DK-28. Ngayene, Nioro du Rip, central-west Senegal **926 ± 32 BP**
AD 1025–1180

Sample collected at 20–40 m depth.

DK-29. Ngayene, Nioro du Rip, central-west Senegal **960 ± 27 BP**
AD 1064–1155

Sample collected at 61–80 m depth.

DK-32. Ngayene, Nioro du Rip, central-west Senegal **3823 ± 60 BP**
1075–1155 BC

Sample collected at 190–200 cm depth.

Comment: The purpose of this sequence of dates (DK-17, -18, -19, -28, -29, and -32) is the construction of a chronological timeframe of the Megalithic cemetery of Ngayene. These dates were made in a cooperative study between the ¹⁴C laboratory of IFAN (Dakar) and the LOCEAN laboratory (University Paris VI, France).

Grande Côte Series

These samples were collected by Hamady Bocoum, archaeologist at the Laboratory of Protohistory and Prehistory of the IFAN, and Cheikh Anta Diop of Dakar. Dates below are made at the Grande côte, Senegal (Figure 3) for small stations behind the offshore bar of the big coast, which had not been studied prior.

DK-20. Grande côte, Senegal **3657 ± 81 BP**
1871–2235 BC

Charcoal from wood collected at 1.5 m depth, 1600 m north of Mbanar 1 (18°48'N, 15°29'W).

DK-21. Grande côte, Senegal **2005 ± 50 BP**
121 BC–AD 85

Charcoal from wood collected between 190 and 200 cm depth, 1600 m north of Mbanar 1.

DK-22. Grande côte, Senegal **2014 ± 75 BP**
206 BC–AD 137

Charcoal from wood collected at 1.3 m depth at Mbanar 1. The tumulus is located near the village school (15°15'N, 16°20'W).

DK-23. Grande côte, Senegal **1970 ± 35 BP**
45 BC–AD 87

Charcoal from wood collected at 1.55 m depth, 1600 m north of Mbanar 1. The tumulus is situated near the village school (15°15'N, 16°20'W).

DK-24. Grande côte, Senegal **1643 ± 49 BP**
AD 322–539

Charcoal from wood collected at the village of Keur Abdou Ndoeye, Kayar (15°15'N, 16°20'W).

DK-25. Grande côte, Senegal **5900 ± 66 BP**
4601–4944 BC

Shell samples collected at an ocean station connecting Kayar and Mboro.

DK-25a. Grande côte, Senegal **1324 ± 70 BP**
AD 602–878

Shell samples collected on the road leading to Lac Rose.

DK-25b. Grande côte, Senegal **3823 ± 60 BP**
2133–2467 BC

Shell samples collected at Yallay-Mbanar-Mbow site VII.

Comment: Sequence DK-20, DK-21: Analysis of the progress of the excavation allows us to construct a relatively coherent succession of the early sequences of the tumulus. Two dates obtained at approximately the same depth are very close (1970 ± 35 and 2014 ± 37 BP) and place us at the beginning of the Christian era.

GEOLOGICAL SAMPLES

The following samples were collected in eastern Senegal by Alain Pearson, researcher at the University Paris VI (France).

DK-30. East Senegal **5033 ± 33 BP**
3760–3947 BC

Concretion samples of lakeside limestone collected in Ferlo, a site in eastern Senegal (15°27'97"N, 13°39'32"W).

DK-31. East Senegal **4517 ± 47 BP**
3089–3365 BC

Pedological carbonated concretion samples brought out by the dismantling of a wind turbine in Falémé (14°22'78"N, 12°13'99"W).

MARINE SAMPLES

The following samples were analyzed to determine $\Delta^{14}\text{C}$ (Stuiver and Polach 1977; Mook and van der Plicht 1999) and time variation for the Senegalese coastal region. The samples were collected by Mr Marchad and processed by the Marine Biology Laboratory of IFAN.

DK-26, DK-27. Dakar (Senegal) $\Delta^{14}\text{C} = 44\text{‰}$
Shell sample collected in 1951.

DK-33. Dakar (Senegal) $\Delta^{14}\text{C} = 43\text{‰}$
Shell sample collected in 1951.

DK-34. Dakar (Senegal) $\Delta^{14}\text{C} = 45\text{‰}$
Shell sample collected in 1956.

DK-35. Iles du saloum (Senegal) $\Delta^{14}\text{C} = 44\text{‰}$
Coral sample collected in 1950.

DK-36. Dakar, Senegal $\Delta^{14}\text{C} = 65\text{‰}$

Coral sample collected in 1974.

Comment: These ^{14}C values correspond with the predicted values according to their collection dates (Broecker et al. 1978).

ACKNOWLEDGMENTS

We thank all the researchers and technicians who contributed to the production of these dates and who allowed their publication. Particularly, our thanks go to Dr Hamady Bocoum of the IFAN Laboratory of Archaeology at Cheikh Anta Diop University, and Jean-François Saliège from the Laboratory of Oceanography at the University Paris 6 (France) for his constant support and cooperation.

REFERENCES

- Bocoum H, McIntosh SK, McIntosh RJ. 1992. L'âge du fer dans la moyenne vallée du fleuve: chronologie et integration. Communication présentée au cinquième colloque de l'association ouest africaine d'archéologie, Ouagadougou, 27 July–1 August 1992.
- Broecker WS, Peng T-H, Stuiver M. 1978. An estimate of the upwelling rate in the equatorial Atlantic based on the distribution of bomb radiocarbon. *Journal of Geophysical Research* 83(C12):6179–86.
- Cook GT, Hold AG, Naysmith P, Anderson R. 1990. Applicability of 'new technology' scintillation counters (Packard 2000 CA/LL and 2260 XL) for ^{14}C dating. *Radiocarbon* 32(2):233–4.
- Diop CA. 1974. Physique nucléaire et chronologie absolue. Initiations et Etudes Africaines XXXI. Université de Dakar–IFAN.
- Horrocks DL. 1974. *Applications of Liquid Scintillation Counting*. New York: Academic Press.
- Mook WG, van der Plicht J. 1999. Reporting ^{14}C activities and concentrations. *Radiocarbon* 41(3):227–39.
- Ndeye M, Ka O, Bocoum H, Diallo AO. 2003. Rehabilitation of the Laboratoire de carbone 14-Dakar (Senegal) with a super low-level liquid scintillation counting system. *Radiocarbon* 46(1):117–22.
- Ravisé A. 1970. Industries en os de la région de St-louis. *Notes africaines* 128:98–102.
- Rozanski K, Stichler W, Gonfiatini R, Scott EM, Beukens RP, Kromer B, van der Plicht J. 1990. The IAEA ^{14}C intercomparison exercise 1990. *Radiocarbon* 34(3):506–19.
- Reimer PJ, Reimer RW. 2001. A marine reservoir correction database and on-line interface. *Radiocarbon* 43(2A):461–3.
- Stuiver M, Polach HA. 1977. Discussion: reporting of ^{14}C data. *Radiocarbon* 19(3):355–63.
- Vernet P. 1993. *Préhistoire de la Mauritanie*. CCF/Sepia. Paris: Nouakchott.