MOSCOW MV LOMONOSOV STATE UNIVERSITY RADIOCARBON DATES II SEA LEVEL INDICATORS FROM COASTAL USSR

N I GLUSHANKOVA, O B PARUNIN, A O SELIVANOV, A I SHLUKOV, and T A TIMASHKOVA

Laboratory of Recent Sediments and Pleistocene Paleogeography, Moscow State University, Moscow 117234, USSR

INTRODUCTION

The following list summarizes the post-1970 laboratory results of ¹⁴C dating of ancient sea-level indicators from the coasts of the Soviet Union. One of the aims of the International Geologic Correlation Programme Project No. 61 "Sea level movements during the last deglacial hemicycle" is the global cataloguing and mapping of ancient sea levels. The laboratory, which acts as the USSR National curator for these age measurements obtains dates sampled from its own expeditions and from other institutions of the country.

The list of ¹⁴C dated sea-level indicators from the Soviet coasts was sent to Walter S Newman, Queens College, City University of New York, at his request for use in an enlarged world catalogue and maps.

Global mapping and modeling require reference to the laboratory code (MGU) number¹, latitude-longitude position, present elevation, date, and material of each sea-level indicator. A brief site description is published below. Some of the dated sea levels are evidently tectonically deformed but that is not our concern here. For samples younger than 7000 ¹⁴C years, we adduce not only the direct age measurements but also the corrected dates in parentheses. We have chosen for correction the dendrochronologic calibration curve of Michael and Ralph (1973).

¹⁴C dating was performed by the liquid scintillation method. To exclude from dating recrystallized mollusk shells, we use an x-ray diffractometric method. For a brief description of measurement procedures and methodologic aspects, see Glushankova *et al* (1980) and Kaplin (1976).

ACKNOWLEDGMENTS

We thank P A Kaplin, the Head of the Laboratory, the Head of the National working group on Project No. 61 IGCP, for advice in our research. We are also greatly indebted to all of those who submitted samples for dating.

Arctic Ocean Coasts

MGU-IOAN-129. Spitzbergen Is

 8360 ± 140

Wood from ancient driftwood trunk, NE Barentz I. (78° 6′ N, 22° 2′ E), from scarp of limnodeltaic terrace, at 48m alt. Sample subm by M G Groswald, Inst Geog, Acad Sci, USSR.

¹ Dates obtained jointly with the Inst Oceanology, Acad Sci, USSR, are lettered MGU-IOAN.

Laptev Sea series

Samples subm by A Sidorchuk, Moscow State Univ.

 840 ± 250

MGU-326. Yana River delta

 (880 ± 250)

Wood from main channel, left bank, 84km from mouth (71° 1′ N, 136° 1′ E). Sample from marine terrace at 8m alt.

 1970 ± 150

MGU-327. Yana River delta

 (1870 ± 150)

Wood from main channel, left bank, 32km from mouth $(71^{\circ} 4' N, 135^{\circ} 8' E)$. Sample from marine terrace at 6m alt.

 3640 ± 300

MGU-329. Yana River delta

 (3850 ± 300)

Peat from main channel, right bank, 13km from mouth (71° 5′ N, 136° 2′ E). Sample from marine terrace at 2m alt.

Longa Strait series

Samples subm by L V Tarakanov, Moscow State Univ.

 1370 ± 140

MGU-574. Billings Cape

 (1459 ± 140)

Peat from N lagoon coast, 0.9km W of Billings settlement (69° 8′ N, 176° 1′ E). Sample from ledge of bar at 2.5m alt.

 3007 ± 170

MGU-575. Billings Cape

 (3116 ± 170)

Peat from N lagoon coast, 2.7km E of Billings settlement (69° 8′ N, 176° 2′ E). Sample from ledge of bar at 2.1m alt.

 3650 ± 170

MGU-576. Billings Cape

 (3850 ± 170)

Peat from N lagoon coast, 7.2km E of Billings settlement (69° 8′ N, 176° 0′ E). Sample from ledge of bar at 1.8m alt.

Taimir series

Samples subm by M G Groswald.

MGU-450. Taimir Lowland

 $25,000 \pm 250$

Mollusk shells, 100% aragonite (Mya truncata, Cytrodaria jenissea, Hiatella arctica, Astarte montagui striata) from Ulakhan-Gurakh, right bank, 6km from Novaya R confluence, 45km NNW of Khatanga settlement (72° 3′ N, 102° 3′ E). Sample from marine terrace at 50 to 55m alt.

MGU-451. Taimir Lowland

 $30,000 \pm 300$

Mollusk shells (Mya truncata, Hiatella arctica, Astarte montagui striata, Macoma calcarea, Cyrtodaria sp) from Selebir R, upper reaches, 40km NW of Khatanga settlement (72° 2′ N, 102° 0′ E). Sample from marine terrace at 60m alt.

MGU-452. Taimir Lowland

 $15,300 \pm 500$

Mollusk shells (*Mya* sp, *Astarte* sp, *Macoma* sp, *Hiatella* sp) from Ushkan Kamen Heights, Dudypta R, left bank, 90km from mouth of Boganida R at 120m alt (71° 8′ N, 93° 2′ E).

Far East Coasts

Western Kamchatka series

Samples coll by lab expedition; subm by V F Ivanov and L G Nikiforov.

MGU-IOAN-68. W Kamchatka

 35.000 ± 700

Wood from 5.5km N of Kihchik R mouth, 350m from lake shore (53° 6′ N, 156° 0′ E). Sample from scarp of terrace at 1m alt.

MGU-IOAN-185. W Kamchatka

 $37,000 \pm 2100$

Buried peat from Utka R near Mitoga settlement (53° 0′ N, 156° 0′ E). Sample from scarp of terrace at 25 to 30m alt.

MGU-60. W Kamchatka

 $31,000 \pm 900$

Wood from peaty layer from modern sea cliff in marine terrace at 6.2m alt (52° 9′ N, 156° 0′ E).

MGU-202. W Kamchatka

45,000

Wood from same loc as MGU-IOAN-185 at 12.3m alt.

Chukotka series

Samples coll by lab expedition; subm by A A Svitoch and V S Khorev.

MGU-201. Chukotka

 $27,000 \pm 2000$

Plant remains from Anadyr estuary, Dionisiy cape (64° 7′ N, 177° 3′ E). Sample from scarp of terrace at 23m alt.

MGU-311. Chukotka

 $31,500 \pm 850$

Peat from Osinovaya R, right bank, 2km from mouth (64° 7′ N, 175° 0′ E). Sample from scarp of plane surface at 21.5m alt.

MGU-312. Chukotka

44,000

Peat from same loc as MGU-311.

MGU-314. Chukotka

 $37,500 \pm 800$

Wood from SW coast of Anadyr estuary, 1.5km N of Khluzny stream (64° 6′ N, 177° 3′ E). Sample from scarp of marine terrace at 7m alt.

 7060 ± 200

MGU-320. Chukotka

 (7740 ± 200)

Peat from Anadyr estuary coast, 2km N of fishing factory (64° 6′ N, 177° 3′ E). Sample from scarp of 1st marine terrace at 2.2m alt.

 7010 ± 160

MGU-321. Chukotka

 (7680 ± 160)

Plant remains from same loc as MGU-320 at 2.3m alt.

Moscow MV Lomonosov State University Radiocarbon Dates II 895

MGU-340. Chukotka

 $11,800 \pm 200$

Peat from Onemen Bay coast (64° 5′ N, 176° 8′ E). Sample from scarp of marine terrace at 4.8m alt.

MGU-341. Chukotka

 8950 ± 200

Peat from same loc as MGU-340 at 2.9m alt.

MGU-342. Chukotka

 8200 ± 100

Peat and plant remains from same loc as MGU-320 (64° 6′ N, 177° 3′ E) at 3.9m alt.

MGU-383. Chukotka

 $11,600 \pm 100$

Peat from Koluchinskaya inlet, rear part of Belyanka spit (67° 1' N, 174° 7' W) at 0.5m alt. Subm by L A Zhindarev, MGU.

Kresta Bay series

MGU-384. Kresta Bay

 $30,000 \pm 1100$

Mollusk shell, 100% aragonite from Kresta (Cross) Bay, E coast, S of Perkla stream mouth (65° 7′ N, 178° 7′ W). Sample from scarp of marine terrace at 10.5m alt.

MGU-385. Kresta Bay

 $30,000 \pm 380$

Mollusk shell, 100% aragonite from Kresta Bay, E coast, 8km N of Konergino settlement (66° 1′ N, 178° 6′ W). Sample from slope of watershed plain at 7.2m alt.

MGU-386. Kresta Bay

 $35,000 \pm 530$

Mollusk shell, 100% aragonite from same loc as MGU-384 (65° 6′ N, 178° 7′ W) at 17.5m alt.

 5900 ± 110

MGU-393. Kresta Bay

 (6480 ± 110)

Peat from same loc as MGU-320 (64° 6′ N, 177° 3′ E) at 1.6m alt.

MGU-394. Kresta Bay

 7700 ± 320

Plant remains from same loc as MGU-393 at 2.4m alt.

 6120 ± 220

MGU-398. Kresta Bay

 (6770 ± 220)

Peat from same loc as MGU-394 at 1.6m alt.

Dionisiya River Mouth series

 3150 ± 80

MGU-402. Dionisiya R

 (3300 ± 80)

Plant remains from excavation of ancient spit at 2.8m alt, 1.5km N of mouth (64° 7′ N, 177° 3′ E).

 4180 ± 100

MGU-403. Dionisiya R

 (4400 ± 100)

Plant remains from same loc as MGU-402 at 2m alt.

 4000 ± 100

MGU-454. Dionisiya R

 (4360 ± 100)

Peat from recent marine terrace at 0.5 to 1m alt (64° 7′ N, 177° 3′ E).

MGU-478. Dionisiya R

 9900 ± 500

Peat from crest of ancient spit (64° 6′ N, 177° 3′ E) at 4.5m alt.

 4200 ± 120

MGU-455. Dionisiya R

 (4500 ± 120)

Peat from same loc as MGU-454 at 0.6m alt.

Japan Sea series

Samples coll by lab expedition and A M Korotky, Vladivostok, Geol Inst, Far East Sci Center, Acad Sci, USSR.

MGU-IOAN-227. Vostok Gulf

11,000

Peat from 16m depth at 1.7 to 2m from basal surface (42° 8′ N, 132° 7′ E). Subm by A M Korotky.

MGU-IOAN-229. Barabashevka R mouth

 7360 ± 160

Peat from lagoonal deposits in scarp of terrace at 0.2m alt (42° 9′ N, 131° 4′ E). Subm by A M Korotky.

 6260 ± 110

MGU-263. Partizanskaya R mouth

 (6980 ± 110)

Mollusk shells from marine terrace (42° 8′ N, 133° 0′ E). Sample from well at -9m alt; subm by A I Shlukov, MGU.

 5790 ± 110

MGU-264. Partizanskaya R mouth

 (6370 ± 110)

Mollusk shells from same loc as MGU-263 at -6m alt.

MGU-305. Kievka Bay

38,000

Peat from Chinchasovaya inlet, 2nd submerged terrace, 30m depth, 1.5km from coast at 0.15 to 3.4m from basal surface (42° 9′ N, 134° 0′ E).

MGU-307. Vostok Gulf

42.00

Peaty silt from 2nd submerged terrace, 26m depth, 0.5 to 1m from basal surface (42° 8′ N, 132° 7′ E).

 2880 ± 200

MGU-323. Chernaya R mouth

 (2950 ± 200)

Peat from scarp of lagoonal terrace at 3m alt (43° 1′ N, 134° 3′ E). Subm by A M Korotky.

 4450 ± 100

MGU-324. Chernaya R mouth

 (4890 ± 100)

Peat from same loc as MGU-323 at 2.1m alt.

MGU-325. Zolotoy Rog Gulf (Golden Gulf) $29,000 \pm 250$

Algae peat from 7.3m depth, 6.5 to 6.8m from basal surface (43° 1' N, 131° 3' E).

MGU-408. Zerkalnaya Bay

 2950 ± 60

Mollusk shells from 20m depth, 3.2 to 3.3m from basal surface (44° 1' N, 135° 8' E).

 3300 ± 100

MGU-409. Rudnaya Bay

 (3490 ± 100)

Mollusk shells from 20m depth, 0.91 to 1.4m from basal surface (44° 2′ N, 135° 9′ E).

 3950 ± 800

MGU-461. Zerkalnaya R mouth

 (4320 ± 800)

Coal overlying marine pebble from archaeol site on scarp of terrace at 1.8m alt (44° 0′ N, 135° 6′ E). Subm by V P Stepanov, MGU.

 5700 ± 80

MGU-501. Zerkalnaya R mouth

 (6300 ± 80)

Wood from scarp of low flood plain at 1.5m alt (44° 0′ N, 135° 6′ E). Subm by S S Karpukhin, MGU.

MGU-515. Zerkalnaya Bay

 $10,680 \pm 260$

Silt for analysis of humic acid fraction, 35m depth, in submerged terrace, 4 to 3.8m from basal surface (44° 1′ N, 135° 8′ E). Subm by A I Vvedenskaya, MGU.

MGU-516. Rudnaya Bay

 $12,190 \pm 700$

Silt for analysis of humic acid fraction, 20m depth, in submerged terrace, 4.1 to 4.5m from basal surface (44° 2′ N, 135° 9′ E). Subm by A I Vvedenskaya.

MGU-518. Rudnaya Bay

 $10,300 \pm 260$

Silt for analysis of humic acid fraction, 33m depth, in submerged terrace, 3.3 to 3.4m from basal surface (44° 2′ N, 135° 9′ E). Subm by A I Vvedenskaya.

 1227 ± 120

MGU-538. Kema R mouth

 (1270 ± 120)

Wood from scarp of terrace at 1.8m alt, 0.6km from coastline (45° 4′ N, 137° 3′ E). Subm by A M Korotky.

 1270 ± 30

MGU-539. Kema R mouth

 (1320 ± 30)

Wood from same loc as MGU-538 at 1.1m alt.

 2270 ± 120

MGU-540. Kema R mouth

 (2851 ± 120)

Wood from 2.5km S of mouth from scarp of marine terrace at 3.7m alt. Subm by A M Korotky.

 4320 ± 90

MGU-544. Valentin Bay

 (4670 ± 90)

Coal overlying marine pebble from archaeol site at Titova tombolo at 1.8m alt. Subm by V P Stepanov, MGU.

MGU-546. Kema R

 2870 ± 50

 (2940 ± 50)

Peat from Khomushina stream mouth from scarp of lagoonal terrace at 1.2m alt (45° 3′ N, 137° 3′ E). Subm by A M Korotky.

 2500 ± 170

MGU-559. Adimi Bay

 (2538 ± 170)

Wood from 2.5km S of Adimi R mouth from scarp of marine terrace at 2.2m alt (47° 2′ N, 139° 0′ E). Subm by A M Korotky.

 $\mathbf{2910} \pm \mathbf{600}$

MGU-608. Rudnaya Bay

 (3000 ± 600)

Silt lens in sand deposits from ancient spit at 1.7m alt, 1km S of Rudnaya R mouth (44° 1′ N, 135° 8′ E). Subm by A I Shlukov, MGU.

MGU-638. Peter the Great Gulf

28,000

Peaty silt from 22m depth of submerged terrace, 3.3m from basal surface (42° 6′ N, 131° 3′ E). Subm by A V Porotov, MGU.

REFERENCES

Glushankova, N I, Parunin, O B, Timashkova, T A, Khait, V Z, and Shlukov, A I, 1980, Moscow MV Lomonosov State University radiocarbon dates I: Radiocarbon, v 22, p 82-90.

Kaplin, P A, ed, 1976, Rukovodstvo po Izucheniu Novejshih Otlozhenij M, Moscow Univ Pub House (Manual on recent sediments research).

Michael, H N and Ralph, E K, 1973, Discussion of radiocarbon dates obtained from precisely dated sequoia and bristlecone pine samples, in Rafter, T H and Grant-Taylor, T, eds, Internatl conf on ¹⁴C dating, 8th, Proc: Wellington, Royal Soc New Zealand, p 27-43.